

## **Pediatric Products**

Handbook

Leading the way in science-based pediatric nutrition products, to help give infants and toddlers the best start in life.



Hospital Resource Centre: 1-800-263-1143
Order Desk: 1-877-307-2984





## **Pediatric Products Handbook**



For more than 100 years, Mead Johnson has been providing science-based infant formulas and other nutritional products developed to help people lead healthier lives. Our company was founded by Edward Mead Johnson, whose son, Ted, suffered from severe feeding difficulties and barely survived infancy. The experience helped inspire E. Mead to develop products that would provide nutritional excellence to healthy infants as well as those with special dietary requirements. The same principles that guided the company during its early days are still the

Our Vision is to be the world's leading nutrition company for babies and children. We are dedicated to nourishing the world's children for the best start in life.

basis for its vision today:

IMPORTANT CONT	ACT INFORMATION
MJN Representati	ve
Phone Number	
HEALTHCARE PRO	DFESSIONAL CONTACT INFORMATION
1-800-263-1143	Hospital Resource Centre
1-877-307-2984	Order Desk

Intended for Canadian Healthcare Professionals Only. ©2024 Mead Johnson & Company, LLC

enfamil.ca/medical

The information in this handbook is provided as a service to healthcare professionals. Every measure is taken to assure that this is the most current and accurate information available. Because our products are subject to revision, we ask that you please refer to the product label for the most accurate information.





## **Table of Contents**

#### PREMATURE PRODUCTS

Infamil	Enlamil
Spirit in the lateral	Human Milk Forder Fordiert de let maternel
	-
	Marian.

Enfamil® Human Milk Fortifier12
Powdered breast milk supplement for premature
or low birth weight infants.



## Enfamil® Liquid Human Milk Fortifier Standard Protein, 5 mL Sachet ..........16 Please contact your Hospital Account Manager for



Enfamil® Liquid Human Milk Fortifier
High Protein, 5 mL Sachet ......26

Please contact your Hospital Account Manager for product access and contract inclusion.

product access and contract inclusion.





Enfamil® Liquid Human Milk
Fortifier High Protein ......39



#### Enfamil A<sup>+®</sup> Premature High Protein Nursette<sup>®</sup> Bottle .....43

Cow's milk-based, iron-fortified infant formula specifically designed for premature or low birth weight infants that require a higher level of protein.





**Enfamil A<sup>+®</sup> NeuroPro<sup>™</sup> Gentlease<sup>®</sup> ......78**Partially hydrolyzed formula designed to reduce fussiness, crying, and gas.

Control of the contro	Enfamil A <sup>+®</sup> For feeding babies who frequently Spit Up <sup>‡</sup>
	<b>Enfamil A<sup>+®</sup> Lactose Free</b>
	<b>Enfamil A<sup>+®</sup> Soy95</b> Soy-based, iron fortified lactose-free infant formula designed for infants who need a milk-free formula due to health or cultural reasons.
Enfamil	Enfamil A+® Serenity
Nutramigen  Coc  Megnadugan, 199-darphopan  Megn	Nutramigen® A+® with LGG®
Nutramigen  Papolinguis 1 Spaningsing  Papolinguis 2 Spaningsing  Papolinguis 3 Spaningsing  Papolinguis 2 Spaningsing  Papolinguis 3 Spaningsing  Papolingu	Nutramigen® A+®
PURAMINO  Pyckelryic  Ryspelryic  Was de Sale  Was and was de Sale  Was	PURAMINO A <sup>+®</sup> Hypoallergenic, 100% free amino acid-based, iron fortified infant formula for the dietary management of infants and toddlers with severe cow's milk protein and multiple food allergies or other conditions requiring an amino acid-based formulation.



5% alucose in water. For oral use only.

#### **TODDLER PRODUCTS**



#### **MEDICAL PRODUCTS**



Portagen®																	•	•								1	4	8
1 milk-protein	h	2	c	2	1 -	20	١١٨	10	ما	r	۱۸	/it	·h	r	n	۵,	4;	 m	١_	را	٠.	si.	n					

A milk-protein based powder with medium-chain triglycerides (MCT) for children and adults.

Light cherry flavour.



## Enfamil® Enfalyte® Oral Electrolyte Solution ................152 Oral electrolyte maintenance solution.

#### **METABOLICS**



## Phenyl-Free® 1 ......156

Phenylalanine-free, iron fortified infant formula with DHA and ARA for infants and young children with documented phenylketonuria (PKU).



### Phenyl-Free® 2 ......160

Phenylalanine-free powder for dietary management of children and adults (3 years and older) with phenylketonuria (PKU).



### Phenyl-Free® 2 HP ......164

High protein phenylalanine-free powder for dietary management of children and adults (3 years and older) with phenylketonuria (PKU).



#### TYROS1 ......168

Tyrosine-and phenylalanine-free, iron fortified infant formula with DHA and ARA for infants and young children with documented tyrosinemia type I and type II.



### 

Phenylalanine and tyrosine-free powder for dietary management of children and adults (3 years and older) with inborn errors of tyrosine metabolism including tyrosinemia type II.



(MSUD).

amino acid metabolism

metabolism.

An isoleucine-, leucine-, and valine-free, iron fortified infant formula with DHA and ARA for the dietary management of infants and young children with documented Maple Syrup Urine Disease





\* This product does contain taurine, a non-protein building amino acid.



Protein- and amino acid-free powder\* for dietary management of children and adults (3 years and older) with documented amino acid metabolic disorders.

\* This product does contain taurine, a non-protein building amino acid.

#### **VITAMINS**



Enfamil® D-Vi-Sol® (Liquid) ......193
Vitamin D Supplement

Liquid vitamin D supplement for all full-term breastfed infants.



Enfamil® Tri-Vi-Sol® (Liquid) .............195
Multi-Vitamin Supplement
of Vitamins A. D and C

Liquid vitamin A, D, and C supplement for infants and children under 3 years of age. For the maintenance of good health for growing infants and toddlers.



Concentrated Liquid Infant Formula Dilutions ......223

Kosher and Halal Products238
<b>Hospital Instructions</b>
Miscellaneous Nutrient Information246
ADDITIONAL INFORMATION
Alberta Reimbursement List248
Ontario Reimbursement List249
Quebec Reimbursement List250
Saskatchewan Reimbursement List251



## **Premature Products**



Date published 11/2024 – Please reference Mead Johnson Nutrition's online Product Handbook for the most up to date product details (www.enfamil.ca/medical).



## Enfamil® Human Milk Fortifier (Powder)

Powdered breast milk supplement for premature or low birth weight infants.

OD (M) IFANCA

#### INDICATION

Powdered breast milk supplement for premature or low birth weight infants. Four (4) packets of Enfamil® Human Milk Fortifier, the amount usually added to 100 mL preterm human milk, supply the following:

#### **NUTRIENTS**

Analysis	per 4 packets				
Energy (kcal/kJ)	14/59				
Protein (g)	1.1				
% of total energy	32				
Source: Milk protein isolate, whey protein isolate hydr	olysate				
Carbohydrate (g)	0.4				
% of total energy	6				
Source: Minerals, corn syrup solids, citrates, lactose					
Fat (g)	1				
% of total energy	62				
Source: MCT oil (70%), soy oil (30%)					
Linoleic Acid (mg)	140				
Linolenic Acid (mg)	17				
Minerals					
Calcium (mg) (mMol)	90 2.2				
Phosphorus (mg) (mMol)	50 1.61				
Magnesium (mg) (mMol)	1 0.04				



Analysis	per 4 packets
Iron (mg)	1.44
Zinc (mg)	0.72
Manganese (μg)	10
Copper (µg)	44
Sodium (mg) (mMol)	16 0.7
Potassium (mg) (mMol)	29 0.74
Chloride (mg) (mMol)	13 0.37
Vitamins	
Vitamin A (IU)	950
Vitamin D (IU)	150
Vitamin E (IU)	4.6
Vitamin K (μg)	4.4
Vitamin C (mg)	12
Thiamine (µg)	150
Riboflavin (µg)	220
Niacin (µg)	3000
Pantothenic Acid (µg)	730
Vitamin B <sub>6</sub> (μg)	115
Folic Acid (µg)	25
Vitamin B <sub>12</sub> (μg)	0.18
Biotin (µg)	2.7

Other Characteristics							
Potential Renal Solute Load (m0sm/4 packets) <sup>1</sup>	9.8*						
Osmolality (mOsm/kg H <sub>2</sub> O)	+ 35†						
Osmolarity (mOsm/L)	Not available						

<sup>\*</sup> When added to preterm human milk as recommended (4 packets/100 mL), Enfamil® Human Milk Fortifier increases potential renal solute load by approximately 9.8 mOsm/4 packets.

<sup>†</sup> When added to preterm human milk as recommended (4 packets/100 mL), Enfamil® Human Milk Fortifier increases osmolality by approximately 35 m0sm/kg water.

<sup>1.</sup> Fomon SJ, Ziegler EE. Renal solute load and potential renal solute load in infancy. *J Pediatr.* 1999;134:11-14. Nutrient values are subject to change. Consult actual product labels for most current information.

#### **INGREDIENTS**

Medium chain triglycerides (MCT oil), Modified milk ingredients (cow's milk), Hydrolyzed whey, Soy oil, Corn syrup solids, Soy lecithin, Citric acid, **Minerals** (calcium gluconate, calcium glycerophosphate, calcium phosphate, cupric sulfate, ferrous sulfate, magnesium phosphate, potassium chloride, potassium citrate, potassium phosphate, sodium citrate, zinc sulfate), **Vitamins** (ascorbic acid, biotin, calcium D-pantothenate, dl- $\alpha$ -tocopheryl acetate, folic acid, niacinamide, pyridoxine hydrochloride, riboflavin, sodium ascorbate, thiamine hydrochloride, vitamin A palmitate, vitamin B<sub>12</sub>, vitamin D<sub>3</sub>, vitamin K<sub>1</sub>).

#### PRODUCT FORMS

Product	Format	Unit Size	Shelf Life	ltem Number	Availability
Enfamil® Human Milk Fortifier	Powder	Four (4) packets 0.71g per packet	12 months	3221214	Hospital & available on discharge by hospital dispensing

#### PREPARATION OF FEEDINGS

Enfamil® Human Milk Fortifier is designed for preterm infants as a nutritional supplement to be added to mother's human milk.

**CAUTION:** Nutritionally incomplete. To be used only under the supervision of a physician.

**WARNING:** The baby's health depends on carefully following these instructions. Use only as directed by a medical professional. Improper hygiene, preparation, dilution, use or storage may result in severe harm. Although this powder is formulated for premature infants, nutritional powders are not sterile and should **not** be fed to premature infants or infants who might have immune problems unless directed and supervised by a doctor.

- Follow hospital rules or the doctor's instructions for the safe handling of human milk.
- To aid mixing, agitate the human milk well. Pour the desired amount into a sterile container and warm to feeding temperature.
- Add the powder to the human milk according to the following chart:

Additional Calories Desired	Human Milk	Enfamil® Human Milk Fortifier
2 kcal/30 mL	50 mL	1 packet
4 kcal/30mL	25 mL	1 packet

The baby's doctor will provide instructions for the desired amount of calories to add.

- Failure to follow these instructions could result in severe harm. Once prepared, fortified breast milk can spoil quickly.
- Either feed fortified human milk immediately or cover and store in refrigerator at 2-4°C (35-40°F) for no longer than 24 hours. Agitate before each use.
- For tube feeding: Once fortified human milk is prepared it can safely remain at room temperature for 4 hours.<sup>2</sup> The Academy of Nutrition and Dietetics (formerly known as the American Dietetic Association) recommends a hang time for fortified breast milk of no longer than 4 hours at room temperature (25°C/77°F).<sup>3</sup>
- For bottle feeding: Pour only the amount of fortified human milk to be fed
  into a feeding container and feed immediately. Do not use fortified human
  milk if it is unrefrigerated for more than a total of 2 hours. After feeding
  begins, do not refrigerate feeding bottle. You must use within 1 hour or
  discard.
- WARNING: Do not use a microwave oven to warm formula. Serious burns may result.

#### When Using This Product:

- Monitor and regularly assess blood levels such as serum calcium, sodium, urea nitrogen, albumin, and others. In case of an elevated level, reduce the amount of Enfamil® Human Milk Fortifier used to supplement breast milk. In case of deficiency, additional supplementation with the deficient nutrient may be indicated.
- 2. Do not administer other nutrient supplements in addition to Enfamil® Human Milk Fortifier unless appropriate tests indicate a need for further supplementation with a particular nutrient.
- 3. Do not add Enfamil® Human Milk Fortifier to breast milk in a ratio greater than 1 packet/25 mL.
- 4. Preterm breast milk collected less than 2 weeks postpartum is particularly rich in nutrients, so fortification with Enfamil® Human Milk Fortifier is generally not required.

If more than 25 packets daily are used, the infant should be monitored for evidence of excessive vitamin A and D intake.

**STORAGE:** Store Enfamil® Human Milk Fortifier powder at room temperature. Avoid freezing and excessive heat. Use by date on package.

Neither expressed human milk nor nutritional powders are sterile.

**CAUTION:** Regarding use in extremely-low-birth-weight infants (ELBW-1 kg or less): Hypercalcemia has been reported in some of these infants on full enteral feeds of mother's milk supplemented with human milk fortifiers.

Telang S, Berseth CL, Ferguson PW, et al. Fortifying fresh human milk with commercial powdered human milk fortifiers does not affect bacterial growth during 6 hours at room temperature. J Am Diet Assoc. 2005;105:1567-1572.

<sup>3.</sup> Robbins ST, Meyers R, eds. Infant Feedings: Guidelines for Preparation of Human Milk and Formula in Health Care Facilities. Chicago, Ill: American Dietetic Association; 2011.



## Enfamil® Liquid Human Milk Fortifier Standard Protein, 5 mL Sachet

Please contact your Hospital Account Manager for product access and contract inclusion.

O D (M) IFANCA

#### INDICATION

Enfamil® Liquid Human Milk Fortifier Standard Protein is added to expressed breast milk for feeding premature and late preterm infants. It increases protein, energy, calcium, phosphorus, iron, vitamin D and other nutrients to help promote growth in premature infants. It has 3.4 g of protein per 100 Cal when mixed with preterm human milk at the 24 Calorie per fluid ounce concentration.\*

#### **PRODUCT BENEFITS**

Enfamil® Liquid Human Milk Fortifier is designed for premature infants as a nutritional supplement to be added to breast milk.

- Convenient, single-serve options for flexibility in delivery
- Easy-to-mix commercially sterile liquid
- Ultra-concentrated formulation minimizes dilution of breast milk
- DHA and ARA to help support brain and eye development
- Non-acidified formulation
- · Transparent sachet for viewing product

#### **NUTRIENTS**

Analysis	Per 5 mL Fortifier only	20 mL Plus 100 mL Preterm Human Milk (120 mL mixed to 24 Cal/fl oz concentration) **	Per 100 Cal Fortified Preterm Human Milk when mixed to 24 Cal/fl oz concentration <sup>†1</sup>
Calories, kcal	7.5	97	100
Protein, g	0.42	3.3	3.4
Fat, g	0.58	5.8	6
Linoleic acid, mg	58	710	730
Carbohydrate, g	0.28	8.4	8.7
Water, g	3.8	104	107



<sup>\*</sup> When mixed as directed at 5 mL HMF + 25 mL preterm human milk.

Analysis	20 mL Plus 100 mL Preterm Per 5 mL Fortifier only (120 mL mixed to 24 Cal/fl oz concentration) **		Per 100 Cal Fortified Preterm Human Milk when mixed to 24 Cal/fl oz concentration <sup>11</sup>	
Vitamins/Other Nutrie	ents			
Vitamin A, IU	290	1210	1240	
Vitamin D, IU	47	200	200	
Vitamin E, IU	1.4	6	6.2	
Vitamin K, mcg	1.42	7.7	7.9	
Thiamin (Vitamin $B_1$ ), mcg	46	193	200	
Riboflavin (Vitamin B <sub>2</sub> ), mcg	66	290	300	
Vitamin B <sub>6</sub> , mcg	35	146	150	
Vitamin B <sub>12</sub> , mcg	0.16	0.66	0.68	
Niacin, mcg	920 3900		4000	
Folic Acid (Folacin), mcg	7.7		35	
Pantothenic acid, mcg	230	1150	1190	
Biotin, mcg	0.84	3.9	4.1	
Vitamin C (Ascorbic acid), mg	3.8	20	20	
Minerals				
Calcium, mg	29	141	145	
Phosphorus, mg	15.8	78	80	
Magnesium, mg	2	11.3	11.6	
Iron, mg	0.44	1.85	1.9	
Zinc, mg	nc, mg 0.24 1.33		1.37	
Manganese, mcg	2.5	10.4	10.7	
Copper, mcg	15	98	101	
Sodium, mg	6.8	55	57	
Potassium, mg	11.3	95	98	
Chloride, mg	6.9	86	88	

<sup>†</sup> Calculated using values for preterm human breast milk compiled from multiple published studies; protein value used for preterm human milk is 1.62 g/100 mL.

Koletzko B, Cheah F-C, Domellöf M, van Goudoever JB, Poindexter BB, Vain N. Scientific Basis and Practical Application of Nutritional Care for Preterm Infants. World Rev Nutr Diet. 2021;122:XIII-XIV.

### FORTIFIED HUMAN MILK INFORMATION

Fortified Preterm Human Milk Prepared to 22 or 24 Cal/fl oz	Per 100 Cal at 22 Cal/fl oz	Per 100 Cal at 24 Cal/fl oz	
Recipe			
Preterm human milk + Enfamil® Liquid HMF Standard Protein	50 mL + 5 mL	25 mL + 5 mL	
Final volume, mL	55	30	
Nutrients			
Protein, g	3	3.4	
Fat, g	5.7	6	
Carbohydrate, g	9.6	8.7	
Water, g	118	107	
Vitamins			
Vitamin A activity, IU	770	1240	
Vitamin D, IU	124	200	
Vitamin E, IU	3.9	6.2	
Vitamin K, mcg	5.9	7.9	
Thiamin (Vitamin B <sub>1</sub> ), mcg	123	200	
Riboflavin (Vitamin B <sub>2</sub> ), mcg	194	300	
Vitamin B <sub>6</sub> , mcg	93	150	
Vitamin B <sub>12</sub> , mcg	0.41	0.68	
Niacin, mcg	2500	4000	
Folic acid (Folacin), mcg	23	35	
Pantothenic acid, mcg	840	1190	
Biotin, mcg	2.7	4.1	
Vitamin C (Ascorbic acid), mg	14.6	20	
Minerals			
Calcium, mg	101	145	
Phosphorus, mg	56	80	
Magnesium, mg	8.9	11.6	
Iron, mg	1.18	1.9	
Zinc, mg	1.04	1.37	
Manganese, mcg	6.5	10.7	
Copper, mcg	83	101	

Fortified Preterm Human Milk Prepared to 22 or 24 Cal/fl oz	Per 100 Cal at 22 Cal/fl oz	Per 100 Cal at 24 Cal/fl oz	
lodine, mcg	22	18.4	
Selenium, mcg	2.9	2.5	
Sodium, mg	51	57	
Potassium, mg	89	98	
Chloride, mg	88	88	
Linoleic acid, mg	730	730	
Linolenic acid, mg	54	60	
ARA, mg	32	38	
DHA, mg	21	24	

Fortified Preterm Human Milk Prepared to 22 or 24 Cal/fl oz	Per 100 mL at 22 Cal/fl oz	Per 100 mL at 24 Cal/fl oz	
Recipe			
Preterm human milk + Enfamil® Liquid HMF Standard Protein	50 mL + 5 mL	25 mL + 5 mL	
Final volume, mL	55	30	
Nutrients			
Energy, Cal	75	81	
Protein, g	2.2	2.8	
Fat, g	4.2	4.8	
Carbohydrate, g	7.1	7	
Water, g	88	87	
Vitamins/Other Nutrients			
Vitamin A activity, IU	570	1010	
Vitamin D, IU	93 167		
Vitamin E, IU	2.9	5	
Vitamin K, mcg	4.4	6.4	
Thiamin (Vitamin B <sub>1</sub> ), mcg	92	161	
Riboflavin (Vitamin B <sub>2</sub> ), mcg	145	240	
Vitamin B <sub>6</sub> , mcg	69	122	
Vitamin B <sub>12</sub> , mcg	0.31	0.55	
Niacin, mcg	1860	3300	

Fortified Preterm Human Milk Prepared to 22 or 24 Cal/fl oz	Per 100 mL at 22 Cal/fl oz	Per 100 mL at 24 Cal/fl oz	
Folic acid (Folacin), mcg	16.8	28	
Pantothenic acid, mcg	630	960	
Biotin, mcg	2	3.3	
Vitamin C (Ascorbic acid), mg	10.9	16.7	
Minerals			
Calcium, mg	75	118	
Phosphorus, mg	42	65	
Magnesium, mg	6.6	9.4	
Iron, mg	0.88	1.54	
Zinc, mg	0.77	1.11	
Manganese, mcg	4.9	8.7	
Copper, mcg	62	82	
lodine, mcg	16.2	14.8	
Selenium, mcg	2.2	2	
Sodium, mg	38	46	
Potassium, mg	66	79	
Chloride, mg	65	72	
Additional Nutrients			
Linoleic acid, mg	540	590	
Linolenic acid, mg	40	48	
ARA, mg	24	31	
DHA, mg	15.6	19.2	

Fortified Term Human Milk <sup>1-5</sup> Prepared to 22 or 24 Cal/fl oz	Per 100 Cal at 22 Cal/fl oz	Per 100 Cal at 24 Cal/fl oz	
Recipe			
Term human milk + Enfamil® Liquid HMF Standard Protein	50 mL + 5 mL	25 mL + 5 mL	
Final volume, mL	55 30		
Nutrients			
Protein <sup>1</sup> , g	2.1	2.7	
Fat <sup>1</sup> , g	5.6	6	

Fortified Term Human Milk <sup>1-5</sup> Prepared to 22 or 24 Cal/fl oz	Per 100 Cal at 22 Cal/fl oz	Per 100 Cal at 24 Cal/fl oz	
Carbohydrate <sup>1</sup> , g	10.4	9.4	
Water, g	118	108	
Vitamins/Other Nutrients			
Vitamin A activity <sup>1</sup> , IU	880	1340	
Vitamin D¹, IU	116	194	
Vitamin E¹, IU	4.4	6.6	
Vitamin K <sup>1</sup> , mcg	3.7	6.2	
Thiamin (Vitamin B <sub>1</sub> )¹, mcg	136	210	
Riboflavin (Vitamin B <sub>2</sub> ) <sup>1</sup> , mcg	220	320	
Vitamin B <sub>6</sub> ¹, mcg	109	164	
Vitamin B <sub>12</sub> 1, mcg	0.48	0.74	
Niacin¹, mcg	2700	4200	
Folic acid (Folacin) <sup>1</sup> , mcg	32	43	
Pantothenic acid <sup>1</sup> , mcg	830	1180	
Biotin <sup>1</sup> , mcg	2.9	4.2	
Vitamin C (Ascorbic acid) <sup>1</sup> , mg	21	26	
Choline², mg	19.4	16.4	
Inositol³, mg	32	27	
Minerals			
Calcium¹, mg	98	143	
Phosphorus <sup>1</sup> , mg	54	78	
Magnesium <sup>1</sup> , mg	8.8	11.6	
Iron <sup>1</sup> , mg	1.14	1.87	
Zinc¹, mg	0.82	1.19	
Manganese <sup>1</sup> , mcg	6.4	10.6	
Copper <sup>1</sup> , mcg	73	92	
lodine <sup>1</sup> , mcg	18.2	15.4	
Selenium <sup>1</sup> , mcg	2.4	2.1	
Sodium <sup>1</sup> , mg	39	47	
Potassium <sup>1</sup> , mg	85	95	
Chloride <sup>1</sup> , mg	68	73	

Fortified Term Human Milk <sup>1-5</sup> Prepared to 22 or 24 Cal/fl oz	Per 100 Cal at 22 Cal/fl oz	Per 100 Cal at 24 Cal/fl oz	
Additional Nutrients			
Linoleic acid <sup>4</sup> , mg	830	820	
Linolenic acid <sup>4</sup> , mg	62	67	
ARA <sup>5</sup> , mg	32	38	
DHA <sup>5</sup> , mg	21	24	

Fortified Term Human Milk <sup>1-5</sup> Prepared to 22 or 24 Cal/fl oz	Per 100 mL at 22 Cal/fl oz	Per 100 mL at 24 Cal/fl oz
Recipe		
Term human milk + Enfamil® Liquid HMF Standard Protein	50 mL + 5 mL	25 mL + 5 mL
Final volume, mL	55	30
Nutrients		
Energy¹, Cal	75	81
Protein <sup>1</sup> , g	1.58	2.2
Fat <sup>1</sup> , g	4.2	4.8
Carbohydrate <sup>1</sup> , g	7.8	7.6
Water, g	89	88
Vitamins/Other Nutrients		
Vitamin A activity <sup>1</sup> , IU	660	1090
Vitamin D¹, IU	87	158
Vitamin E <sup>1</sup> , IU	3.3	5.3
Vitamin K <sup>1</sup> , mcg	2.8 5	
Thiamin (Vitamin B <sub>1</sub> )¹, mcg	102	167
Riboflavin (Vitamin B <sub>2</sub> ) <sup>1</sup> , mcg	165	260
Vitamin B <sub>6</sub> ¹, mcg	82	133
Vitamin B <sub>12</sub> 1, mcg	0.36	0.6
Niacin¹, mcg	2000	3400
Folic acid (Folacin)¹, mcg	24	35
Pantothenic acid¹, mcg	620	960
Biotin <sup>1</sup> , mcg	2.2	3.4
Vitamin C (Ascorbic acid) <sup>1</sup> , mg	16	21

Fortified Term Human Milk <sup>1-5</sup> Prepared to 22 or 24 Cal/fl oz	Per 100 mL at 22 Cal/fl oz	Per 100 mL at 24 Cal/fl oz
Choline², mg	14.5	13.3
Inositol³, mg	24	22
Minerals		
Calcium¹, mg	73	116
Phosphorus <sup>1</sup> , mg	41	63
Magnesium¹, mg	6.6	9.4
Iron <sup>1</sup> , mg	0.85	1.52
Zinc¹, mg	0.62	0.97
Manganese <sup>1</sup> , mcg	4.8	8.6
Copper <sup>1</sup> , mcg	55	75
lodine <sup>1</sup> , mcg	13.6	12.5
Selenium <sup>1</sup> , mcg	1.82	1.67
Sodium <sup>1</sup> , mg	29	38
Potassium¹, mg	64	78
Chloride <sup>1</sup> , mg	51	59
Additional Nutrients		
Linoleic acid <sup>4</sup> , mg	620	670
Linolenic acid <sup>4</sup> , mg	46	54
ARA⁵, mg	24	31
DHA <sup>5</sup> , mg	15.6	19.2

- 1. Picciano MF. Nutrient composition of human milk. Pediatr Clin North Am. 2001 Feb;48(1):53-67, 263-264.
- 2. Institute of Medicine (US) Standing Committee on the Scientific Evaluation of Dietary Reference Intakes and its Panel on Folate, Other B Vitamins, and Choline. Dietary Reference Intakes for Thiamin, Riboflavin, Niacin, Vitamin B6, Folate, Vitamin B12, Pantothenic Acid, Biotin, and Choline. Washington (DC): National Academies Press (US); 1998.
- 3. Pereira GR, Baker L, Egler J, Corcoran L, Chiavacci R. Serum myoinositol concentrations in premature infants fed human milk, formula for infants, and parenteral nutrition. Am J Clin Nutr. 1990;51[4]:589-93.
- 4. Innis SM. Human milk and formula fatty acids. J Pediatr. 1992; 120(4 Pt 2):S56-61.
- Brenna JT, Varamini B, Jensen RG, Diersen-Schade DA, Boettcher JA, Arterburn LM. Docosahexaenoic and arachidonic acid concentrations in human breast milk worldwide. Am J Clin Nutr. 2007;85(6):1457-64.

Nutrient Density	Fortified Preterm Human Milk to 24 Cal (1:5 ratio HMF to preterm milk)	Fortified Preterm Human Milk to 22 Cal (1:10 ratio HMF to preterm milk)
Potential Renal Solute Load (m0sm/100 Calories) <sup>6</sup>	30	26
Osmolality (mOsm/kg H <sub>2</sub> O)	300-360	280-340
рН*	5.5-6.3	5.8-6.3

<sup>\*</sup> The pH of any product can vary depending upon environmental factors such as temperature or as the product ages. This value represents the pH of products over shelf life.

#### **INGREDIENTS**

Water, whey protein isolate hydrolysate (milk), medium chain triglycerides (MCT oil), maltodextrin, soy oil, and less than 2%: Mortierella alpina oil $^{\dagger}$ , Schizochytrium sp. oil $^{\ddagger}$ , magnesium oxide, calcium citrate, calcium phosphate, potassium citrate, calcium chloride, sodium citrate, magnesium phosphate, ferrous sulfate, zinc sulfate, cupric sulfate, pectin, ascorbic acid, vitamin E acetate, niacinamide, calcium pantothenate, vitamin A palmitate, riboflavin, thiamin hydrochloride, vitamin B $_{6}$  hydrochloride, folic acid, vitamin K $_{1}$ , biotin, vitamin B $_{12}$ , vitamin D $_{3}$ .

- † A source of arachidonic acid (ARA).
- ± A source of docosahexaenoic acid (DHA).

#### **ALLERGEN INFORMATION**

Enfamil® Liquid Human Milk Fortifier Standard Protein contains milk and soy.

#### **PRODUCT FORMS**

Product	Format	Unit Size	Approximate Product Yield at Normal Dilution	Shelf Life	Item Number	Availability
Enfamil® Liquid Human Milk Fortifier Standard Protein Sachet	Liquid	5 mL Sachets /144 per case	1 sachet = 5 mL of liquid HMF + 25 mL of breast milk	12 months	3180545	Hospital & available on discharge by hospital dispensing

#### **CONCENTRATIONS AND PREPARATION**

The baby's health depends on carefully following the instructions below. Use only as directed by a medical professional. Improper hygiene, preparation, concentration, use or storage may result in severe harm.

<sup>6.</sup> Fomon SJ, Ziegler EE. Renal solute load and potential renal solute load in infancy. J Pediatr. 1999;134(1):11-14.

Follow hospital rules or the baby's doctor's instructions for safe handling and fortification of breast milk. Opened sachets can spoil quickly. Use fortifier immediately and discard any remaining or unused fortifier.

This product is commercially sterile as produced.

#### **Directions for Use:**

- Pour desired quantity of breast milk into a sterile feeding container.
- Gently massage HMF sachet until uniform as some product separation is normal.
- Use perforation to completely tear open sachet.
- Add desired amount of HMF to breast milk in sterile feeding container.
- Shake until mixed well. Fortified breast milk can spoil quickly. Either feed fortified breast milk immediately or cover and store in refrigerator at 35-40°F (2-4°C) for no longer than 24 hours. After feeding begins, use fortified breast milk within one hour or discard.

#### Directions for When Stored:

Refrigerate unused fortified breast milk and use within 24 hours. After refrigeration, gently remix and feed immediately. Discard remaining fortified breast milk promptly after bottle feeding session.

#### Storage:

Store unopened sachets in carton at room temperature. Avoid excessive heat and prolonged exposure to light. Do not freeze.

**WARNING:** Do not use a microwave oven to warm the fortified breast milk. Serious burns may result.

WARNING: Do not administer directly - must add to breast milk.

WARNING: Not for parenteral (I.V.) use.

Do not use if sachet is damaged. Use by date on carton or sachet.

Additional Calories Desired	Breast Milk	Enfamil® Liquid Human Milk Fortifier Standard Protein	Total Concentration (breast milk + LHMF)
2 Calories/ fl oz	50 mL	5 mL	22 Cal/fl oz concentration
4 Calories/ fl oz	25 mL	5 mL	24 Cal/fl oz concentration



## Enfamil® Liquid Human Milk Fortifier High Protein, 5 mL Sachet

Please contact your Hospital Account Manager for product access and contract inclusion.

OD M IFANCA

#### INDICATION

Enfamil® Liquid Human Milk Fortifier High Protein is added to expressed breast milk for feeding premature and low-birth-weight infants. It provides enriched nutrition needed by these special babies while still allowing them to receive their mother's milk. Enfamil® Liquid Human Milk Fortifier High Protein increases the levels of protein, energy, calcium, phosphorus, iron, vitamin D and other nutrients, producing a diet more suited to the nutritional needs of the rapidly growing premature infant. It has 4 g of protein per 100 Cal when mixed with preterm human milk at the 24 Calorie per fluid ounce concentration.\*

#### PRODUCT BENEFITS

Enfamil® Liquid Human Milk Fortifier is designed for premature infants as a nutritional supplement to be added to breast milk.

- Convenient, single-serve options for flexibility in delivery
- Easy-to-mix commercially sterile liquid
- Ultra-concentrated formulation minimizes dilution of breast milk
- DHA and ARA to help support brain and eye development
- Non-acidified formulation
- Transparent sachet for viewing product

#### **NUTRIENTS**

Analysis	Per 5 mL Fortifier only	20 mL Plus 100 mL Preterm Human Milk (120 mL mixed to 24 Cal/fl oz concentration) **	Per 100 Cal Fortified Preterm Human Milk when mixed to 24 Cal/fl oz concentration <sup>†1</sup>
Calories, kcal	7.5	97	100
Protein, g	0.56	3.9	4
Fat, g	0.58	5.8	6
Linoleic acid, mg	58	710	730
Carbohydrate, g	0.11	7.7	7.9
Water, g	3.8	104	107



<sup>\*</sup> When mixed as directed at 5 mL HMF + 25 mL preterm human milk.

Analysis	Per 5 mL Fortifier only	20 mL Plus 100 mL Preterm Human Milk (120 mL mixed to 24 Cal/fl oz concentration) †1	Per 100 Cal Fortified Preterm Human Milk when mixed to 24 Cal/fl oz concentration <sup>11</sup>
Vitamins/Other Nutrie	ents		
Vitamin A, IU	290	1210	1240
Vitamin D, IU	47	200	200
Vitamin E, IU	1.4	6	6.2
Vitamin K, mcg	1.42	7.7	7.9
Thiamin (Vitamin $B_1$ ), mcg	46	193	200
Riboflavin (Vitamin B <sub>2</sub> ), mcg	66	290	300
Vitamin B <sub>6</sub> , mcg	35	146	150
Vitamin B <sub>12</sub> , mcg	0.16	0.66	0.68
Niacin, mcg	920	3900	4000
Folic acid (Folacin), mcg	7.7	34	35
Pantothenic acid, mcg	230	1150	1190
Biotin, mcg	0.84	3.9	4.1
Vitamin C (Ascorbic acid), mg	3.8	20	20
Minerals			
Calcium, mg	29	141	145
Phosphorus, mg	15.8	78	80
Magnesium, mg	2	11.3	11.6
Iron, mg	0.44	1.85	1.9
Zinc, mg	0.24	1.33	1.37
Manganese, mcg	2.5	10.4	10.7
Copper, mcg	15	98	101
Sodium, mg	6.8	55	57
Potassium, mg	11.3	95	98
Chloride, mg	6.9	86	88

<sup>†</sup> Calculated using values for preterm human breast milk compiled from multiple published studies; protein value used for preterm human milk is 1.62 g/100 mL.

Koletzko B, Cheah F-C, Domellöf M, van Goudoever JB, Poindexter BB, Vain N. Scientific Basis and Practical Application of Nutritional Care for Preterm Infants. World Rev Nutr Diet. 2021;122:XIII-XIV.

### FORTIFIED HUMAN MILK INFORMATION

Fortified Preterm Human Milk Prepared to 22 or 24 Cal/fl oz	Per 100 Cal at 22 Cal/fl oz	Per 100 Cal at 24 Cal/fl oz
Recipe		
Preterm human milk + Enfamil® Liquid HMF High Protein	50 mL + 5 mL	25 mL + 5 mL
Final volume, mL	55	30
Nutrients		
Protein, g	3.3	4
Fat, g	5.7	6
Carbohydrate, g	9.2	7.9
Water, g	118	107
Vitamins		
Vitamin A activity, IU	770	1240
Vitamin D, IU	124	200
Vitamin E, IU	3.9	6.2
Vitamin K, mcg	5.9	7.9
Thiamin (Vitamin B <sub>1</sub> ), mcg	123	200
Riboflavin (Vitamin B <sub>2</sub> ), mcg	194	300
Vitamin B <sub>6</sub> , mcg	93	150
Vitamin B <sub>12</sub> , mcg	0.41	0.68
Niacin, mcg	2500	4000
Folic acid (Folacin), mcg	23	35
Pantothenic acid, mcg	840	1190
Biotin, mcg	2.7	4.1
Vitamin C (Ascorbic acid), mg	14.6	20
Minerals		
Calcium, mg	101	145
Phosphorus, mg	56	80
Magnesium, mg	8.9	11.6
Iron, mg	1.18	1.9
Zinc, mg	1.04	1.37
Manganese, mcg	6.5	10.7
Copper, mcg	83	101

Fortified Preterm Human Milk Prepared to 22 or 24 Cal/fl oz	Per 100 Cal at 22 Cal/fl oz	Per 100 Cal at 24 Cal/fl oz
lodine, mcg	22	18.4
Selenium, mcg	2.9	2.5
Sodium, mg	51	57
Potassium, mg	89	98
Chloride, mg	88	88
Linoleic acid, mg	730	730
Linolenic acid, mg	54	60
ARA, mg	32	38
DHA, mg	21	24

Fortified Preterm Human Milk Prepared to 22 or 24 Cal/fl oz	Per 100 mL at 22 Cal/fl oz	Per 100 mL at 24 Cal/fl oz
Recipe		
Preterm human milk + Enfamil® Liquid HMF High Protein	50 mL + 5 mL	25 mL + 5 mL
Final volume, mL	55	30
Nutrients		
Energy, Cal	75	81
Protein, g	2.5	3.3
Fat, g	4.2	4.8
Carbohydrate, g	6.8	6.4
Water, g	88	87
Vitamins/Other Nutrients		
Vitamin A activity, IU	570	1010
Vitamin D, IU	93	167
Vitamin E, IU	2.9	5
Vitamin K, mcg	4.4	6.4
Thiamin (Vitamin B <sub>1</sub> ), mcg	92	161
Riboflavin (Vitamin B <sub>2</sub> ), mcg	145	240
Vitamin B <sub>6</sub> , mcg	69	122
Vitamin B <sub>12</sub> , mcg	0.31	0.55
Niacin, mcg	1860	3300

Fortified Preterm Human Milk Prepared to 22 or 24 Cal/fl oz	Per 100 mL at 22 Cal/fl oz	Per 100 mL at 24 Cal/fl oz
Folic acid (Folacin), mcg	16.8	28
Pantothenic acid, mcg	630	960
Biotin, mcg	2	3.3
Vitamin C (Ascorbic acid), mg	10.9	16.7
Minerals		
Calcium, mg	75	118
Phosphorus, mg	42	65
Magnesium, mg	6.6	9.4
Iron, mg	0.88	1.54
Zinc, mg	0.77	1.11
Manganese, mcg	4.9	8.7
Copper, mcg	62	82
lodine, mcg	16.2	14.8
Selenium, mcg	2.2	2
Sodium, mg	38	46
Potassium, mg	66	79
Chloride, mg	65	72
Additional Nutrients		
Linoleic acid, mg	540	590
Linolenic acid, mg	40	48
ARA, mg	24	31
DHA, mg	15.6	19.2

Fortified Term Human Milk <sup>1-5</sup> Prepared to 22 or 24 Cal/fl oz	Per 100 Cal at 22 Cal/fl oz	Per 100 Cal at 24 Cal/fl oz
Recipe		
Term human milk + Enfamil® Liquid HMF High Protein	50 mL + 5 mL	25 mL + 5 mL
Final volume, mL	55	30
Nutrients		
Protein <sup>1</sup> , g	2.4	3.3
Fat <sup>1</sup> , g	5.6	6

Fortified Term Human Milk <sup>1-5</sup> Prepared to 22 or 24 Cal/fl oz	Per 100 Cal at 22 Cal/fl oz	Per 100 Cal at 24 Cal/fl oz
Carbohydrate <sup>1</sup> , g	10	8.7
Water, g	118	108
Vitamins/Other Nutrients		
Vitamin A activity <sup>1</sup> , IU	880	1340
Vitamin D <sup>1</sup> , IU	116	194
Vitamin E <sup>1</sup> , IU	4.4	6.6
Vitamin K <sup>1</sup> , mcg	3.7	6.2
Thiamin (Vitamin $B_1$ ) $^1$ , mcg	122	200
Riboflavin (Vitamin B <sub>2</sub> )¹, mcg	220	320
Vitamin B <sub>6</sub> ¹, mcg	109	164
Vitamin B <sub>12</sub> 1, mcg	0.48	0.74
Niacin <sup>1</sup> , mcg	2700	4200
Folic acid (Folacin)¹, mcg	32	43
Pantothenic acid <sup>1</sup> , mcg	830	1180
Biotin <sup>1</sup> , mcg	2.9	4.2
Vitamin C (Ascorbic acid) <sup>1</sup> , mg	21	26
Choline², mg	19.4	16.4
Inositol³, mg	32	27
Minerals		
Calcium¹, mg	98	143
Phosphorus <sup>1</sup> , mg	54	78
Magnesium <sup>1</sup> , mg	8.8	11.6
Iron <sup>1</sup> , mg	1.14	1.87
Zinc¹, mg	0.82	1.19
Manganese <sup>1</sup> , mcg	6.4	10.6
Copper <sup>1</sup> , mcg	73	92
lodine <sup>1</sup> , mcg	18.2	15.4
Selenium <sup>1</sup> , mcg	2.4	2.1
Sodium <sup>1</sup> , mg	39	47
Potassium¹, mg	85	95
Chloride <sup>1</sup> , mg	68	73

Fortified Term Human Milk <sup>1-5</sup> Prepared to 22 or 24 Cal/fl oz	Per 100 Cal at 22 Cal/fl oz	Per 100 Cal at 24 Cal/fl oz
Additional Nutrients		
Linoleic acid <sup>4</sup> , mg	830	820
Linolenic acid <sup>4</sup> , mg	62	67
ARA⁵, mg	32	38
DHA <sup>5</sup> , mg	21	24

Fortified Term Human Milk <sup>1-5</sup> Prepared to 22 or 24 Cal/fl oz	Per 100 mL at 22 Cal/fl oz	Per 100 mL at 24 Cal/fl oz
Recipe		
Term human milk + Enfamil Liquid HMF High Protein	50 mL + 5 mL	25 mL + 5 mL
Final volume, mL	55	30
Nutrients		
Energy¹, Cal	75	81
Protein <sup>1</sup> , g	1.84	2.7
Fat <sup>1</sup> , g	4.2	4.8
Carbohydrate <sup>1</sup> , g	7.5	7
Water, g	89	88
Vitamins/Other Nutrients		
Vitamin A activity <sup>1</sup> , IU	660	1090
Vitamin D¹, IU	87	158
Vitamin E¹, IU	3.3	5.3
Vitamin K <sup>1</sup> , mcg	2.8	5
Thiamin (Vitamin B <sub>1</sub> )¹, mcg	92	161
Riboflavin (Vitamin B <sub>2</sub> ) <sup>1</sup> , mcg	165	260
Vitamin B <sub>6</sub> ¹, mcg	82	133
Vitamin B <sub>12</sub> 1, mcg	0.36	0.6
Niacin¹, mcg	2000	3400
Folic acid (Folacin) <sup>1</sup> , mcg	24	35
Pantothenic acid¹, mcg	620	960
Biotin <sup>1</sup> , mcg	2.2	3.4
Vitamin C (Ascorbic acid) <sup>1</sup> , mg	16	21

Fortified Term Human Milk <sup>1-5</sup> Prepared to 22 or 24 Cal/fl oz	Per 100 mL at 22 Cal/fl oz	Per 100 mL at 24 Cal/fl oz
Choline², mg	14.5	13.3
Inositol³, mg	24	22
Minerals		
Calcium¹, mg	73	116
Phosphorus <sup>1</sup> , mg	41	63
Magnesium <sup>1</sup> , mg	6.6	9.4
Iron <sup>1</sup> , mg	0.85	1.52
Zinc¹, mg	0.62	0.97
Manganese <sup>1</sup> , mcg	4.8	8.6
Copper <sup>1</sup> , mcg	55	75
lodine <sup>1</sup> , mcg	13.6	12.5
Selenium¹, mcg	1.82	1.67
Sodium <sup>1</sup> , mg	29	38
Potassium¹, mg	64	78
Chloride <sup>1</sup> , mg	51	59
Additional Nutrients		
Linoleic acid <sup>4</sup> , mg	620	670
Linolenic acid <sup>4</sup> , mg	46	54
ARA <sup>5</sup> , mg	24	31
DHA⁵, mg	15.6	19.2

- 1. Picciano MF. Nutrient composition of human milk. Pediatr Clin North Am. 2001 Feb;48(1):53-67, 263-264.
- 2. Institute of Medicine (US) Standing Committee on the Scientific Evaluation of Dietary Reference Intakes and its Panel on Folate, Other B Vitamins, and Choline. Dietary Reference Intakes for Thiamin, Riboflavin, Niacin, Vitamin B6, Folate, Vitamin B12, Pantothenic Acid, Biotin, and Choline. Washington (DC): National Academies Press (US); 1998.
- 3. Pereira GR, Baker L, Egler J, Corcoran L, Chiavacci R. Serum myoinositol concentrations in premature infants fed human milk, formula for infants, and parenteral nutrition. Am J Clin Nutr. 1990;51(4):589-93.
- 4. Innis SM. Human milk and formula fatty acids. J Pediatr. 1992; 120(4 Pt 2):S56-61.
- Brenna JT, Varamini B, Jensen RG, Diersen-Schade DA, Boettcher JA, Arterburn LM. Docosahexaenoic and arachidonic acid concentrations in human breast milk worldwide. Am J Clin Nutr. 2007;85(6):1457-64.

Nutrient Density	Fortified Preterm Human Milk to 24 Cal (1:5 ratio HMF to preterm milk)	Fortified Preterm Human Milk to 22 Cal (1:10 ratio HMF to preterm milk)
Potential Renal Solute Load (m0sm/100 Calories) <sup>6</sup>	33	28
Osmolality (mOsm/kg H <sub>2</sub> O)	290-350	280-340
рН*	5.5-6.3	5.8-6.3

<sup>\*</sup> The pH of any product can vary depending upon environmental factors such as temperature or as the product ages. This value represents the pH of products over shelf life.

#### **INGREDIENTS**

Water, whey protein isolate hydrolysate (milk), medium chain triglycerides (MCT oil), soy oil and less than 2%: Mortierella alpina oil<sup>†</sup>, Schizochytrium Sp. oil<sup>‡</sup>, magnesium oxide, pectin, vitamin A palmitate, vitamin E acetate, vitamin D<sub>3</sub>, vitamin K<sub>1</sub>, thiamin hydrochloride, riboflavin, vitamin B<sub>6</sub> hydrochloride, vitamin B<sub>12</sub>, niacinamide, folic acid, calcium pantothenate, biotin, ascorbic acid, calcium citrate, calcium phosphate, cupric sulfate, sodium citrate, potassium citrate, calcium chloride, ferrous sulfate, magnesium phosphate, zinc sulfate.

#### **ALLERGEN INFORMATION**

Enfamil® Liquid Human Milk Fortifier High Protein contains milk and soy.

#### **PRODUCT FORMS**

Product	Format	Unit Size	Approximate Product Yield at Normal Dilution	Shelf Life	Item Number	Availability
Enfamil® Liquid Human Milk Fortifier High Protein Sachet	Liquid	5 mL Sachets /144 per case	1 sachet = 5 mL of liquid HMF + 25 mL of breast milk	12 months	3180315	Hospital & available on discharge by hospital dispensing

#### **CONCENTRATIONS AND PREPARATION**

The baby's health depends on carefully following the instructions below. Use only as directed by a medical professional. Improper hygiene, preparation, concentration, use or storage may result in severe harm.

<sup>6.</sup> Fomon SJ, Ziegler EE. Renal solute load and potential renal solute load in infancy. J Pediatr. 1999;134[1]:11-14.

<sup>†</sup> A source of arachidonic acid (ARA).

<sup>±</sup> A source of docosahexaenoic acid (DHA).

Follow hospital rules or the baby's doctor's instructions for the safe handling and fortification of breast milk. Opened sachets can spoil quickly. Use fortifier immediately and discard any remaining, unused fortifier.

This product is commercially sterile as produced.

#### **Directions for Use:**

- Pour desired quantity of breast milk into a sterile feeding container.
- Gently massage HMF sachet until uniform as some product separation is normal.
- Use perforation to completely tear open sachet.
- Add desired amount of HMF to breast milk in sterile feeding container.
- Shake until mixed well. Fortified breast milk can spoil quickly. Either feed fortified breast milk immediately or cover and store in refrigerator at 35-40°F (2-4°C) for no longer than 24 hours. After feeding begins, use fortified breast milk within one hour or discard.

#### **Directions for When Stored:**

Refrigerate unused fortified breast milk and use within 24 hours. After refrigeration, gently remix and feed immediately. Discard remaining fortified breast milk promptly after bottle feeding session.

#### Storage:

Store unopened sachets in carton at room temperature. Avoid excessive heat and prolonged exposure to light. Do not freeze.

**WARNING:** Do not use a microwave oven to warm the fortified breast milk. Serious burns may result.

WARNING: Do not administer directly - must add to breast milk.

WARNING: Not for parenteral (I.V.) use.

Do not use if sachet is damaged. Use by date on carton or sachet.

Additional Calories Desired	Breast Milk	Enfamil Liquid Human Milk Fortifier High Protein	Total Concentration (breast milk + LHMF)
2 Calories/ fl oz	50 mL	5 mL	22 Cal/fl oz concentration
4 Calories/ fl oz	25 mL	5 mL	24 Cal/fl oz concentration



# Enfamil® Liquid Human Milk Fortifier Standard Protein

OD (M) IFANCA

#### INDICATION

Enfamil® Liquid Human Milk Fortifier Standard Protein is added to expressed breast milk for feeding premature and late preterm infants. It increases protein, energy, calcium, phosphorus, iron, vitamin D and other nutrients to help promote growth in premature infants. It has 3.4 g of protein per 100 Cal when mixed with preterm human milk at the 24 Calorie per fluid ounce concentration\*/0.81 kcal/mL concentration.

#### **NUTRIENTS**

Analysis	Per 5 mL Fortifier only	20 mL Plus 100 mL Preterm Breast Milk (120 mL mixed to 24 Cal/fl oz or 0.81 kcal/mL concentration)†	Per 100 Cal Fortified Preterm Breast Milk when mixed to 24 Cal/fl oz or 0.81 kcal/mL concentration†		
Calories, kcal	7.5	97	100		
Protein, g	0.42	3.3	3.4		
Fat, g	0.58	5.8	6		
Linoleic acid, mg	58	710	730		
Carbohydrate, g	0.28	8.4	8.7		
Water, g	3.8	104	107		
Vitamins/Other Nutrients					
Vitamin A, IU	290	1210	1240		
Vitamin D, IU	47	200	200		
Vitamin E, IU	1.4	6	6.2		
Vitamin K, mcg	1.42	7.7	7.9		
Thiamin (Vitamin B <sub>1</sub> ), mcg	46	193	200		
Riboflavin (Vitamin B <sub>2</sub> ), mcg	66	290	300		





<sup>\*</sup> When mixed as directed at 5 mL HMF + 25 mL preterm human milk.

Analysis	Per 5 mL Fortifier only	20 mL Plus 100 mL Preterm Breast Milk (120 mL mixed to 24 Cal/fl oz or 0.81 kcal/mL concentration) <sup>†</sup>	Per 100 Cal Fortified Preterm Breast Milk when mixed to 24 Cal/fl oz or 0.81 kcal/mL concentration <sup>†</sup>
Vitamin B <sub>6</sub> , mcg	35	146	150
Vitamin B <sub>12</sub> , mcg	0.16	0.66	0.68
Niacin, mcg	920	3900	4000
Folic Acid (Folacin), mcg	7.7	34	35
Pantothenic Acid, mcg	230	1150	1190
Biotin, mcg	0.84	3.9	4.1
Vitamin C (Ascorbic Acid), mg	3.8	20	20
Minerals			
Calcium, mg	29	141	145
Phosphorus, mg	15.8	78	80
Magnesium, mg	2	11.3	11.6
Iron, mg	11.6	2	11.3
Zinc, mg	0.24	1.33	1.37
Manganese, mcg	2.5	10.4	10.7
Copper, mcg	15	98	101
Sodium, mg	6.8	55	57
Potassium, mg	11.3	95	98
Chloride, mg	6.9	86	88

<sup>†</sup> Calculated using values for preterm human breast milk compiled from multiple published studies; protein value used for preterm human milk is 1.62 g/100 mL.¹

Water, whey protein isolate hydrolysate (milk), medium chain triglycerides (MCT oil), maltodextrin, soy oil, and less than 2%: Mortierella alpina oil<sup>‡</sup>, Schizochytrium sp. oil<sup>§</sup>, calcium citrate, calcium phosphate, potassium citrate, calcium chloride, sodium citrate, magnesium phosphate, ferrous sulfate, zinc sulfate, cupric sulfate, pectin, ascorbic acid, vitamin E acetate, niacinamide, calcium pantothenate, vitamin A palmitate, riboflavin, thiamin hydrochloride, vitamin B<sub>6</sub> hydrochloride, folic acid, vitamin K<sub>1</sub>, biotin, vitamin B<sub>12</sub>, vitamin D<sub>3</sub>.

<sup>‡</sup> A source of arachidonic acid (ARA).

<sup>§</sup> A source of docosahexaenoic acid (DHA).

# **POTENTIAL ALLERGENS**

Enfamil® Liquid Human Milk Fortifier Standard Protein contains milk and soy. **CAUTION:** Nutritionally incomplete. To be used only under the supervision of a

#### PRODUCT FORMS

physician.

Product	Format	Unit Size	Approximate Product Yield at Normal Dilution	Shelf Life	Item Number	Availability
Enfamil® Liquid Human Milk Fortifier Standard Protein	Liquid	5.5 fl oz/ 163 mL per bottle	Approx. 32 doses per bottle based on a 24 kcal/ fl oz or 0.81 kcal/mL formulation of 5 mL of liquid HMF + 25 mL of breast milk	15 months	3279585	Hospital only

# PREPARATION OF FEEDINGS

Use as directed by the baby's doctor. Shake gently before using. Do not shake vigorously to minimize product foaming. Foaming is normal with this product. Remove protective seal around cap, remove cap and foil seal. This product is commercially sterile as produced. Use proper aseptic techniques.<sup>1</sup>

A variety of aseptic preparation methods are available including the use of sterile containers, or sterile transfer lids and sterile, single-use, enteral syringes. When using sterile containers, after opening bottle pour into disposable, sterile container and draw up needed volume into a sterile, single-use, enteral syringe. Dispense into measured breast milk and swirl.

Failure to follow these instructions could result in severe harm. Opened bottles and fortified breast milk can spoil quickly. Either use immediately or replace fortifier bottle cap and store in refrigerator at 35-40°F (2-4°C) for no longer than 48 hours. Gently agitate product before each use. After feeding begins, use fortified breast milk within one hour or discard. After 48 hours of opening, discard any unused product. Do not freeze.

The baby's doctor will provide instructions for the desired amount of calories to add.

- **WARNING:** Do not use product that has unusual characteristics.
- WARNING: Do not use a microwave oven to warm the fortified breast milk.
   Serious burns may result.
- WARNING: Not for parenteral (I.V.) use. Do not administer directly—must add to breast milk

Infant and Pediatric Feedings: Guidelines for Preparation of Human Milk and Formula in Health Care Facilities. 3rd ed.; 2018.



# Enfamil® Liquid Human Milk Fortifier High Protein

OD (M) IFANCA

#### INDICATION

Enfamil® Liquid Human Milk Fortifier High Protein is added to expressed breast milk for feeding premature and low-birth-weight infants who often need fortified breast milk to meet their increased caloric protein, fat, and other micronutrient needs. Enfamil Liquid Human Milk Fortifier High Protein increases the levels of protein, energy, calcium, phosphorus, iron, vitamin D and other nutrients, producing a diet more suited to the nutritional needs of the rapidly growing premature infant. It has 4 g of protein per 100 Cal when mixed with preterm human milk at the 24 Calorie per fluid ounce concentration/0.81 kcal/mL concentration.\*

# **NUTRIENTS**

Analysis	Per 5 mL Fortifier only	20 mL Plus 100 mL Preterm Breast Milk (120 mL mixed to 0.81 kcal/mL or 24 Cal/fl oz concentration)†	Per 100 Cal Fortified Preterm Breast Milk when mixed to 0.81 kcal/mL or 24 Cal/fl oz concentration†
Calories, kcal	7.5	97	100
Protein, g	0.56	3.9	4
Fat, g	0.58	5.8	6
Linoleic acid, mg	58	710	730
Carbohydrate, g	0.11	7.7	7.9
Water, g	3.8	104	107
Vitamins/Other Nutri	ents		
Vitamin A, IU	290	1210	1240
Vitamin D, IU	47	200	200
Vitamin E, IU	1.4	6	6.2
Vitamin K, mcg	1.42	7.7	7.9

<sup>†</sup> Calculated using values for preterm human breast milk compiled from multiple published studies; protein value used for preterm human milk is 1.62 g/100 mL.¹





<sup>\*</sup> When mixed as directed at 5 mL HMF + 25 mL preterm human milk.

Analysis	Per 5 mL Fortifier only	20 mL Plus 100 mL Preterm Breast Milk (120 mL mixed to 0.81 kcal/mL or 24 Cal/fl oz concentration)*	Per 100 Cal Fortified Preterm Breast Milk when mixed to 0.81 kcal/mL or 24 Cal/fl oz concentration*
Thiamin (Vitamin $B_1$ ), mcg	46	193	200
Riboflavin (Vitamin B <sub>2</sub> ), mcg	66	290	300
Vitamin B <sub>6</sub> , mcg	35	146	150
Vitamin B <sub>12</sub> , mcg	0.16	0.66	0.68
Niacin, mcg	920	3900	4000
Folic Acid (Folacin), mcg	7.7	34	35
Pantothenic Acid, mcg	230	1150	1190
Biotin, mcg	0.84	3.9	4.1
Vitamin C (Ascorbic Acid), mg	3.8	20	20
Minerals			
Calcium, mg	29	141	145
Phosphorus, mg	15.8	78	80
Magnesium, mg	2	11.3	11.6
Iron, mg	2	11.3	11.6
Zinc, mg	0.24	1.33	1.37
Manganese, mcg	2.5	10.4	10.7
Copper, mcg	15	98	101
Sodium, mg	6.8	55	57
Potassium, mg	11.3	95	98
Chloride, mg	6.9	86	88

<sup>\*</sup> Calculated using values for preterm human breast milk compiled from multiple published studies; protein value used for preterm human milk is 1.62 g/100 mL.<sup>1</sup>

Water, whey protein isolate hydrolysate (milk), medium chain triglycerides (MCT oil), soy oil and less than 2%: Mortierella alpina oil<sup>†</sup>, Schizochytrium Sp oil<sup>‡</sup>, pectin, vitamin A palmitate, vitamin E acetate, vitamin  $D_3$ , vitamin  $K_1$ , thiamin hydrochloride, riboflavin, vitamin  $B_6$  hydrochloride, vitamin  $B_{12}$ , niacinamide,

folic acid, calcium pantothenate, biotin, ascorbic acid, calcium citrate, calcium phosphate, cupric sulfate, sodium citrate, potassium citrate, calcium chloride, ferrous sulfate, magnesium phosphate, zinc sulfate.

- † A source of arachidonic acid (ARA).
- ‡ A source of docosahexaenoic acid (DHA).

#### **POTENTIAL ALLERGENS**

Enfamil® Liquid Human Milk Fortifier High Protein contains milk and soy. **CAUTION:** Nutritionally incomplete. To be used only under the supervision of a physician.

### PRODUCT FORMS

Product	Format	Unit Size	Approximate Product Yield at Normal Dilution	Shelf Life	Item Number	Availability
Enfamil® Liquid Human Milk Fortifier High Protein	Liquid	5.5 fl oz/ 163 mL per bottle	Approx. 32 doses per bottle based on a 24 kcal/ fl oz or 0.81 kcal/mL formulation of 5 mL of liquid HMF + 25 mL of breast milk	15 months	3279712	Hospital only

#### PREPARATION OF FEEDINGS

Use as directed by the baby's doctor. Shake gently before using. Do not shake vigorously to minimize product foaming. Foaming is normal with this product. Remove protective seal around cap, remove cap and foil seal. This product is commercially sterile as produced. Use proper aseptic techniques.<sup>1</sup>

A variety of aseptic preparation methods are available including the use of sterile containers, or sterile transfer lids and sterile, single-use, enteral syringes. When using sterile containers, after opening bottle pour into disposable, sterile container and draw up needed volume into a sterile, single-use, enteral syringe. Dispense into measured breast milk and swirl.

Failure to follow these instructions could result in severe harm. Opened bottles and fortified breast milk can spoil quickly. Either use immediately or replace fortifier bottle cap and store in refrigerator at 35-40°F (2-4°C) for no longer than 48 hours. Gently agitate product before each use. After feeding begins, use fortified breast milk within one hour or discard. After 48 hours of opening, discard any unused product. Do not freeze.

Infant and Pediatric Feedings: Guidelines for Preparation of Human Milk and Formula in Health Care Facilities. 3rd ed.; 2018.

The baby's doctor will provide instructions for the desired amount of calories to add.

- WARNING: Do not use product that has unusual characteristics.
- WARNING: Do not use a microwave oven to warm the fortified breast milk.
   Serious burns may result.
- WARNING: Not for parenteral (I.V.) use. Do not administer directly—must add to breast milk



# Enfamil A<sup>+®</sup> Premature High Protein Nursette<sup>®</sup> Bottle

Cow's milk-based, iron-fortified infant formula specifically designed for premature or low birth weight infants that require a higher level of protein.

# **INDICATION**

Cow's milk-based, iron-fortified infant formula specifically designed for premature or low birth weight infants that require a higher level of protein. Contains DHA (a type of Omega-3 fat). For institutional use only.

# **NUTRIENTS**

Analysis	per 100 mL	per 100 kcal			
Energy (kcal/kJ)	81/340	100/420			
Protein (g)	2.8	3.5			
% of total energy	14	14			
Source: Whey protein concentrate, skim m	nilk				
Carbohydrate (g)	8.5	10.5			
% of total energy	42	42			
Source: Corn syrup solids, lactose					
Fat (g)	4.1	5.1			
% of total energy	44	44			
Source: MCT (40%), soy (31%), high oleic sunflower or safflower oils (26.5%), ARA & DHA single cell oil blend (2.5%)					
Linoleic Acid (g)	0.66	0.81			
Linolenic Acid (g)	0.073	0.09			
ARA (mg)	28	34			
DHA (mg)	13.8	17			
Minerals					
Calcium (mg)	134	165			
(mMol)	3.3	4.1			
Phosphorus (mg)	67	83			
(mMol)	2.2	2.7			





Analysis	per 100 mL	per 100 kcal
Magnesium (mg)	7.3	9
(mMol)	0.3	0.37
Iron (mg)	1.46	1.8
Zinc (mg)	1.22	1.5
Manganese (mg)	0.0051	0.0063
Copper (mg)	0.097	0.12
lodine (mg)	0.02	0.025
Sodium (mg) (mMol)	47 2.0	58 2.5
Potassium (mg) (mMol)	80 2.0	98 2.5
Chloride (mg) (mMol)	73 2.1	90 2.5
Vitamins		
Vitamin A (IU)	1020	1250
Vitamin D (IU)	195	240
Vitamin E (IU)	5.1	6.3
Vitamin K (mg)	0.0073	0.009
Vitamin C (mg)	16.2	20
Thiamine (mg)	0.162	0.2
Riboflavin (mg)	0.24	0.3
Niacin (mg)	3.2	4
Pantothenic Acid (mg)	0.97	1.2
Vitamin B <sub>6</sub> (mg)	0.122	0.15
Folic Acid (mg)	0.032	0.04
Vitamin B <sub>12</sub> (mg)	0.0002	0.00025
Biotin (mg)	0.0032	0.004
Choline (mg)	16.2	20
Inositol (mg)	36	44
Carnitine (mg)	1.95	2.4
Taurine (mg)	4.9	6
Nucleotides (mg)	3.4	4.2

Other Characteristics	
Potential Renal Solute Load (m0sm/100 mL) <sup>1</sup>	24
Osmolality (mOsm/kg H <sub>2</sub> O)	300
Osmolarity (mOsm/L)	260
Water (g/100 mL)	88

Use under medical supervision. See Advisory on the product carton. Do not exceed a protein intake of 4.0-4.5 g/kg/d for infants up to 1000 g [3.6-4.1 g/100 kcal) and 3.5-4.0 g/kg/day for infants from 1000-1800 g [3.2-3.6 g/100 kcal). This product could contain up to 3.85 g of protein/100 Kcal. Not intended for feeding premature infants after they reach a weight of approximately 3600 g, unless directed by a physician.

1. Fomon SJ, Ziegler EE. Renal solute load and potential renal solute load in infancy. *J Pediatr*. 1999;134:11-14. Nutrient values are subject to change. Consult actual product labels for most current information.

### **INGREDIENTS**

Water, Modified Milk Ingredients, Corn Syrup Solids, Medium Chain Triglycerides (Fractionated Coconut Oil), Soy Oil, Lactose, High Oleic Vegetable Oil (Sunflower And/Or Safflower Oil), Mono- And Diglycerides, Mortierella Alpina Oil\*, Carrageenan, Crypthecodinium Cohnii Oil\*, Soy Lecithin, Maltodextrin, Taurine, Nucleotides (Adenosine 5'-Monophosphate, Cytidine 5'-Monophosphate, Disodium Guanosine 5'-Monophosphate, Disodium Uridine5'-Monophosphate), L-Carnitine, Ascorbyl Palmitate, Mixed Tocopherols (May Contain Potassium Hydroxide), Minerals (Calcium Carbonate, Calcium Chloride, Calcium Hydroxide, Calcium Phosphate, Cupric Sulfate, Ferrous Sulfate, Magnesium Phosphate, Potassium Citrate, Potassium Iodide, Sodium Chloride, Sodium Citrate And Zinc Sulfate), Vitamins (Biotin, Calcium D-Pantothenate, Choline Chloride, Dl- $\alpha$ -Tocopheryl Acetate, Folic Acid, Inositol, Niacinamide, Pyridoxine Hydrochloride, Riboflavin, Sodium Ascorbate, Thiamine Hydrochloride, Vitamin A Palmitate, Vitamin B<sub>12</sub>, Vitamin D<sub>3</sub> and Vitamin K<sub>1</sub>).

- \* A source of arachidonic acid (ARA).
- t A source of DHA

### **PRODUCT FORMS**

Product	Format	Unit Size	Approximate Product Yield at Normal Dilution	Shelf Life	Item Number	Availability
Enfamil A+® Premature High Protein	Nursette® Bottle	59 mL	59 mL (2 fl oz)	12 Months	1281250	Hospital only

#### PREPARATION OF FEEDINGS

Your baby's health depends on carefully following the instructions below.

Proper hygiene, preparation, use and storage are important when preparing infant formula. Use as directed by your baby's doctor.

**Inspect** each bottle for signs of damage.



1. Wash hands thoroughly with soap and water before preparing bottle for feeding.



2. **SHAKE BOTTLE WELL** and remove cap.



3. Attach **DISPOSABLE NIPPLE UNIT** (not included) or ask your baby's doctor about the need to boil a clean nipple in water before use.

Failure to follow these instructions could result in severe harm. Opened bottles can spoil quickly. Either feed immediately or replace cap and store in refrigerator at 2-4 °C (35-40°F) for no longer than 24 hours. Do not use opened bottle if it is unrefrigerated for more than a total of 2 hours. Do not freeze.

After feeding begins, use within one hour or discard.

**WARNING:** Do not use a microwave oven to warm formula. Serious burns may result.

**Storage:** Store unopened bottles at room temperature. Avoid excessive heat and prolonged exposure to light. Do not freeze.

DO NOT USE IF CAP RING IS BROKEN OR MISSING

USE BY EXPIRATION DATE ON CARTON AND BOTTLE



# Enfamil A<sup>+®</sup> Premature 20 kcal Nursette<sup>®</sup> Bottle

Cow's milk-based, iron-fortified infant formula with concentrated levels of protein, vitamins and minerals for premature or low birth weight infants.

O D (M) IFANCA

## INDICATION

Cow's milk-based, iron-fortified infant formula with concentrated levels of protein, vitamins and minerals for premature or low birth weight infants. Contains DHA (a type of Omega-3 fat).

## **NUTRIENTS**

Analysis	per 100 mL	per 100 kcal			
Energy (kcal/kJ)	68/280	100/420			
Protein (g)	2	3			
% of total energy	12	12			
Source: Whey protein concentrate, skim m	ilk				
Carbohydrate (g)	7.4	11			
% of total energy	44	44			
Source: Corn syrup solids, lactose					
Fat (g)	3.4	5.1			
% of total energy	44	44			
Source: MCT (40%), soy (31%), high oleic sunflower oil (26.5%), ARA & DHA single cell oil blend (2.5%)					
Linoleic Acid (g)	0.55	0.81			
Linolenic Acid (g)	0.061	0.09			
ARA (mg)	23	34			
DHA (mg)	11.5	17			
Minerals					
Calcium (mg)	112	165			
(mMol)	2.8	4.1			



Analysis	per 100 mL	per 100 kcal
Phosphorus (mg)	56	83
(mMol)	1.81	2.7
Magnesium (mg)	6.1	9
(mMol)	0.25	0.37
Iron (mg)	1.22	1.8
Zinc (mg)	1.01	1.5
Manganese (mg)	0.0043	0.0063
Copper (mg)	0.081	0.12
lodine (mg)	0.0169	0.025
Sodium (mg)	39	58
(mMol)	1.71	2.5
Potassium (mg)	66	98
(mMol)	1.7	2.5
Chloride (mg) (mMol)	61 1.71	90 2.5
Vitamins	1.71	2.5
Vitamin A (IU)	050	1050
( )	850	1250
Vitamin D (IU)	162	240
Vitamin E (IU)	4.3	6.3
Vitamin K (mg)	0.0054	0.008
Vitamin C (mg)	13.5	20
Thiamine (mg)	0.135	0.2
Riboflavin (mg)	0.2	0.3
Niacin (mg)	2.7	4
Pantothenic Acid (mg)	0.81	1.2
Vitamin B <sub>6</sub> (mg)	0.101	0.15
Folic Acid (mg)	0.027	0.04
Vitamin B <sub>12</sub> (mg)	0.00017	0.00025
Biotin (mg)	0.0027	0.004
Choline (mg)	13.5	20
Inositol (mg)	30	44
Carnitine (mg)	1.62	2.4
Taurine (mg)	4.1	6
Nucleotides (mg)	2.8	4.2

Other Characteristics	
Potential Renal Solute Load (m0sm/100 mL) <sup>1</sup>	18.4
Osmolality (m0sm/kg H <sub>2</sub> 0)	240
Osmolarity (mOsm/L)	220
Water (g/100 mL)	90

<sup>1.</sup> Fomon SJ, Ziegler EE. Renal solute load and potential renal solute load in infancy. *J Pediatr*. 1999;134:11-14. Nutrient values are subject to change. Consult actual product labels for most current information.

Water, Corn Syrup Solids, Modified Milk Ingredients, Lactose, Medium Chain Triglycerides (Fractionated Coconut Oil), Soy Oil, High Oleic Vegetable Oil (Sunflower And/Or Safflower Oil), Mono And Diglycerides, Mortierella Alpina Oil\*, Crypthecodinium Cohnii Oil†, Carrageenan, Maltodextrin, Taurine, Nucleotides (Adenosine 5'- Monophosphate, Cytidine 5'-Monophosphate, Disodium Guanosine 5'-Monophosphate, Disodium Uridine 5'-Monophosphate), L-Carnitine, Ascorbyl Palmitate, Mixed Tocopherols (May Contain Potassium Hydroxide), Minerals (Calcium Carbonate, Calcium Chloride, Calcium Hydroxide, Calcium Phosphate, Cupric Sulfate, Ferrous Sulfate, Magnesium Phosphate, Potassium Citrate, Potassium Iodide, Sodium Chloride, Sodium Citrate And Zinc Sulfate), Vitamins (Biotin, Calcium D-Pantothenate, Choline Chloride, Dl-α-Tocopheryl, Folic Acid, Inositol, Niacinamide, Pyridoxine Hydrochloride, Riboflavin, Sodium Ascorbate, Thiamine Hydrochloride, Vitamin A Palmitate, Vitamin B<sub>12</sub>, Vitamin D<sub>3</sub> and Vitamin K<sub>1</sub>).

- \* A source of arachidonic acid (ARA).
- † A source of DHA.

# **PRODUCT FORMS**

Product	Format	Unit Size	Approximate Product Yield at Normal Dilution	Shelf Life	Item Number	Availability
Enfamil A <sup>+®</sup> Premature 20 kcal	Nursette® Bottle	59 mL	59 mL (2 fl oz)	12 Months	1250747	Hospital only

#### PREPARATION OF FEEDINGS

Your baby's health depends on carefully following the instructions below.

Proper hygiene, preparation, use and storage are important when preparing infant formula. Use as directed by your baby's doctor.

Inspect each bottle for signs of damage.



1. Wash hands thoroughly with soap and water before preparing bottle for feeding.



2. **SHAKE BOTTLE WELL** and remove cap.



3. Attach **DISPOSABLE NIPPLE UNIT** (not included) or ask your baby's doctor about the need to boil a clean nipple in water before use.

Failure to follow these instructions could result in severe harm. Opened bottles can spoil quickly. Either feed immediately or replace cap and store in refrigerator at 2-4°C (35-40°F) for no longer than 24 hours. Do not use opened bottle if it is unrefrigerated for more than a total of 2 hours. Do not freeze.

After feeding begins, use within one hour or discard.

**WARNING:** Do not use a microwave oven to warm formula. Serious burns may result.

**Storage:** Store unopened bottles at room temperature. Avoid excessive heat and prolonged exposure to light. Do not freeze.

DO NOT USE IF CAP RING IS BROKEN OR MISSING

USE BY EXPIRATION DATE ON CARTON AND BOTTLE



# Enfamil A<sup>+®</sup> Premature 24 kcal Nursette<sup>®</sup> Bottle

Cow's milk-based, iron-fortified infant formula with concentrated levels of energy, protein, vitamins and minerals for premature or low birth weight infants.

OD M IFANCA

# **INDICATION**

Cow's milk-based, iron-fortified infant formula with concentrated levels of energy, protein, vitamins and minerals for premature or low birth weight infants. Contains DHA (a type of Omega-3 fat).

# **NUTRIENTS**

Analysis	per 100 mL	per 100 kcal				
Energy (kcal/kJ)	81/340	100/420				
Protein (g)	2.4	3				
% of total energy	12	12				
Source: Whey protein concentrate, skim m	ilk					
Carbohydrate (g)	8.9	11				
% of total energy	44	44				
Source: Corn syrup solids, lactose						
Fat (g)	4.1	5.1				
% of total energy	44	44				
Source: MCT (40%), soy (31%), high oleic sunflower or safflower oils (26.5%), ARA & DHA single cell oil blend (2.5%)						
Linoleic Acid (g)	0.66	0.81				
Linolenic Acid (g)	0.073	0.09				
ARA (mg)	28	34				
DHA (mg)	13.8	17				
Minerals						
Calcium (mg)	134	165				
(mMol)	3.3	4.1				





Analysis	per 100 mL	per 100 kcal
Phosphorus (mg)	67	83
(mMol)	2.2	2.7
Magnesium (mg)	7.3	9
(mMol)	0.3	0.37
Iron (mg)	1.46	1.8
Zinc (mg)	1.22	1.5
Manganese (mg)	0.0051	0.0063
Copper (mg)	0.097	0.12
lodine (mg)	0.02	0.025
Sodium (mg)	47	58
(mMol)	2.0	2.5
Potassium (mg)	80	98
(mMol)	2.0	2.5
Chloride (mg)	73	90
(mMol)	2.1	2.5
Vitamins		
Vitamin A (IU)	1010	1250
Vitamin D (IU)	195	240
Vitamin E (IU)	5.1	6.3
Vitamin K (mg)	0.0065	0.008
Vitamin C (mg)	16.2	20
Thiamine (mg)	0.162	0.2
Riboflavin (mg)	0.24	0.3
Niacin (mg)	3.2	4
Pantothenic Acid (mg)	0.97	1.2
Vitamin B <sub>6</sub> (mg)	0.122	0.15
Folic Acid (mg)	0.032	0.04
Vitamin B <sub>12</sub> (mg)	0.0002	0.00025
Biotin (mg)	0.0032	0.004
Choline (mg)	16.2	20
Inositol (mg)	36	44
Carnitine (mg)	1.95	2.4

Analysis	per 100 kcal					
Taurine (mg)	Taurine (mg) 4.9					
Nucleotides (mg)	4.2					
Other Characteristics						
Potential Renal Solute Load (m0sm/100 m	22					
Osmolality (mOsm/kg H <sub>2</sub> O)	300					
Osmolarity (mOsm/L)	260					
Water (g/100 mL)		88				

<sup>1.</sup> Fomon SJ, Ziegler EE. Renal solute load and potential renal solute load in infancy. *J Pediatr*. 1999;134:11-14. Nutrient values are subject to change. Consult actual product labels for most current information.

Water, Corn Syrup Solids, Modified Milk Ingredients, Lactose, Medium Chain Triglycerides (Fractionated Coconut Oil), Soy Oil, High Oleic Vegetable Oil (Sunflower and/or Safflower Oil), Mono and Diglycerides, Mortierella Alpina Oil\*, Crypthecodinium Cohnii Oil $^{\dagger}$ , Carrageenan, Maltodextrin, Taurine, Nucleotides (Adenosine 5'- Monophosphate, Cytidine 5'-Monophosphate, Disodium Guanosine 5'-Monophosphate, Disodium Uridine 5'-Monophosphate), L-Carnitine, Ascorbyl Palmitate, Mixed Tocopherols (May contain Potassium Hydroxide), Minerals (Calcium Carbonate, Calcium Chloride, Calcium Hydroxide, Calcium Phosphate, Cupric Sulfate, Ferrous Sulfate, Magnesium Phosphate, Potassium Citrate, Potassium Iodide, Sodium Chloride, Sodium Citrate And Zinc Sulfate), Vitamins (Biotin, Calcium D-Pantothenate, Choline Chloride, Dl- $\alpha$ -Tocopheryl Acetate, Folic Acid, Inositol, Niacinamide, Pyridoxine Hydrochloride, Riboflavin, Sodium Ascorbate, Thiamine Hydrochloride, Vitamin A Palmitate, Vitamin B<sub>12</sub>, Vitamin D<sub>3</sub> And Vitamin K<sub>1</sub>).

- \* A source of arachidonic acid (ARA).
- † A source of DHA.

#### **PRODUCT FORMS**

Product	Format	Unit Size	Approximate Product Yield at Normal Dilution	Shelf Life	Item Number	Availability
Enfamil A <sup>+®</sup> Premature 24 kcal	Nursette® Bottle	59 mL	59 mL (2 fl oz)	12 Months	1215248	Hospital only

#### PREPARATION OF FEEDINGS

Your baby's health depends on carefully following the instructions below.

Proper hygiene, preparation, use and storage are important when preparing infant formula. Use as directed by your baby's doctor.

**Inspect** each bottle for signs of damage.



1. Wash hands thoroughly with soap and water before preparing bottle for feeding.



2. **SHAKE BOTTLE WELL** and remove cap.



3. Attach **DISPOSABLE NIPPLE UNIT** (not included) or ask your baby's doctor about the need to boil a clean nipple in water before use.

Failure to follow these instructions could result in severe harm. Opened bottles can spoil quickly. Either feed immediately or replace cap and store in refrigerator at 2-4 °C (35-40°F) for no longer than 24 hours. Do not use opened bottle if it is unrefrigerated for more than a total of 2 hours. Do not freeze.

After feeding begins, use within one hour or discard.

**WARNING:** Do not use a microwave oven to warm formula. Serious burns may result.

**Storage:** Store unopened bottles at room temperature. Avoid excessive heat and prolonged exposure to light. Do not freeze.

DO NOT USE IF CAP RING IS BROKEN OR MISSING

**USE BY EXPIRATION DATE ON CARTON AND BOTTLE** 



# Enfamil A<sup>+®</sup> EnfaCare®

Cow's milk-based, iron fortified post-discharge infant formula for premature or low birth weight infants.

OD (M) IFANCA

# **INDICATION**

Cow's milk-based, iron fortified post-discharge infant formula for premature or low birth weight infants. Contains DHA (a type of Omega-3 fat).

# **NUTRIENTS**

Analysis	per 100 mL (normal dilution)	per 100 g powder	per 100 kcal			
Energy (kcal/kJ)	74/310	490/2060	100/420			
Protein (g)	2.1	13.8	2.8			
% of total energy	11	11	11			
Source: Whey protein concentrate, skim milk						
Carbohydrate (g)	7.7	51	10.4			
% of total energy	42	42	42			
Source: Lactose, corn syrup solids						
Fat (g)	3.9	26	5.3			
% of total energy	47	47	47			

Source: High oleic vegetable (34%), soy (29%), MCT (20%), coconut (14%) oils, ARA and DHA single cell oil blend (3%)

Linoleic Acid (g)	0.64 (Powder) 0.67 (Nursette® Bottle)*	4.2	0.9
Linolenic Acid (g)	0.07 (Powder) 0.067 (Nursette® Bottle)*	0.47	0.09
ARA (mg)	25	168	34
DHA (mg)	12.6	84	17
Minerals			
Calcium (mg) (mMol - Nursette® Bottle)	89 2.2*	590 -	120 3.0

<sup>\*</sup> Per 100 mL undiluted liquid.





Analysis	per 100 mL (normal dilution)	per 100 g powder	per 100 kcal
Phosphorus (mg)	49	330	66
(mMol - Nursette® Bottle)	1.58*	-	2.1
Magnesium (mg)	5.9	39	8
(mMol - Nursette® Bottle)	0.24*	-	0.33
Iron (mg)	1.33	8.9	1.8
Zinc (mg)	0.74	4.9	1
Manganese (mg)	0.0111	0.074	0.015
Copper (mg)	0.089	0.59	0.12
lodine (mg)	0.0155	0.104	0.021
Selenium (mg)	0.0021	0.0138	0.0028
	27 (Powder)		
Sodium (mg)	26 (Nursette® Bottle)*	182	35
(mMol - Nursette® Bottle)	1.13*	-	1.52
Potassium (mg)	78	520	105
(mMol - Nursette® Bottle)	2.0*	-	2.7
Chloride (mg)	58	380	78
(mMol - Nursette® Bottle)	1.63*	-	2.2
Vitamins			
Vitamin A (IU)	330	2200	450
Vitamin D (IU)	52	350	70
Vitamin E (IU)	3	20	4
Vitamin K (mg)	0.0059	0.039	0.008
Vitamin C (mg)	11.8	79	16
Thiamine (mg)	0.148	0.99	0.2
Riboflavin (mg)	0.148	0.99	0.2
Niacin (mg)	0.74 (Powder) 1.48 (Nursette® Bottle)*	4.9	1 (Powder) 2 (Nursette® Bottle)*
Pantothenic Acid (mg)	0.63	4.2	0.85

<sup>\*</sup> Per 100 mL undiluted liquid.

Analysis	per 100 mL (normal dilution)	per 100 g powder	per 100 kcal
Vitamin B <sub>6</sub> (mg)	0.044 (Powder) 0.074 (Nursette® Bottle)*	0.3	0.06 (Powder) 0.1 (Nursette® Bottle)*
Folic Acid (mg)	0.0192	0.128	0.026
Vitamin B <sub>12</sub> (mg)	0.00022	0.00148	0.0003
Biotin (mg)	0.0037 (Powder) 0.0044 (Nursette® Bottle)*	0.025	0.005 (Powder) 0.006 (Nursette® Bottle)*
Choline (mg)	17.8	118	24
Inositol (mg)	22	148	30
Carnitine (mg)	1.48	9.9	2
Taurine (mg)	4.4	30	6
Nucleotides (mg)	3.1	21	4.2

Other Characteristics	
Renal Solute Load (m0sm/100 mL) <sup>1</sup>	18.4 (Powder) 18.3 (Nursette® Bottle)
Osmolality (mOsm/kg H <sub>2</sub> O)	310 (Powder) 250 (Nursette® Bottle)
Osmolarity (m0sm/L)	280 (Powder) 220 (Nursette® Bottle)
Water (g/100 mL)	89

<sup>\*</sup> Per 100 mL undiluted liquid.

Powdered formulas are not sterile. "To minimize the risk of infection, powdered formulas are not recommended for use in premature or immunocompromised infants unless no appropriate nutritional alternative is available, and then only with strict medical supervision and careful preparation, storage and use."<sup>2</sup>

- Fomon SJ, Ziegler EE. Renal solute load and potential renal solute load in infancy. J Pediatr. 1999; 134:11-14.
- 2. International Formula Council. Infant Feeding: Safety Issues for Health Care Professionals. Atlanta, GA: International Formula Council; 2004:4.

Nutrient values are subject to change. Consult actual product labels for most current information.

#### **POWDER**

Modified Milk Ingredients, Corn Syrup Solids, Lactose, High Oleic Vegetable Oil (Sunflower and/or Safflower), Soy Oil, Medium Chain Triglycerides (Fractionated Coconut Oil), Coconut Oil, Mortierella Alpina Oil\*, Crypthecodinium Cohnii Oil†, Maltodextrin, Taurine, Nucleotides (Adenosine 5'-Monophosphate, Cytidine 5'- Monophosphate, Disodium Guanosine 5'-Monophosphate, Disodium Uridine 5'- Monophosphate), L-Carnitine, Ascorbyl Palmitate, Mixed Tocopherols, (May Contain Potassium Hydroxide), Minerals (Calcium Carbonate, Calcium Phosphate, Cupric Sulfate, Ferrous Sulfate, Magnesium Chloride, Manganese Sulfate, Potassium Citrate, Potassium Iodide, Sodium Chloride, Sodium Selenite And Zinc Sulfate), Vitamins (Ascorbic Acid, Biotin, Calcium D-Pantothenate, Choline Chloride, Dl-α-Tocopheryl Acetate, Folic Acid, Inositol, Niacinamide, Pyridoxine Hydrochloride, Riboflavin, Thiamine Hydrochloride, Vitamin A Palmitate, Vitamin B<sub>12</sub>, Vitamin D<sub>3</sub> and Vitamin K<sub>1</sub>).

- \* A source of arachidonic acid (ARA).
- † A source of DHA.

#### 59 mL NURSETTE® BOTTLE

Water, Maltodextrin, Modified Milk Ingredients, Lactose, High Oleic Vegetable Oil (Sunflower and/or Safflower Oil), Soy Oil, Medium Chain Triglycerides (Fractionated Coconut Oil), Coconut Oil, Mono- and Diglycerides, Mortierella Alpina Oil\*, Soy Lecithin, Crypthecodinium Cohnii Oil\*, Carrageenan, Corn Syrup Solids, Taurine, Nucleotides (Adenosine 5'-Monophosphate, Cytidine 5'-Monophosphate, Disodium Guanosine 5'-Monophosphate, Disodium Uridine 5'-Monophosphate), L-Carnitine, Ascorbyl Palmitate, Mixed Tocopherols, (May Contain Potassium Hydroxide), **Minerals** (Calcium Chloride, Calcium Citrate, Calcium Phosphate, Cupric Sulfate, Ferrous Sulfate, Magnesium Chloride, Manganese Sulfate, Potassium Citrate, Potassium Iodide, Sodium Citrate, Sodium Selenite And Zinc Sulfate), **Vitamins** (Ascorbic Acid, Biotin, Calcium D-Pantothenate, Choline Chloride, Dl- $\alpha$ -Tocopheryl Acetate, Folic Acid, Inositol, Niacinamide, Pyridoxine Hydrochloride, Riboflavin, Sodium Ascorbate, Thiamine Hydrochloride, Vitamin A Palmitate, Vitamin B<sub>12</sub>, Vitamin D<sub>3</sub> and Vitamin K<sub>1</sub>).

- \* A source of arachidonic acid (ARA).
- † A source of DHA.

# **PRODUCT FORMS**

Product	Format	Unit Size	Approximate Product Yield at Normal Dilution	Shelf Life	Item Number	Availability
Enfamil A <sup>+®</sup> Enfacare	Powder	363 g	2400 mL (13.5 bottles x 6 fl oz)	18 Months	1171320	Hospital + Retail
Enfamil A <sup>+®</sup> Enfacare	Nursette® Bottle	59 mL	59 mL (2 fl oz)	12 Months	1215387	Hospital + Enfamil.ca (only)

# PREPARATION OF FEEDINGS

#### **POWDER**

Your baby's health depends on carefully following the instructions below.

Proper hygiene, preparation, dilution, use, and storage are important when preparing infant formula. Powdered infant formulas are not sterile and should not be fed to premature infants or infants who might have immune problems unless directed and supervised by your baby's doctor. Ask your baby's doctor which formula is appropriate for your baby.



1. Wash hands thoroughly with soap and water before preparing formula. Boil clean bottles, nipples, caps and utensils in water (2 minutes at a rolling boil).



2. Boil fresh water at a rolling boil for 2 minutes. Cool to room temperature prior to mixing.



3. Pour desired amount of the cooled water into the bottle. Add powder. Cap bottle and **SHAKE WELL**.

Use the chart below for the correct amounts of cooled, boiled water and powder. Use scoop in can to measure powder. Store **DRY** scoop in this can.

To Make	Boiled Water	Powder
60 mL bottle	<b>60 mL</b> (1/4 cup)	1 unpacked level scoop
120 mL bottle	<b>120 mL</b> (1/2 cup)	2 unpacked level scoops
180 mL bottle	<b>180 mL</b> (3/4 cup)	3 unpacked level scoops
240 mL bottle	<b>240 mL</b> (1 cup)	4 unpacked level scoops

**WARNING:** Do not use a microwave oven to warm formula. Serious burns may result.

Failure to follow these instructions could result in severe harm. Once prepared, infant formula can spoil quickly. Either feed immediately or cover and store in refrigerator at 2-4°C (35-40°F) for no longer than 24 hours. Do not use prepared formula if it is unrefrigerated for more than a total of 2 hours. Do not freeze prepared formula. After feeding begins, use formula within one hour or discard.

**Storage/Handling:** Store powder at room temperature. After opening and between uses, keep tightly covered, store in dry area and use contents within 1 month. Do not freeze powder and avoid excessive heat.

#### **NURSETTE® BOTTLES**

Your baby's health depends on carefully following the instructions below.

Proper hygiene, preparation, use and storage are important when preparing infant formula. Use as directed by your baby's doctor. **Inspect** each bottle for signs of damage.



1. Wash hands thoroughly with soap and water before preparing bottle for feeding.



2. **SHAKE BOTTLE WELL** and remove cap.



3. Attach **DISPOSABLE NIPPLE UNIT** (not included) or ask your baby's doctor about the need to boil a clean nipple in water before use.

**WARNING:** Do not use a microwave oven to warm formula. Serious burns may result.

Failure to follow these instructions could result in severe harm. Opened bottles and prepared bottles can spoil quickly. Either feed immediately or replace cap and store in refrigerator at 2-4°C (35-40°F) for no longer than 24 hours. Do not use opened bottle or prepared bottle if it is unrefrigerated for more than a total of 2 hours. Do not freeze formula. After feeding begins, use formula within one hour or discard.

**Storage:** Store unopened bottles at room temperature. Avoid excessive heat and prolonged exposure to light. Do not freeze.

DO NOT USE IF CAP RING IS BROKEN OR MISSING

USE BY EXPIRATION DATE ON CAN, CARTON, AND/OR BOTTLE



Date published 11/2024 – Please reference Mead Johnson Nutrition's online Product Handbook for the most up to date product details (www.enfamil.ca/medical).



# Enfamil A+® NeuroPro™ Infant

OD (M) IFANCA

# INDICATION

Enfamil A+® NeuroPro™ Infant is a milk-based infant formula designed for term infants (0-12 months) with an expert-recommended level of DHA,\*† our exclusive blend‡ including 2'-FL for immune support, and naturally occurring MFGM components.§

- \* World Health Organization recommended amount of DHA is 0.2%-0.36% of total fatty acids. Enfamil® has 0.32% of total fatty acids as DHA, similar to the average amount found in world-wide mature breast milk.
- † DHA supports normal physical brain development.
- ± HuMO6 Blend™ of Polydextrose, GOS, 2'-FL, Vit C & E and Selenium.
- § From whey protein concentrate.

Enfamil A<sup>+®</sup> NeuroPro™ Infant is available through Health Canada's Interim Policy on the importation and sale of infant formulas.

### PRODUCT FORMATS AND FORMS:

Item Number	Product Description	Availability	Format	Approximate Product Yield at Normal Dilution
3290179	Enfamil A+® NeuroPro™ Infant Hospital Nursette® 59 ML (8X6 BTL) CA	Hospital Only	Liquid, Nursette	59 mL (2 fl oz)
3276799	Enfamil A+® NeuroPro™ Infant Powder 587 G (6 BIT) CA	Hospital & Retail	Powder	4381 mL (18 bottles x 8 fl oz)
3285831	Enfamil A+® NeuroPro™ Infant Ready to Use (RTU) 237 ML (4X6 BTL) CA	Hospital & Retail	Liquid, RTF	237 mL (1 bottle x 8 fl oz)
3285832	Enfamil A+® NeuroPro™ Infant Ready to Use (RTU) 237 ML (18 BTL) CA	Hospital & Retail	Liquid, RTF	237 mL (1 bottle x 8 fl oz)

# **NUTRIENTS**

Nutrients (Normal Dilution)	Per 100 mL	Per 100 g Powder
Calories	68	510
Protein, g	1.4	11
Fat, g	3.6	27
Carbohydrate, g	7.6	57
Water, g	90	2.4
Linoleic Acid, g	0.5	3.9
Linolenic Acid, g	0.05	0.4
Arachidonic Acid, mg	17	126
Docosahexaenoic Acid, mg	12 (11 RTF)	86
Vitamin A, IU	203	1515
Vitamin D, IU	47	354
Vitamin E, IU	1.4	10
Vitamin K, mg	0.0061	0.045
Thiamin (Vitamin B <sub>1</sub> ), mg	0.054	0.4
Riboflavin (Vitamin B <sub>2</sub> ), mg	0.095	0.71
Vitamin B <sub>6</sub> , mg	0.041	0.3
Vitamin B <sub>12</sub> , mg	0.0002	0.0015
Niacin, mg	0.68	5.1
Folic Acid (Folacin), mg	0.011	0.081
Pantothenic Acid, mg	0.34	2.5
Biotin, mg	0.002	0.015
Vitamin C (Ascorbic acid), mg	8.1	61
Choline, mg	16	121
Inositol, mg	16	121
L-Carnitine, mg (not declared on label)	1.4	10
Taurine, mg (not declared on label)	4.1	30

Nutrients (Normal Dilution)	Per 100 mL	Per 100 g Powder
Calcium, mg	53	394
Phosphorus, mg	29	217
Magnesium, mg	5.4	40
Iron, mg	1.2	9.1
Zinc, mg	0.68	5.1
Manganese, mg	0.0101	0.076
Copper, mg	0.051	0.38
lodine, mg	0.0101	0.076
Selenium, mg	0.0019	0.014
Sodium, mg (mEq)	18 (0.79)	136 (5.9)
Potassium, mg (mEq)	73 (1.87)	545 (13.9)
Chloride, mg (mEq)	43 (1.2)	318 (9)

Source of Protein: nonfat milk and whey protein concentrate

Source of Fat: Palm olein, soy, coconut, high oleic sunflower oils, and single cell oil blend of DHA and ARA Source of Carbohydrate: Lactose, GOS, PDX

Product nutrient values and ingredients are subject to change. Please see product label for current information.

# **NUTRIENT FACTS**

Nutrient Density	Powered Prepared at Standard Dilution of 0.68 kcal/mL (20 Calories/fl oz)	RTF Prepared at Standard Dilution of 0.68 kcal/mL (20 Calories/fl oz)
Protein (% Calories)	8.5	8.5
Whey: Casein Ratio	65:35	65:35
Fat (% Calories)	48	48
Carbohydrate (% Calories)	43.5	43.5
Potential Renal Solute Load (m0sm/100 Calories) <sup>1</sup>	19.1	19.1
Potential Renal Solute Load (m0sm/100 mL) <sup>1</sup>	12.8	12.8

Nutrient Density	Powered Prepared at Standard Dilution of 0.68 kcal/mL (20 Calories/fl oz)	RTF Prepared at Standard Dilution of 0.68 kcal/mL (20 Calories/fl oz)	
Osmolality (mOsm/kg water)	300	300	
Osmolarity (mOsm/L)	270	270	
Lactose-free	No	No	

<sup>1.</sup> Fomon SJ, Ziegler EE, Renal solute load and potential renal solute load in infancy, J Pediatr, 1999:134:11-14.

#### ENFAMIL A+® NEUROPRO™ INFANT POWDER:

Lactose, Contains three or more of the following (palm olein oil, coconut oil, soy oil, high oleic sunflower oil), Modified milk ingredients\*, Galactooligosaccharide solids (GOS), Polydextrose, Soy lecithin, Mortierella alpina oil $^{\dagger}$ , Corn syrup solids, Schizochytrium sp. oil $^{\dagger}$ , 2'-fucosyllactose, Taurine, L-carnitine, Mixed tocopherols, Ascorbyl palmitate, (may contain potassium hydroxide), **Minerals** (calcium carbonate, calcium phosphate, cupric sulfate, ferrous sulfate, magnesium oxide, manganese sulfate, potassium chloride, potassium citrate, potassium iodide, sodium chloride, sodium selenite and zinc sulfate), **Vitamins** (ascorbic acid, biotin, calcium Dpantothenate, choline chloride, dl- $\alpha$ -tocopheryl acetate, folic acid, inositol, niacinamide, pyridoxine hydrochloride, riboflavin, thiamine hydrochloride, vitamin A palmitate, vitamin B<sub>12</sub>, vitamin D<sub>3</sub> and vitamin K<sub>1</sub>).

# ENFAMIL A+® NEUROPRO™ INFANT READY TO USE (NURSETTE® BOTTLES AND RTU BOTTLES):

Water, Lactose, Contains three or more of the following (palm olein oil, coconut oil, soy oil, high oleic sunflower oil), Modified milk ingredients\*, Polydextrose, Galactooligosaccharide solids (GOS), Mono- and diglycerides, Mortierella alpina oil<sup>†</sup>, Soy lecithin, Carrageenan, 2'-fucosyllactose, Schizochytrium sp. oil<sup>‡</sup>, Corn syrup solids, Taurine, L-carnitine, Mixed tocopherols, Ascorbyl palmitate, (may contain potassium hydroxide), **Minerals** (calcium carbonate, calcium chloride, calcium phosphate, cupric sulfate, ferrous sulfate, magnesium phosphate, manganese sulfate, potassium citrate, potassium iodide, sodium chloride, sodium citrate, sodium selenite and zinc sulfate), **Vitamins** (ascorbic acid, biotin, calcium Dpantothenate, choline chloride, dl- $\alpha$ -tocopheryl acetate, folic acid, inositol, niacinamide, pyridoxine hydrochloride, riboflavin, sodium ascorbate, thiamine hydrochloride, vitamin A palmitate, vitamin B<sub>12</sub>, vitamin D<sub>3</sub> and vitamin K<sub>1</sub>).

#### POTENTIAL ALLERGENS

Enfamil A+® NeuroPro™ Infant contains milk and soy.

<sup>\*</sup> A source of MFGM (Milk Fat Globule Membrane composed of proteins and fats) components.

<sup>†</sup> A source of ARA.

<sup>‡</sup> A source of DHA.

# PREPARATION OF FEEDINGS

#### **POWDER**

Your baby's health depends on carefully following the instructions below.

Proper hygiene, preparation, dilution, use, and storage are important when preparing infant formula. Powdered infant formulas are not sterile and should not be fed to premature infants or infants who might have immune problems unless directed and supervised by your baby's doctor. Ask your baby's doctor which formula is appropriate for your baby.



1. Wash hands thoroughly with soap and water before preparing formula. Boil clean bottles, nipples, caps and utensils in water (2 minutes at a rolling boil).



 Boil fresh water at a rolling boil for
 minutes. Cool to room temperature prior to mixing.



3. Pour desired amount of the cooled water into the bottle. Add powder. Cap bottle and **SHAKE WELL**.

Use the chart below for the correct amounts of cooled, boiled water and powder. Use scoop in container to measure powder. Store **DRY** scoop in this tub.

To Make	Boiled Water	Powder
<b>60 mL</b> bottle	<b>60 mL</b> (1/4 cup)	1 unpacked level scoop
120 mL bottle	<b>120 mL</b> (1/2 cup)	2 unpacked level scoops
180 mL bottle	<b>180 mL</b> (3/4 cup)	3 unpacked level scoops
240 mL bottle	<b>240 mL</b> (1 cup)	4 unpacked level scoops

**WARNING:** Do not use a microwave oven to warm formula. Serious burns may result.

Failure to follow these instructions could result in severe harm. Once prepared, infant formula can spoil quickly. Either feed immediately or cover and store in refrigerator at 2-4°C (35-40°F) for no longer than 24 hours. Do not use prepared formula if it is unrefrigerated for more than a total of 2 hours. Do not freeze prepared formula. After feeding begins, use formula within one hour or discard.

**Storage/Handling:** Store powder at room temperature. After opening and between uses, keep tightly covered, store in dry area and use contents within 1 month. Do not freeze powder and avoid excessive heat.

#### **NURSETTE® BOTTLES**

Your baby's health depends on carefully following the instructions below.

Proper hygiene, preparation, use and storage are important when preparing infant formula. Use as directed by your baby's doctor. **Inspect** each bottle for signs of damage.



1. Wash hands thoroughly with soap and water before preparing bottle for feeding.



2. **SHAKE BOTTLE WELL** and remove cap.



3. Attach **DISPOSABLE NIPPLE UNIT** (not included) or ask your baby's doctor about the need to boil a clean nipple in water before use.

**WARNING:** Do not use a microwave oven to warm formula. Serious burns may result.

Failure to follow these instructions could result in severe harm. Opened bottles and prepared bottles can spoil quickly. Either feed immediately or replace cap and store in refrigerator at 2-4°C (35-40°F) for no longer than 24 hours. Do not use opened bottle or prepared bottle if it is unrefrigerated for more than a total of 2 hours. Do not freeze formula. After feeding begins, use formula within one hour or discard.

**Storage:** Store unopened bottles at room temperature. Avoid excessive heat and prolonged exposure to light. Do not freeze.

DO NOT USE IF CAP RING IS BROKEN OR MISSING

USE BY EXPIRATION DATE ON TUB, CARTON, AND/OR BOTTLE

#### **READY TO FEED**

Your baby's health depends on carefully following the instructions below.

Proper hygiene, preparation, use and storage are important when preparing infant formula. Use as directed by your baby's doctor. **Inspect** each bottle for signs of damage.



1. Wash hands thoroughly with soap and water before preparing bottle for feeding.



 SHAKE BOTTLE WELL, remove protective seal around cap, remove cap and foil seal.



3. Attach **DISPOSABLE NIPPLE UNIT** (not included) or ask your baby's doctor about the need to boil a clean nipple in water before use.

**WARNING:** Do not use a microwave oven to warm formula. Serious burns may result.

Failure to follow these instructions could result in severe harm. Opened bottles and prepared bottles can spoil quickly. Either feed immediately or replace cap and store in refrigerator at 2-4°C (35-40°F) for no longer than 48 hours. Do not use opened bottle or prepared bottle if it is unrefrigerated for more than a total of 2 hours. Do not freeze formula. After feeding begins, use formula within one hour or discard.

**Storage:** Store unopened bottles at room temperature. Avoid excessive heat and prolonged exposure to light. Do not freeze.



# Enfamil A<sup>+®</sup> Hypercaloric Nursette<sup>®</sup> Bottle

Cow's milk-based, iron fortified hypercaloric infant formula for infants with high energy needs or requiring fluid restriction.

OD (M) IFANCA

# INDICATION

Cow's milk-based, iron fortified hypercaloric infant formula for infants with high energy needs or requiring fluid restriction. Contains DHA (a type of Omega-3 fat).

# **NUTRIENTS**

Analysis	per 100 mL		
Energy (kcal/kJ)	81/340		
Protein (g)	1.7		
% of total energy	8.5		
Source: Whey protein concentrate, nonfat milk			
Carbohydrate (g)	8.9		
% of total energy	43.5		
Source: Lactose			
Fat (g)	4.3		
% of total energy	48		
Source: Palm olein (44%), soy (19.5%), coconut (19.5%), high oleic sunflower (14.5%) oils, ARA & DHA single cell oil blend (2.5%)			
Linoleic Acid (g)	0.7		
Linolenic Acid (g)	0.06		
ARA (mg)	28		
DHA (mg)	14		
Minerals			
Calcium (mg)	64		
(mMol)	1.58		
Phosphorus (mg)	35		
(mMol)	1.13		





Analysis	per 100 mL		
Magnesium (mg)	6.5		
(mMol)	0.27		
Iron (mg)	1.47		
Zinc (mg)	0.81		
Manganese (mg)	0.0122		
Copper (mg)	0.061		
lodine (mg)	0.0122		
Selenium (mg)	0.0023		
Sodium (mg)	22		
(mMol)	0.95		
Potassium (mg)	88		
(mMol)	2.2		
Chloride (mg)	51		
(mMol)	1.44		
Vitamins			
Vitamin A (IU)	240		
Vitamin D (IU)	49		
Vitamin E (IU)	1.63		
Vitamin K (mg)	0.0073		
Vitamin C (mg)	9.8		
Thiamine (mg)	0.065		
Riboflavin (mg)	0.114		
Niacin (mg)	0.81		
Pantothenic Acid (mg)	0.41		
Vitamin B <sub>6</sub> (mg)	0.049		
Folic Acid (mg)	0.013		
Vitamin B <sub>12</sub> (mg)	0.00024		
Biotin (mg)	0.0024		
Choline (mg)	20		
Inositol (mg)	4.9		
Carnitine (mg)	1.63		
Taurine (mg)	4.9		
Nucleotides (mg)	3.4		

Other Characteristics			
Potential Renal Solute Load (m0sm/100 mL) <sup>1</sup>	15.4		
Osmolality (mOsm/kg $H_2O$ )	360		
Osmolarity (mOsm/L)	320		
Water (g/100 mL)	88		

<sup>1.</sup> Fomon SJ, Ziegler EE. Renal solute load and potential renal solute load in infancy. *J Pediatr.* 1999;134:11-14. Nutrient values are subject to change. Consult actual product labels for most current information.

Water, Lactose, Modified Milk Ingredients, Palm Olein Oil, Soy Oil, Coconut Oil, High Oleic Sunflower Oil, Mono- And Diglycerides, Soy Lecithin, Mortierella Alpina Oil\*, Carrageenan, Crypthecodinium Cohnii Oil $^{\dagger}$ , Taurine, Maltodextrin, Nucleotides (Adenosine 5'-Monophosphate, Cytidine 5'-Monophosphate, Disodium Guanosine 5'-Monophosphate, Disodium Uridine 5'-Monophosphate), Corn Syrup Solids, L-Carnitine, Ascorbyl Palmitate, Mixed Tocopherols, (May Contain Potassium Hydroxide), **Minerals** (Calcium Carbonate, Calcium Chloride, Calcium Phosphate, Cupric Sulfate, Ferrous Sulfate, Magnesium Phosphate, Manganese Sulfate, Potassium Citrate, Potassium Iodide, Sodium Chloride, Sodium Citrate, Sodium Selenite And Zinc Sulfate), **Vitamins** (Ascorbic Acid, Biotin, Calcium D-Pantothenate, Choline Chloride, DL- $\alpha$ -Tocopheryl Acetate, Folic Acid, Inositol, Niacinamide, Pyridoxine Hydrochloride, Riboflavin, Sodium Ascorbate, Thiamine Hydrochloride, Vitamin A Palmitate, Vitamin B<sub>12</sub>, Vitamin D<sub>3</sub> and Vitamin K<sub>1</sub>).

#### PRODUCT FORMS

Product	Format	Unit Size	Approximate Product Yield at Normal Dilution	Shelf Life	Item Number	Availability
Enfamil A <sup>+®</sup>	Nursette®	59	59 mL	12	1210915	Hospital
Hypercaloric	Bottle	mL	(2 fl oz)	Months		only

<sup>\*</sup> A source of arachidonic acid (ARA).

<sup>†</sup> A source of DHA.

#### PREPARATION OF FEEDINGS

Your baby's health depends on carefully following the instructions below.

Proper hygiene, preparation, use and storage are important when preparing infant formula. Use as directed by your baby's doctor.

Inspect each bottle for signs of damage.



1. Wash hands thoroughly with soap and water before preparing bottle for feeding.



2. **SHAKE BOTTLE WELL** and remove cap.



3. Attach **DISPOSABLE NIPPLE UNIT** (not included) or ask your baby's doctor about the need to boil a clean nipple in water before use.

Failure to follow these instructions could result in severe harm. Opened bottles can spoil quickly. Either feed immediately or replace cap and store in refrigerator at 2-4°C (35-40°F) for no longer than 24 hours. Do not use opened bottle if it is unrefrigerated for more than a total of 2 hours. Do not freeze. After feeding begins, use within one hour or discard.

**WARNING:** Do not use a microwave oven to warm formula. Serious burns may result.

**Storage:** Store unopened bottles at room temperature. Avoid excessive heat and prolonged exposure to light. Do not freeze.

DO NOT USE IF CAP RING IS BROKEN OR MISSING

**USE BY EXPIRATION DATE ON CARTON AND BOTTLE** 



### **Enfamil®**

Cow's milk-based, iron fortified, routine starter formula for healthy term infants.



#### **INDICATION**

Cow's milk-based, iron fortified, routine starter formula for healthy term infants.

Analysis	per 100 mL	per 100 g		
Factory (1,221/1/1)	(normal dilution)	powder		
Energy (kcal/kJ)	68/280	510/2130		
Protein (g)	1.42	10.8		
% of total energy	8.5	8.5		
Source: Nonfat milk, whey protein cor	ncentrate			
Carbohydrate (g)	7.4	56		
% of total energy	43.5	43.5		
Source: Lactose				
Fat (g)	3.6	27		
% of total energy	48	48		
Source: Palm olein (45%), soy (20%), coconut (20%), high oleic sunflower (15%) oils				
Linoleic Acid (g)	0.58	4.4		
Linolenic Acid (g)	0.054	0.41		
Minerals				
Calcium (mg)	53	400		
Phosphorus (mg)	29	220		
Magnesium (mg)	5.4	41		
Iron (mg)	1.22	9.2		
Zinc (mg)	0.68	5.1		
Manganese (mg)	0.0101	0.077		
Copper (mg)	0.051	0.38		





Analysis	per 100 mL (normal dilution)	per 100 g powder		
lodine (mg)	0.0101	0.077		
Selenium (mg)	0.00189	0.0143		
Sodium (mg)	18.3	138		
Potassium (mg)	73	550		
Chloride (mg)	43	320		
Vitamins				
Vitamin A (IU)	200	1540		
Vitamin D (IU)	41	310		
Vitamin E (IU)	1.35	10.2		
Vitamin K (mg)	0.0061	0.046		
Vitamin C (mg)	8.1	61		
Thiamine (mg)	0.054	0.41		
Riboflavin (mg)	0.095	0.72		
Niacin (mg)	0.68	5.1		
Pantothenic Acid (mg)	0.34	2.6		
Vitamin B <sub>6</sub> (mg)	0.041	0.31		
Folic Acid (mg)	0.0108	0.082		
Vitamin B <sub>12</sub> (mg)	0.0002	0.00154		
Biotin (mg)	0.002	0.0154		
Choline (mg)	16.2	123		
Inositol (mg)	4.1	31		
Carnitine (mg)	1.35	10.2		
Taurine (mg)	4.1	31		
Nucleotides (mg)	2.8	22		
Other Characteristics				
Potential Renal Solute Load (m0sm/1	12.9			
Osmolality (mOsm/kg H <sub>2</sub> O)	300			
Osmolarity (mOsm/L)	270			
Water (g/100 mL)		90		

Powdered formulas are not sterile. "To minimize the risk of infection, powdered formulas are not recommended for use in premature or immunocompromised infants unless no appropriate nutritional alternative is available, and then only with strict medical supervision and careful preparation, storage and use."<sup>2</sup>

- 1. Fomon SJ, Ziegler EE. Renal solute load and potential renal solute load in infancy. J Pediatr. 1999;134:11-14.
- 2. International Formula Council. Infant Feeding: Safety Issues for Health Care Professionals. Atlanta, GA: International Formula Council; 2004:4.

Nutrient values are subject to change. Consult actual product labels for most current information.

#### **INGREDIENTS**

Lactose, Modified Milk Ingredients, Palm Olein Oil, Coconut Oil, Soy Oil, High Oleic Sunflower Oil, Soy Lecithin, Corn Syrup Solids, Maltodextrin, Taurine, Nucleotides (Adenosine 5'-Monophosphate, Cytidine 5'-Monophosphate, Disodium Guanosine 5'-Monophosphate, Disodium Uridine 5'-Monophosphate), L-Carnitine, (May Contain Potassium Hydroxide), **Minerals** (Calcium Carbonate, Cupric Sulfate, Ferrous Sulfate, Magnesium Oxide, Manganese Sulfate, Potassium Chloride, Potassium Citrate, Sodium Citrate, Sodium Selenite And Zinc Sulfate), **Vitamins** (Ascorbic Acid, Biotin, Calcium D-Pantothenate, Choline Chloride, DL- $\alpha$ -Tocopheryl Acetate, Folic Acid, Inositol, Niacinamide, Pyridoxine Hydrochloride, Riboflavin, Thiamine Hydrochloride, Vitamin A Palmitate, Vitamin B<sub>12</sub>, Vitamin D<sub>3</sub> and Vitamin K<sub>1</sub>).

#### PRODUCT FORMS

Product	Format	Unit Size	Approximate Product Yield at Normal Dilution	Shelf Life	Item Number	Availability
Enfamil®	Powder	900 g	6900 mL (29 bottles x 8 fl oz)	36 Months	1274015	Retail only

#### PREPARATION OF FEEDINGS

#### Your baby's health depends on carefully following the instructions below.

Proper hygiene, preparation, dilution, use and storage are important when preparing infant formula. Powdered infant formulas are not sterile and should <u>not</u> be fed to premature infants or infants who might have immune problems unless directed and supervised by your baby's doctor. Ask your baby's doctor which formula is appropriate for your baby.



1. Wash hands thoroughly with soap and water before preparing formula. Boil clean bottles, nipples, caps and utensils in water (2 minutes at a rolling boil).



2. Boil fresh water at a rolling boil for 2 minutes. Cool to room temperature prior to mixing.



3. Pour desired amount of the cooled water into the bottle. Add powder. Cap bottle and **SHAKE WELL.** 

Use the chart below for the correct amounts of cooled, boiled water and powder. Use scoop in can to measure powder. Store **DRY** scoop in this can.

To Make	Boiled Water	Powder
60 mL bottle	<b>60 mL</b> (1/4 cup)	1 unpacked level scoop
120 mL bottle	<b>120 mL</b> (1/2 cup)	2 unpacked level scoops
180 mL bottle	<b>180 mL</b> (3/4 cup)	3 unpacked level scoops
240 mL bottle	<b>240 mL</b> (1 cup)	4 unpacked level scoops

**WARNING:** Do not use a microwave oven to warm formula. Serious burns may result.

Failure to follow these instructions could result in severe harm. Once prepared, infant formula can spoil quickly. Either feed immediately or cover and store in refrigerator at  $2-4^{\circ}\text{C}$  ( $35-40^{\circ}\text{F}$ ) for no longer than 24 hours. Do not use prepared formula if it is unrefrigerated for more than a total of 2 hours. Do not freeze prepared formula. After feeding begins, use formula within one hour or discard.

**Storage/Handling:** Store cans at room temperature. After opening and between uses, keep tightly covered, store in dry area and use contents within 1 month. Do not freeze powder and avoid excessive heat.

#### **USE BY EXPIRATION DATE ON BOTTOM OF CAN**



### Enfamil A+® NeuroPro™ Gentlease®

OD (M) IFANCA

#### **INDICATION**

Enfamil A+® NeuroPro™ Gentlease® is a milk-based formula designed to reduce fussiness, crying, and gas in 24 hours.¹ With easy-to-digest partially hydrolyzed proteins,\* it also contains an expert-recommended level of brain-building DHA,¹¹ naturally occurring MFGM components,§ and our exclusive fibre blend# shown to promote soft stools.¹¹² The partially hydrolyzed formula of Enfamil A+® NeuroPro™ Gentlease® has lower lactose levels (1/5th) as compared to routine milk-based formula

- \* Similar to all infant formulas.
- † DHA supports normal physical brain development.
- ‡ World Health Organization recommended amount of DHA is 0.2%-0.36% of total fatty acids. Enfamil® has 0.32% of total fatty acids as DHA, similar to the average amount found in world-wide mature breast milk. § From whey protein concentrate.
- # Blend of GOS, Polydextrose in powder format only.
- ¶ Compared to Enfamil® without fibre blend.
- 1. Berseth et al. study, 2009.
- 2. Ziegler et al. study, 2007.

#### PRODUCT LONG TERM USAGE

Enfamil A+® NeuroPro™ Gentlease® is designed to provide the sole source of nutrition for infants up to age 6 months and provide a major source of nutrition for the remainder of the first year.

Nutrients (Normal Dilution)	Per 100 mL	Per 100 g Powder
Protein, g	1.6	12
Fat, g	3.6	27
Carbohydrate, g	7.3	57
Calories	68	510
Water, g	90	2.1
Linoleic Acid, g	0.5	4
Linolenic Acid, g	0.05	0.4
Arachidonic Acid, mg	17	128





Nutrients (Normal Dilution)	Per 100 mL	Per 100 g Powder
Docosahexaenoic Acid, mg	11	87
Vitamin A, IU	203 (202 RTF)	1530
Vitamin D, IU	47 (44 RTF)	357
Vitamin E, IU	1.4	10
Vitamin K, mg	0.0061	0.046
Thiamin (Vitamin B <sub>1</sub> ), mg	0.054	0.41
Riboflavin (Vitamin B <sub>2</sub> ), mg	0.095	0.71
Vitamin B <sub>6</sub> , mg	0.041	0.31
Vitamin B <sub>12</sub> , mg	0.0002	0.0015
Niacin, mg	0.68	5.1
Folic Acid (folacin), mg	0.011	0.082
Pantothenic Acid, mg	0.34	2.6
Biotin, mg	0.002	0.015
Vitamin C (Ascorbic Acid), mg	8.1	61
Choline, mg	16	122
Inositol, mg	16	122
L-Carnitine, mg	1.4	10
Taurine, mg	4.1	31
Calcium, mg	55	418
Phosphorus, mg	31	235
Magnesium, mg	5.4	41
Iron, mg	1.2	9.2
Zinc, mg	0.68	5.1
Manganese, mg	0.0101	0.077
Copper, mg	0.051	0.38
lodine, mg	0.0101	0.077
Selenium, mg	0.0019	0.014

Nutrients (Normal Dilution)	Per 100 mL	Per 100 g Powder
Sodium, mg	27	204
Potassium, mg	73	551
Chloride, mg	43	321

Source of Protein: Partially hydrolyzed nonfat milk and whey protein concentrate Source of Fat: Palm olein, soy, coconut, high oleic sunflower oils, and single cell oil blend of DHA and ARA Source of Carbohydrate: Powder - Corn syrup solids, lactose; RTU - Corn syrup solids, lactose, rice starch Product nutrient values and ingredients are subject to change. Please see product label for current information.

#### **NUTRIENT FACTS**

Nutrient Density	Powered Prepared at Standard Dilution of 0.68 kcal/mL (20 Calories/fl oz)
Protein (% Calories)	9
Fat (% Calories)	48
Carbohydrate (% Calories)	43
Potential Renal Solute Load (m0sm/100 Calories)¹	21
Potential Renal Solute Load (m0sm/100 mL)¹	14.4
Osmolality (mOsm/kg water)	200
Osmolarity (mOsm/L)	180
Lactose-free	No

<sup>1.</sup> Fomon SJ, Ziegler EE. Renal solute load and potential renal solute load in infancy. J Pediatr. 1999;134:11-14.

#### **INGREDIENTS**

#### ENFAMIL A+® NEUROPRO™ GENTLEASE® POWDER:

Corn syrup solids, Partially hydrolyzed modified milk ingredients (soy)\*, Contains three or more of the following (palm olein oil, coconut oil, soy oil, high oleic sunflower oil), Galactooligosaccharide solids (GOS), Polydextrose, Soy lecithin, Lactose, Mortierella alpina oil\*, Schizochytrium sp. oil\*, 2'- fucosyllactose, Taurine, L-carnitine, Mixed tocopherols, Ascorbyl palmitate, (may contain potassium hydroxide), Minerals (calcium carbonate, calcium phosphate, cupric sulfate, ferrous sulfate, magnesium phosphate, manganese sulfate, potassium chloride, potassium iodide, sodium citrate, sodium hydroxide, sodium selenite and zinc sulfate), Vitamins (ascorbic acid, biotin, calcium D-pantothenate, choline

chloride,  $dl-\alpha$ -tocopheryl acetate, folic acid, inositol, niacinamide, pyridoxine hydrochloride, riboflavin, thiamine hydrochloride, vitamin A palmitate, vitamin  $B_{12}$ , vitamin  $D_3$  and vitamin  $K_1$ ).

#### ENFAMIL A+® NEUROPRO™ GENTLEASE® LIQUID (RTF):

Water, Corn syrup solids, Partially hydrolyzed modified milk ingredients (soy)\*, Contains three or more of the following (palm olein oil, coconut oil, soy oil, high oleic sunflower oil), Rice starch, Acetylated tartaric acid esters of mono- and diglycerides, Mortierella alpina oil†, 2'-fucosyllactose, Schizochytrium sp. oil‡, Soy Lecithin, Taurine, L-carnitine, Lactose, Mixed tocopherols, Ascorbyl palmitate, (may contain potassium hydroxide), **Minerals** (calcium carbonate, calcium phosphate, cupric sulfate, ferrous sulfate, magnesium phosphate, manganese sulfate, potassium chloride, potassium citrate, potassium iodide, sodium chloride, sodium citrate, sodium hydroxide, sodium selenite and zinc sulfate), **Vitamins** (ascorbic acid, biotin, calcium D-pantothenate, choline chloride, dl- $\alpha$ -tocopheryl acetate, folic acid, inositol, niacinamide, pyridoxine hydrochloride, riboflavin, thiamine hydrochloride, vitamin A palmitate, vitamin B<sub>12</sub>, vitamin D<sub>3</sub> and vitamin K<sub>1</sub>).

Potential Allergens: Enfamil A+® NeuroPro™ Gentlease® contains milk and soy.

- \* A source of MFGM (Milk Fat Globule Membrane composed of proteins and fats) components.
- † A source of ARA.
- ‡ A source of DHA.

#### PRODUCT FORMS

Item	Description	Availability	Format	Approximate Product Yield at Normal Dilution
3286052	Enfamil A+® NeuroPro™ Gentlease® Powder 553 G (4 BIT) CA	Hospital & Retail	Powder	Not available as NeuroPro™ formulation yet
3276297	Enfamil A+® NeuroPro™ Gentlease® Ready to Feed (RTF) 237 ML (4X6 BTL) CA	Hospital & Retail	Liquid, RTF	237 mL

#### PREPARATION OF FEEDINGS

#### POWDER

Your baby's health depends on carefully following the instructions below.

Proper hygiene, preparation, dilution, use, and storage are important when preparing infant formula. Powdered infant formulas are not sterile and should not be fed to premature infants or infants who might have immune problems unless directed and supervised by your baby's doctor. Ask your baby's doctor which formula is appropriate for your baby.



1. Wash hands thoroughly with soap and water before preparing formula. Boil clean bottles, nipples, caps and utensils in water (2 minutes at a rolling boil).



 Boil fresh water at a rolling boil for
 minutes. Cool to room temperature prior to mixing.



3. Pour desired amount of the cooled water into the bottle. Add powder. Cap bottle and **SHAKE WELL**.

Use the chart below for the correct amounts of cooled, boiled water and powder. Use scoop in container to measure powder. Store **DRY** scoop in this tub.

To Make	Boiled Water	Powder
<b>60 mL</b> bottle	<b>60 mL</b> (1/4 cup)	1 unpacked level scoop
120 mL bottle	<b>120 mL</b> (1/2 cup)	2 unpacked level scoops
180 mL bottle	<b>180 mL</b> (3/4 cup)	3 unpacked level scoops
240 mL bottle	<b>240 mL</b> (1 cup)	4 unpacked level scoops

**WARNING:** Do not use a microwave oven to warm formula. Serious burns may result.

Failure to follow these instructions could result in severe harm. Once prepared, infant formula can spoil quickly. Either feed immediately or cover and store in refrigerator at 2-4°C (35-40°F) for no longer than 24 hours. Do not use prepared formula if it is unrefrigerated for more than a total of 2 hours. Do not freeze prepared formula. After feeding begins, use formula within one hour or discard.

**Storage/Handling:** Store powder at room temperature. After opening and between uses, keep tightly covered, store in dry area and use contents within 1 month. Do not freeze powder and avoid excessive heat.

#### **NURSETTE® BOTTLES**

Your baby's health depends on carefully following the instructions below.

Proper hygiene, preparation, use and storage are important when preparing infant formula. Use as directed by your baby's doctor. **Inspect** each bottle for signs of damage.



1. Wash hands thoroughly with soap and water before preparing bottle for feeding.



2. **SHAKE BOTTLE WELL** and remove cap.



3. Attach **DISPOSABLE NIPPLE UNIT** (not included) or ask your baby's doctor about the need to boil a clean nipple in water before use.

**WARNING:** Do not use a microwave oven to warm formula. Serious burns may result.

Failure to follow these instructions could result in severe harm. Opened bottles and prepared bottles can spoil quickly. Either feed immediately or replace cap and store in refrigerator at 2-4°C (35-40°F) for no longer than 24 hours. Do not use opened bottle or prepared bottle if it is unrefrigerated for more than a total of 2 hours. Do not freeze formula. After feeding begins, use formula within one hour or discard.

**Storage:** Store unopened bottles at room temperature. Avoid excessive heat and prolonged exposure to light. Do not freeze.

DO NOT USE IF CAP RING IS BROKEN OR MISSING

USE BY EXPIRATION DATE ON TUB, CARTON AND/OR BOTTLE

#### **READY TO FEED**

Your baby's health depends on carefully following the instructions below.

Proper hygiene, preparation, use and storage are important when preparing infant formula. Use as directed by your baby's doctor. **Inspect** each bottle for signs of damage.



1. Wash hands thoroughly with soap and water before preparing bottle for feeding.



2. SHAKE BOTTLE WELL, remove protective seal around cap, remove cap and foil seal.



3. Attach **DISPOSABLE NIPPLE UNIT** (not included) or ask your baby's doctor about the need to boil a clean nipple in water before use.

**WARNING:** Do not use a microwave oven to warm formula. Serious burns may result.

Failure to follow these instructions could result in severe harm. Opened bottles and prepared bottles can spoil quickly. Either feed immediately or replace cap and store in refrigerator at 2-4°C (35-40°F) for no longer than 48 hours. Do not use opened bottle or prepared bottle if it is unrefrigerated for more than a total of 2 hours. Do not freeze formula. After feeding begins, use formula within one hour or discard.

**Storage:** Store unopened bottles at room temperature. Avoid excessive heat and prolonged exposure to light. Do not freeze.



# Enfamil A<sup>+®</sup> For feeding babies who frequently Spit Up\*

Cow's milk-based, iron fortified infant formula with added rice starch for infants who spit up frequently or who otherwise need a thickened formula.

\* For babies who spit up more than 4 times per day.

(U) D (M) IFANCA

#### INDICATION

Cow's milk-based, iron fortified infant formula with added rice starch for infants who spit up frequently or who otherwise need a thickened formula. Contains two dietary fibres (galactooligosaccharides and polydextrose) and DHA (a type of Omega-3 fat).

Analysis	per 100 mL (normal dilution)	per 100 g		
Energy (kcal/kJ)	68/280	500/2090		
Protein (g)	1.7	12		
% of total energy	10	10		
Source: Nonfat milk				
Carbohydrate (g)	7.6	56		
% of total energy	44	44		
Galactooligosaccharides (g)	0.2	1.5		
Polydextrose (g)	0.2	1.5		
Source: Lactose, rice starch, maltodextrin, galactooligosaccharides, polydextrose				
Fat (g)	3.4	25		
% of total energy	46	46		
Source: Palm olein (44%), soy (19.5%), coconut (19.5%), high oleic sunflower (14.5%) oils, ARA & DHA single cell oil blend (2.5%)				
Linoleic Acid (g)	0.5	3.9		
Linolenic Acid (g)	0.05	0.35		
ARA (mg)	23	169		
DHA (mg)	11	85		





Analysis	per 100 mL (normal dilution)	per 100 g
Minerals		
Calcium (mg)	53	390
Phosphorus (mg)	36	260
Magnesium (mg)	5.4	40
Iron (mg)	1.22	9
Zinc (mg)	0.68	5
Manganese (mg)	0.0101	0.075
Copper (mg)	0.051	0.37
lodine (mg)	0.0101	0.075
Selenium (mg)	0.00189	0.0139
Sodium (mg)	27	200
Potassium (mg)	73	540
Chloride (mg)	51	370
Vitamins		
Vitamin A (IU)	200	1490
Vitamin D (IU)	41	300
Vitamin E (IU)	1.35	9.9
Vitamin K (mg)	0.0061	0.045
Vitamin C (mg)	8.1	60
Thiamine (mg)	0.054	0.4
Riboflavin (mg)	0.095	0.7
Niacin (mg)	0.68	5
Pantothenic Acid (mg)	0.34	2.5
Vitamin B <sub>6</sub> (mg)	0.041	0.3
Folic Acid (mg)	0.0108	0.08
Vitamin B <sub>12</sub> (mg)	0.0002	0.00149
Biotin (mg)	0.002	0.0149
Choline (mg)	16.2	119
Inositol (mg)	4.1	30
Carnitine (mg)	1.35	9.9
Taurine (mg)	4.1	30

Other Characteristics	
Potential Renal Solute Load (m0sm/100 mL) <sup>1</sup>	15.3
Osmolality (mOsm/kg H <sub>2</sub> O)	230
Osmolarity (mOsm/L)	210
Water (g/100 mL)	90

Powdered formulas are not sterile. "To minimize the risk of infection, powdered formulas are not recommended for use in premature or immunocompromised infants unless no appropriate nutritional alternative is available, and then only with strict medical supervision and careful preparation, storage and use." 2

- 1. Fomon SJ, Ziegler EE. Renal solute load and potential renal solute load in infancy. J Pediatr. 1999;134:11-14.
- International Formula Council. Infant Feeding: Safety Issues for Health Care Professionals. Atlanta, GA: International Formula Council; 2004:4.

Nutrient values are subject to change. Consult actual product labels for most current information.

#### **INGREDIENTS**

Milk ingredients, Contains three or more of the following (palm olein oil, coconut oil, soy oil, high oleic sunflower oil), Rice starch, Lactose, Maltodextrin, Polydextrose, Galactooligosaccharide solids (GOS),Mortierella alpina oil\*, Corn syrup solids, Crypthecodinium cohnii oil†, Taurine, L-carnitine, Ascorbyl palmitate, Mixed tocopherols, (may contain potassium hydroxide), **Minerals** (calcium carbonate, calcium phosphate, cupric sulfate, ferrous sulfate, manganese sulfate, potassium iodide, sodium citrate, sodium selenite and zinc sulfate), **Vitamins** (ascorbic acid, biotin, calcium D-pantothenate, choline chloride, dl- $\alpha$ -tocopheryl acetate, folic acid, inositol, niacinamide, pyridoxine hydrochloride, riboflavin, thiamine hydrochloride, vitamin A palmitate, vitamin B<sub>12</sub>, vitamin D<sub>3</sub> and vitamin K<sub>1</sub>).

- \* A source of ARA.
- † A source of DHA

#### **PRODUCT FORMS**

Product	Format	Unit Size	Approximate Product Yield at Normal Dilution	Shelf Life	Item Number	Availability
Enfamil A+® For feeding babies who frequently Spit Up‡	Powder	629 g	5000 mL (21 bottles x 8 fl oz)	18 Months	3271466	Hospital, Retail, Online

<sup>‡</sup> For babies who spit up more than 4 times per day.

#### PREPARATION OF FEEDINGS

Your baby's health depends on carefully following the instructions below.

Proper hygiene, preparation, dilution, use and storage are important when preparing infant formula. Powdered infant formulas are not sterile and should

**<u>not</u>** be fed to premature infants or infants who might have immune problems unless directed and supervised by your baby's doctor. Ask your baby's doctor which formula is appropriate for your baby.

Carefully follow preparation instructions on the label. LET BOTTLE SIT FOR 5 MINUTES PRIOR TO FEEDING.



1. Wash hands thoroughly with soap and water before preparing formula. Boil clean bottles, nipples, caps and utensils in water (2 minutes at a rolling boil).



2. Boil fresh water at a rolling boil for 2 minutes. Cool to room temperature prior to mixing.



3. Pour desired amount of the cooled water into the bottle. Add powder. Cap bottle and **SHAKE WELL**. LET BOTTLE SIT 5 MINUTES.

Use the chart below for the correct amounts of cooled, boiled water and powder. Use scoop in tub to measure powder. Store **DRY** scoop in this tub.

To Make	Boiled Water	Powder
<b>60 mL</b> bottle	<b>60 mL</b> (1/4 cup)	1 unpacked level scoop
120 mL bottle	<b>120 mL</b> (1/2 cup)	2 unpacked level scoops
180 mL bottle	<b>180 mL</b> (3/4 cup)	3 unpacked level scoops
240 mL bottle	<b>240 mL</b> (1 cup)	4 unpacked level scoops

**WARNING:** Do not use a microwave oven to warm formula. Serious burns may result.

Failure to follow these instructions could result in severe harm. Once prepared, infant formula can spoil quickly. Either feed immediately or cover and store in refrigerator at  $2-4^{\circ}\text{C}$  ( $35-40^{\circ}\text{F}$ ) for no longer than 24 hours. Do not use prepared formula if it is unrefrigerated for more than a total of 2 hours. Do not freeze prepared formula. After feeding begins, use formula within one hour or discard.

**Storage/Handling:** Store powder at room temperature; avoid extreme temperatures. After opening and between uses, keep pouch and tub lid tightly closed, store in a dry area, and use contents within 1 month. Do not store loose powder in tub. Use tub with Enfamil A<sup>+®</sup> For feeding babies who frequently Spit Up formula only. Keep powder fresh and prevent bacterial growth by ensuring tub is clean and completely dry. Completely empty tub and wipe with a clean, dry cloth before reusing. Each pouch has a new batch code and expiration date sticker that must be retained until the contents of the pouch have been consumed.



# Enfamil A<sup>+®</sup> Lactose Free

Cow's milk-based, iron fortified lactose-free infant formula designed for infants sensitive to lactose.

① D (M) IFANCA

#### INDICATION

Cow's milk-based, iron fortified lactose-free infant formula designed for infants sensitive to lactose. Contains DHA (a type of Omega-3 fat). Not intended for use by infants or children with galactosemia.

Analysis	per 100 mL (normal dilution) and per 100 mL RTF		
Energy (kcal/kJ)	68/280		
Protein (g)	1.42 (Nursette® Bottle) 1.4 (Conc)		
% of total energy	8.4 (Nursette® Bottle) 8.5 (Conc)		
Source: Milk protein isolate			
Carbohydrate (g)	7.4		
% of total energy	43.5 (Nursette® Bottle) 44 (Conc)		
Source: Corn syrup solids			
Fat (g)	3.6		
% of total energy	48		
Source: Palm olein (44%), soy (19.5%), coconut (14.5%) oils, ARA & DHA single cell oil blend (2.			
Linoleic Acid (g)	0.54 (Nursette® Bottle) 0.6 (Conc)		
Linolenic Acid (g)	0.051 (Nursette® Bottle) 0.05 (Conc)		
ARA (mg)	23 (Nursette® Bottle) 17 (Conc)		
DHA (mg)	11.5 (Nursette® Bottle) 12 (Conc)		



Analysis	per 100 mL (normal dilution) and per 100 mL RTF
Minerals	
Calcium (mg)	55 (Nursette® Bottle) 56 (Conc)
(mMol - Nursette® Bottle)	1.38
Phosphorus (mg) (mMol - Nursette® Bottle)	31 1.0
Magnesium (mg) (mMol - Nursette® Bottle)	5.4 0.22
Iron (mg)	1.22 (Nursette® Bottle) 1.2 (Conc)
Zinc (mg)	0.68
Manganese (mg)	0.0101 (Nursette® Bottle) 0.0102 (Conc)
Copper (mg)	0.051
lodine (mg)	0.0101 (Nursette® Bottle) 0.0102 (Conc)
Selenium (mg)	0.00189 (Nursette® Bottle) 0.0019 (Conc)
Sodium (mg) (mMol - Nursette® Bottle)	20 0.88
Potassium (mg)	74 (Nursette® Bottle)
(mMol - Nursette® Bottle)	75 (Conc) 1.9
Chloride (mg)	45 (Nursette® Bottle)
(mMol - Nursette® Bottle)	46 (Conc) 1.28
Vitamins	
Vitamin A (IU)	200 (Nursette® Bottle) 204 (Conc)
Vitamin D (IU)	41
Vitamin E (IU)	1.35 (Nursette® Bottle) 1.4 (Conc)
Vitamin K (mg)	0.0061
Vitamin C (mg)	8.1 (Nursette® Bottle) 8.2 (Conc)
Thiamine (mg)	0.054

Analysis	per 100 mL (normal dilution) and per 100 mL RTF
Riboflavin (mg)	0.095
Niacin (mg)	0.68
Pantothenic Acid (mg)	0.34
Vitamin B <sub>6</sub> (mg)	0.041
Folic Acid (mg)	0.0108 (Nursette® Bottle) 0.011(Conc)
Vitamin B <sub>12</sub> (mg)	0.0002
Biotin (mg)	0.002
Choline (mg)	16.2 (Nursette® Bottle) 16 (Conc)
Inositol (mg)	4.1
Carnitine (mg)	1.35 (Nursette® Bottle) 1.4 (Conc)
Taurine (mg)	4.1
Nucleotides (mg)	2.8 (Nursette® Bottle) 0 (Conc)

Other Characteristics	
Potential Renal Solute Load (m0sm/100 mL) <sup>1</sup>	13.2 (Nursette® Bottle) 13.1 (Conc)
Osmolality (mOsm/kg H <sub>2</sub> O)	200 (Nursette® Bottle) 300 (Conc)
Osmolarity (mOsm/L)	182 (Nursette® Bottle) 270 (Conc)
Water (g/100 mL)	91 (Nursette® Bottle) 90 (Conc)

<sup>1.</sup> Fomon SJ, Ziegler EE. Renal solute load and potential renal solute load in infancy. *J Pediatr*. 1999;134:11-14. Nutrient values are subject to change. Consult actual product labels for most current information.

#### **INGREDIENTS**

#### CONCENTRATE

Water, Corn syrup solids, Contains three or more of the following (palm olein oil, coconut oil, soy oil, high oleic sunflower oil), Milk protein isolate, Soy lecithin, Mono- and diglycerides, Mortierella alpina oil\*, Schizochytrium sp. oil\*, Carrageenan, Taurine, L-carnitine, Maltodextrin, Ascorbyl palmitate, Mixed tocopherols, (may contain potassium hydroxide), **Minerals** (calcium carbonate, calcium phosphate, cupric sulfate, ferrous sulfate, magnesium phosphate,

manganese sulfate, potassium chloride, potassium citrate, potassium iodide, sodium citrate, sodium selenite and zinc sulfate), **Vitamins** (ascorbic acid, biotin, calcium D-pantothenate, choline chloride, dl- $\alpha$ -tocopheryl acetate, folic acid, inositol, niacinamide, pyridoxine hydrochloride, riboflavin, thiamine hydrochloride, vitamin A palmitate, vitamin B<sub>12</sub>, vitamin D<sub>3</sub> and vitamin K<sub>1</sub>).

#### 59 mL NURSETTE® BOTTLE

Water, Corn Syrup Solids, Milk Protein Isolate, Palm Olein Oil, Soy Oil, Coconut Oil, High Oleic Sunflower Oil, Mono- And Diglycerides, Soy Lecithin, Mortierella Alpina Oil\*, Crypthecodinium Cohnii Oil\*, Carrageenan, Maltodextrin, Taurine, Nucleotides (Adenosine 5'-Monophosphate, Cytidine 5'-Monophosphate, Disodium Guanosine 5'-Monophosphate, Disodium Uridine 5'-Monophosphate), L- Carnitine, Ascorbyl Palmitate, Mixed Tocopherols, (May Contain Potassium Hydroxide), Minerals (Calcium Carbonate, Calcium Phosphate, Cupric Sulfate, Ferrous Sulfate, Magnesium Chloride, Manganese Sulfate, Potassium Chloride, Potassium Citrate, Potassium Iodide, Sodium Citrate, Sodium Selenite And Zinc Sulfate), Vitamins (Ascorbic Acid, Biotin, Calcium D-Pantothenate, Choline Chloride, DL- $\alpha$ -Tocopheryl Acetate, Folic Acid, Inositol, Niacinamide, Pyridoxine Hydrochloride, Riboflavin, Sodium Ascorbate, Thiamine Hydrochloride, Vitamin A Palmitate, Vitamin B<sub>12</sub>, Vitamin D<sub>3</sub> and Vitamin K<sub>1</sub>).

#### **PRODUCT FORMS**

Product	Format	Unit Size	Approximate Product Yield at Normal Dilution	Shelf Life	Item Number	Availability
Enfamil A <sup>+®</sup> Lactose Free	Concentrate	385 mL	770 mL (3 bottles x 8 fl oz)	12 Months	3278010	Hospital & Retail
	Nursette® Bottle	59 mL	59 mL (2 fl oz)	12 Months	1271623	Hospital only

#### PREPARATION OF FEEDINGS

#### CONCENTRATE

Your baby's health depends on carefully following the instructions below.

Proper hygiene, preparation, dilution, use and storage are important when preparing infant formula. Use as directed by your baby's doctor.

<sup>\*</sup> A source of ARA.

<sup>†</sup> A source of DHA.

<sup>\*</sup> A source of ARA.

<sup>†</sup> A source of DHA.



1. Wash hands thoroughly with soap and water before preparing formula. Boil clean bottles, nipples, caps, can opener and utensils in water [2 minutes at a rolling hoil].



2. Boil fresh water at a rolling boil for 2 minutes. Cool to room temperature prior to mixing. Clean top of can with previously boiled water, **SHAKE WELL** and open.



3. Pour desired amount of the cooled water into the bottle. Add an **equal** amount of concentrated liquid. Cap bottle and **SHAKE WELL**.

**WARNING:** Do not use a microwave oven to warm formula. Serious burns may result.

Failure to follow these instructions could result in severe harm. Opened cans and prepared formula can spoil quickly. Either feed immediately or cover and store in refrigerator at 2-4°C (35-40°F) for no longer than 48 hours. Do not use opened can and/or prepared formula if they are unrefrigerated for more than a total of 2 hours. Do not freeze formula. After feeding begins, use formula within one hour or discard.

**Storage:** Store unopened cans at room temperature. Avoid excessive heat. Do not freeze.

#### **NURSETTE® BOTTLES**

Your baby's health depends on carefully following the instructions below. Proper hygiene, preparation, use and storage are important when preparing infant formula. Use as directed by your baby's doctor.

Inspect each bottle for signs of damage.



1. Wash hands thoroughly with soap and water before preparing bottle for feeding.



2. **SHAKE BOTTLE WELL** and remove cap.



3. Attach **DISPOSABLE NIPPLE UNIT** (not included) or ask your baby's doctor about the need to boil a clean nipple in water before use.

Failure to follow these instructions could result in severe harm. Opened bottles can spoil quickly. Either feed immediately or replace cap and store in refrigerator at 2-4°C (35-40°F) for no longer than 24 hours. Do not use opened bottle if it is unrefrigerated for more than a total of 2 hours. Do not freeze. After feeding begins, use formula within one hour or discard.

**WARNING:** Do not use a microwave oven to warm formula. Serious burns may result.

**Storage:** Store unopened bottles at room temperature. Avoid excessive heat and prolonged exposure to light. Do not freeze.

DO NOT USE IF CAP RING IS BROKEN OR MISSING

USE BY EXPIRATION DATE ON CAN, CARTON AND/OR BOTTLE



# Enfamil A+® Soy

Soy-based, iron fortified lactose-free infant formula designed for infants who need a milk-free formula due to health or cultural reasons.

O Pareve manufactured on dairy equipment (M) IFANCA

#### **INDICATION**

Soy-based, iron fortified lactose-free infant formula designed for infants who need a milk-free formula due to health or cultural reasons. Contains DHA (a type of Omega-3 fat).

Analysis	per 100 mL (normal dilution)
Energy (kcal/kJ)	68/280
Protein (g)	1.7
% of total energy	10
Source: Soy protein isolate (with added L-methi	onine)
Carbohydrate (g)	7.2
% of total energy	42
Source: Corn syrup solids	
Fat (g)	3.6
% of total energy	48
Source: Palm olein (44%), soy (19.5%), coconut (14.5%) oils, ARA & DHA single cell oil blend (2.	
Linoleic Acid (g)	0.6
Linolenic Acid (g)	0.05 (Conc) 0.054 (Nursette® Bottle)
ARA (mg)	17 (Conc) 23 (Nursette® Bottle)
DHA (mg)	12 (Conc) 11.5 (Nursette® Bottle)
Minerals	
Calcium (mg)	71
(mMol - Nursette® Bottle)	1.77



Analysis	per 100 mL (normal dilution)
Phosphorus (mg) (mMol - Nursette® Bottle)	47 1.51
Magnesium (mg) (mMol - Nursette® Bottle)	5.4 0.22
Iron (mg)	1.2
Zinc (mg)	0.81
Manganese (mg)	0.017 (Conc) 0.0169 (Nursette® Bottle)
Copper (mg)	0.051
lodine (mg)	0.0101
Selenium (mg)	0.0019
Sodium (mg) (mMol - Nursette® Bottle)	24 1.06
Potassium (mg) (mMol - Nursette® Bottle)	81 2.1
Chloride (mg) (mMol - Nursette® Bottle)	54 1.52
Vitamins	
Vitamin A (IU)	200
Vitamin D (IU)	41
Vitamin E (IU)	1.4 (Conc) 1.35 (Nursette® Bottle)
Vitamin K (mg)	0.0061
Vitamin C (mg)	8.1
Thiamine (mg)	0.054
Riboflavin (mg)	0.061
Niacin (mg)	0.68
Pantothenic Acid (mg)	0.34
Vitamin B <sub>6</sub> (mg)	0.041
Folic Acid (mg)	0.011 (Conc) 0.0108 (Nursette® Bottle)
Vitamin B <sub>12</sub> (mg)	0.0002
Biotin (mg)	0.002

Analysis	per 100 mL (normal dilution)
Choline (mg)	16 (Conc) 16.2 (Nursette® Bottle)
Inositol (mg)	4.1
Carnitine (mg)	1.4 (Conc) 1.35 (Nursette® Bottle)
Taurine (mg)	4.1
Other Characteristics	
Potential Renal Solute Load (m0sm/100 mL) <sup>1</sup>	15.9 (Conc) 15.6 (Nursette® Bottle)
Osmolality (m0sm/kg H <sub>2</sub> 0)	170 (Conc) 200 (Nursette® Bottle)
Osmolarity (mOsm/L)	153 (Conc) 180 (Nursette® Bottle)
Water (g/100 mL)	90

<sup>1.</sup> Fomon SJ, Ziegler EE. Renal solute load and potential renal solute load in infancy. *J Pediatr*. 1999;134:11-14. Nutrient values are subject to change. Consult actual product labels for most current information.

#### **INGREDIENTS**

#### CONCENTRATE

Water, Corn syrup solids, Contains three or more of the following (palm olein oil, coconut oil, soy oil, high oleic sunflower oil), Soy protein isolate, Soy lecithin, Mono- and diglycerides, Mortierella alpina oil\*, Schizochytrium sp. oil†, L- methionine, Taurine, Carrageenan, L-carnitine, Maltodextrin, Ascorbyl palmitate, Mixed tocopherols, (may contain potassium hydroxide), **Minerals** (calcium carbonate, calcium phosphate, cupric sulfate, ferrous sulfate, magnesium chloride, magnesium phosphate, manganese sulfate, potassium chloride, potassium citrate, sodium selenite and zinc sulfate), **Vitamins** (ascorbic acid, biotin, calcium D-pantothenate, choline chloride, dl- $\alpha$ -tocopheryl acetate, folic acid, inositol, niacinamide, pyridoxine hydrochloride, riboflavin, thiamine hydrochloride, vitamin A palmitate, vitamin B<sub>12</sub>, vitamin D<sub>3</sub> and vitamin K<sub>1</sub>).

#### 59 mL NURSETTE® BOTTLE

Water, Corn Syrup Solids, Soy Protein Isolate, Palm Olein Oil, Soy Oil, Coconut Oil, High Oleic Sunflower Oil, Soy Lecithin, Mono- And Diglycerides, Mortierella Alpina Oil\*, L-Methionine, Carrageenan, Crypthecodinium Cohnii Oil\*, Taurine,

<sup>\*</sup> A source of ARA.

<sup>†</sup> A source of DHA.

L-Carnitine, Ascorbyl Palmitate, Mixed Tocopherols, (May Contain Potassium Hydroxide), Minerals (Calcium Carbonate, Calcium Phosphate, Cupric Sulfate, Ferrous Sulfate, Magnesium Chloride, Magnesium Phosphate, Potassium Chloride, Potassium Citrate, Potassium Iodide, Sodium Chloride, Sodium Citrate, Sodium Selenite And Zinc Sulfate), Vitamins (Ascorbic Acid, Biotin, Calcium D-Pantothenate, Choline Chloride, DL- $\alpha$ -Tocopheryl Acetate, Folic Acid, Inositol, Niacinamide, Pyridoxine Hydrochloride, Riboflavin, Thiamine Hydrochloride, vitamin A Palmitate, vitamin B12, vitamin D3 and vitamin K1).


#### **PRODUCT FORMS**

Product	Format	Unit Size	Approximate Product Yield at Normal Dilution	Shelf Life	Item Number	Availability
Enfamil A <sup>+®</sup> Soy	Concentrate	385 mL	770 mL (3 bottles x 8 fl oz)	15 Months	3276942	Hospital & Online
	Nursette® Bottle	59 mL	59 mL (2 fl oz)	12 Months	1271716	Hospital only

#### PREPARATION OF FEEDINGS

#### CONCENTRATE

Your baby's health depends on carefully following the instructions below. Proper hygiene, preparation, dilution, use and storage are important when preparing infant formula. Use as directed by your baby's doctor.



**WARNING:** Do not use a microwave oven to warm formula. Serious burns may result.

<sup>\*</sup> A source of ARA.

<sup>†</sup> A source of DHA.

Failure to follow these instructions could result in severe harm. Opened cans and prepared formula can spoil quickly. Either feed immediately or cover and store in refrigerator at  $2-4^{\circ}\text{C}$  (35-40°F) for no longer than 48 hours. Do not use opened can and/or prepared formula if it is unrefrigerated for more than a total of 2 hours. Do not freeze formula. After feeding begins, use formula within one hour or discard.

**Storage:** Store unopened cans at room temperature. Avoid excessive heat. Do not freeze.

#### **NURSETTE® BOTTLES**

Your baby's health depends on carefully following the instructions below.

Proper hygiene, preparation, use and storage are important when preparing infant formula. Use as directed by your baby's doctor.

**Inspect** each bottle for signs of damage.



Wash hands thoroughly with soap and water before preparing bottle for feeding.



2. **SHAKE BOTTLE WELL** and remove cap.



3. Attach **DISPOSABLE NIPPLE UNIT** (not included) or ask your baby's doctor about the need to boil a clean nipple in water before use.

Failure to follow these instructions could result in severe harm. Opened bottles can spoil quickly. Either feed immediately or replace cap and store in refrigerator at 2-4°C (35-40°F) for no longer than 24 hours. Do not use opened bottle if it is unrefrigerated for more than a total of 2 hours. Do not freeze.

After feeding begins, use within one hour or discard.

**WARNING:** Do not use a microwave oven to warm formula. Serious burns may result.

**Storage:** Store unopened bottles at room temperature. Avoid excessive heat and prolonged exposure to light. Do not freeze formula. After feeding begins, use formula within one hour or discard.

DO NOT USE IF CAP RING IS BROKEN OR MISSING

USE BY EXPIRATION DATE ON CAN, CARTON AND/OR BOTTLE



# Enfamil A<sup>+®</sup> Serenity

Enfamil A+® Serenity is designed to promote soft stools.

O D (M) IFANCA

#### INDICATION

Enfamil A+® Serenity is designed to promote soft stools.

#### **BENEFITS**

A fibre blend (polydextrose & GOS) clinically shown to promote soft stools.\* Lower in iron than Enfamil  $A^{+0,+}$  Contains a milk protein blend, patterned after breast milk, that that has been partially broken down.‡ Contains DHA, a type of Omega-3 fat and an important building block of the brain. Complete nutrition including brain-building DHA. Lower in lactose than Enfamil  $A^{+0,\$}$  The reusable tub has a snap-in scoop which is easy to find and easy to use with less mess.

- \* Studied in Enfamil A+® infant formula compared to same formula without fibre blend.
- † 17% lower iron as compared to Enfamil A+®.
- ‡ Based on whey:casein ratio of typical mature breast milk (15 days to 6 months after birth).
- § 1/2 of the lactose compared to Enfamil A+®.

Analysis	per 100 mL (normal dilution)	per 100 g powder
Energy (kcal/kJ)	68/280	510/2130
Protein (g)	1.56	11.7
Carbohydrate (g)	7.5	56
Galactooligosaccharides (g)	0.2	1.5
Polydextrose (g)	0.2	1.5
Fat (g)	3.6	27
Linoleic Acid (g)	0.5	4
Linolenic Acid (g)	0.05	0.37
Arachidonic Acid (ARA) (mg)	23	173
Docosahexaenoic Acid (DHA) (mg)	11.5	86
Minerals		
Ash (g)	0.4	3
Calcium (mg)	55	420
Phosphorus (mg)	31	230



Analysis	per 100 mL (normal dilution)	per 100 g powder
Magnesium (mg)	5.4	41
Iron (mg)	1.01	7.6
Zinc (mg)	0.67	5.1
Manganese (mg)	0.0101	0.076
Copper (mg)	0.051	0.38
lodine (mg)	0.0101	0.076
Selenium (mg)	0.00189	0.0142
Sodium (mg)	24	183
Potassium (mg)	73	549
Chloride (mg)	42	320
Vitamins		
Vitamin A (IU)	200	1520
Vitamin D (IU)	40	300
Vitamin E (IU)	1.35	10.2
Vitamin K (mg)	0.0061	0.046
Vitamin C (mg)	8.1	61
Thiamine (mg)	0.054	0.41
Riboflavin (mg)	0.094	0.71
Niacin (mg)	0.67	5.1
Pantothenic Acid (mg)	0.34	2.5
Vitamin B <sub>6</sub> (mg)	0.04	0.3
Folic Acid (mg)	0.0108	0.081
Vitamin B <sub>12</sub> (mg)	0.0002	0.00152
Biotin (mg)	0.002	0.0152
Choline (mg)	16.2	122
Inositol (mg)	4	30
Carnitine (mg)	1.35	10.2
Taurine (mg)	4	30

Other Characteristics	
Potential Renal Solute Load (m0sm/100 mL) <sup>1</sup>	14
Osmolality (mOsm/kg H <sub>2</sub> O)	250
Osmolarity (mOsm/L)	230
Water (g/100 mL)	90

Fomon SJ, Ziegler EE. Renal solute load and potential renal solute load in infancy. J Pediatr. 1999;134:11-14.
 Nutrient values are subject to change. Consult actual product labels for most current information.

#### **INGREDIENTS**

Partially Hydrolyzed Modified Milk Ingredients (Soy), Corn Syrup Solids, Lactose, Palm Olein Oil, Coconut Oil, Soy Oil, High Oleic Sunflower Oil, Polydextrose, Galactooligosaccharide Solids (GOS), Mortierella Alpina Oil\*, Crypthecodinium Cohnii Oil\*, Taurine, L-Carnitine, Ascorbyl Palmitate, Mixed Tocopherols, (May Contain Potassium Hydroxide), **Minerals** (Calcium Carbonate, Calcium Phosphate, Cupric Sulfate, Ferrous Sulfate, Magnesium Phosphate, Manganese Sulfate, Potassium Chloride, Potassium Iodide, Sodium Citrate, Sodium Hydroxide, Sodium Selenite And Zinc Sulfate), **Vitamins** (Ascorbic Acid, Biotin, Calcium D-Pantothenate, Choline Chloride, DL- $\alpha$ -Tocopheryl Acetate, Folic Acid, Inositol, Niacinamide, Pyridoxine Hydrochloride, Riboflavin, Thiamine Hydrochloride, Vitamin A Palmitate, Vitamin B12, Vitamin D3 and Vitamin K1).

#### **PRODUCT FORMS**

Product	Format	Unit Size	Approximate Product Yield at Normal Dilution	Shelf Life	Item Number	Availability
Enfamil A <sup>+®</sup> Serenity	Powder	578 g	3986 mL	18 Months	Not Available For Hospital Orders‡	Retail only

<sup>‡</sup> For more info, feel free to connect with your Hospital Account Manager.

#### PREPARATION OF FEEDINGS

Your baby's health depends on carefully following the instructions below.

Proper hygiene, preparation, dilution, use and storage are important when preparing infant formula. Powdered infant formulas are not sterile and should <u>not</u> be fed to premature infants or infants who might have immune problems unless directed and supervised by your baby's doctor. Ask your baby's doctor which formula is appropriate for your baby.

<sup>\*</sup> A source of ARA.

<sup>†</sup> A source of DHA.



1. Wash hands thoroughly with soap and water before preparing formula. Boil clean bottles, nipples, caps and utensils in water (2 minutes at a rolling boil).



Boil fresh water at a rolling boil for 2 minutes.Cool to room temperature prior to mixing.



3. Pour desired amount of the cooled water into the bottle. Add powder. Cap bottle and **SHAKE WELL.** 

Use the chart below for the correct amounts of cooled, boiled water and powder. Use scoop in tub to measure powder. Store **DRY** scoop in this tub.

To Make	Boiled Water	Powder
60 mL bottle	<b>60 mL</b> (1/4 cup)	1 packed level scoop
120 mL bottle	<b>120 mL</b> (1/2 cup)	2 packed level scoops
180 mL bottle	<b>180 mL</b> (3/4 cup)	3 packed level scoops
240 mL bottle	<b>240 mL</b> (1 cup)	4 packed level scoops

**WARNING:** Do not use a microwave oven to warm formula. Serious burns may result.

Failure to follow these instructions could result in severe harm. Once prepared, infant formula can spoil quickly. Either feed immediately or cover and store in refrigerator at 2-4°C (35-40°F) for no longer than 24 hours. Do not use prepared formula if it is unrefrigerated for more than a total of 2 hours. Do not freeze prepared formula. After feeding begins, use formula within one hour or discard.

**Storage/Handling:** Store powder at room temperature; avoid extreme temperatures. After opening and between uses, keep pouch and tub lid tightly closed, store in dry area and use contents within 1 month. Do not store loose powder in tub. Use with Enfamil A+® Serenity formula only. Keep powder fresh and prevent bacterial growth by ensuring tub is clean and completely dry. Completely empty tub and wipe with a clean, dry cloth before reusing. Top rack dishwasher safe.

Each pouch has a new batch code and expiration date sticker that must be retained until the contents of the pouch have been consumed.

Do not freeze powder and avoid excessive heat.



# Nutramigen® A+® with LGG®

Hypoallergenic, extensively hydrolyzed, iron fortified infant formula, providing special feeding for babies with cow's milk protein allergy including colicky\* babies.

\* Due to cow's milk protein allergy. Not Kosher

#### INDICATION

Hypoallergenic, extensively hydrolyzed, iron fortified infant formula, providing special feeding for babies with cow's milk protein allergy including colicky\* babies.

#### **NUTRIENTS**

Analysis	per 100 mL (normal dilution)	per 100 g powder		
Energy (kcal/kJ)	68/280	500/2090		
Protein (g)	1.9	14		
% of total energy	11	11		
Source: Hydrolyzed casein, amino aci	d premix			
Carbohydrate (g)	7	52		
% of total energy	41	41		
Source: Corn syrup solids, modified c	orn starch			
Fat (g)	3.6	27		
% of total energy	48	48		
Source: Palm olein oil (44%), soy oil (19.5%), coconut oil (19.5%), high olein sunflower oil (14.5%), ARA & DHA single cell oil blend (2.5%)				
Linoleic Acid (g)	0.5	3.9		
Linolenic Acid (g)	0.05	0.4		
ARA (mg)	17	125		

LGG® is a registered trademark of Chr. Hansen A/S.

Lactobacillus rhamnosus GG (cfu)



DHA (mg)



85

1 x 108

12

1.35 x 107

<sup>\*</sup> Due to cow's milk protein allergy.

Analysis	per 100 mL (normal dilution)	per 100 g powder
Minerals		
Calcium (mg)	64	470
Phosphorus (mg)	35	260
Magnesium (mg)	5.4	40
Iron (mg)	1.22	9
Zinc (mg)	0.68	5
Manganese (mg)	0.017	0.125
Copper (mg)	0.051	0.38
lodine (mg)	0.0102	0.075
Selenium (mg)	0.0019	0.014
Sodium (mg)	32	240
Potassium (mg)	75	550
Chloride (mg)	58	430
Vitamins		
Vitamin A (IU)	200	1500
Vitamin D (IU)	41	300
Vitamin E (IU)	1.36	10
Vitamin K (mg)	0.0061	0.045
Vitamin C (mg)	8.2	60
Thiamine (mg)	0.054	0.4
Riboflavin (mg)	0.061	0.45
Niacin (mg)	0.68	5
Pantothenic Acid (mg)	0.34	2.5
Vitamin B <sub>6</sub> (mg)	0.041	0.3
Folic Acid (mg)	0.0109	0.08
Vitamin B <sub>12</sub> (mg)	0.0002	0.0015
Biotin (mg)	0.002	0.015
Choline (mg)	16.3	120
Inositol (mg)	16.3	120
Carnitine (mg)	1.36	10
Taurine (mg)	4.1	30

Other Characteristics	
Potential Renal Solute Load (m0sm/100 mL) <sup>1</sup>	16.9
Osmolality (m0sm/kg H <sub>2</sub> 0)	300
Osmolarity (m0sm/L)	270
Water (g/100 mL)	89

Nutrient values are subject to change. Consult actual product labels for most current information.

Powdered formulas are not sterile. "To minimize the risk of infection, powdered formulas are not recommended for use in premature or immunocompromised infants unless no appropriate nutritional alternative is available, and then only with strict medical supervision and careful preparation, storage and use  $^{"2}$ 

- 1. Fomon SJ, Ziegler EE. Renal solute load and potential renal solute load in infancy. J Pediatr. 1999;134:11-14.
- International Formula Council. Infant Feeding: Safety Issues for Health Care Professionals. Atlanta, GA: International Formula Council; 2004:4.

#### **INGREDIENTS**

Corn Syrup Solids, Extensively Hydrolyzed Casein\*(From Milk), Palm Olein Oil, Coconut Oil, Soy Oil, High Oleic Sunflower Oil, Modified Corn Starch, Mortierella Alpina Oil $^{\dagger}$ , L-Cystine, Schizochytrium Sp. Oil $^{\dagger}$ , L-Cystine, L-Tyrosine, L-Tryptophan, Taurine, Lactobacillus Rhamnosus GG, L-Carnitine, Ascorbyl Palmitate, Mixed Tocopherols, (May Contain Potassium Hydroxide), **Minerals** (Calcium Citrate, Calcium Hydroxide, Calcium Phosphate, Cupric Sulfate, Ferrous Sulfate, Magnesium Oxide, Manganese Sulfate, Potassium Chloride, Potassium Citrate, Sodium Citrate, Sodium Iodide, Sodium Selenite, And Zinc Sulfate), **Vitamins** (Ascorbic Acid, Biotin, Calcium D-Pantothenate, Choline Chloride, DL- $\alpha$ -Tocopheryl Acetate, Folic Acid, Inositol, Niacinamide, Pyridoxine Hydrochloride, Riboflavin, Thiamine Hydrochloride, Vitamin A Palmitate, Vitamin B<sub>12</sub>, Vitamin D<sub>3</sub> and Vitamin K<sub>1</sub>).

- \* Modified to be better tolerated in milk allergic babies.
- † A source of ARA.
- ‡ A source of DHA.

#### **PRODUCT FORMS**

Product	Format	Unit Size	Approximate Product Yield at Normal Dilution	Shelf Life	Item Number	Availability
Nutramigen® A+® with LGG®	Powder	561 g	4100 mL (17.5 bottles x 237 mL / 8 fl oz)	18 Months	3230626	Hospital & Retail

#### PREPARATION OF FEEDINGS

#### **POWDER**

Your baby's health depends on carefully following the instructions below. Proper hygiene, preparation, dilution, use, and storage are important when preparing infant formula. Powdered infant formulas are not sterile. Ask your baby's doctor which formula is appropriate for your baby. Important Notice: Nutramigen® A+® with LGG® should only be used under medical supervision. Powdered infant formulas and those with added micro-organisms such as LGG® are not sterile and should <u>not</u> be fed to premature infants or infants who might have immune problems unless directed and supervised by your baby's doctor.



1. Wash hands thoroughly with soap and water before preparing formula. Boil clean bottles, nipples, caps, scoop and utensils in water (2 minutes at a rolling boil).



2. Boil fresh water at a rolling boil for 2 minutes. Cool to room temperature prior to mixing.



3. Pour desired amount of the cooled water into the bottle. Add powder. Cap bottle and **SHAKE WELL**.

Use the chart below for the correct amounts of cooled, boiled water and powder. Use scoop in can to measure powder. Store **DRY** scoop in this can.

To Make	Boiled Water	Powder
<b>60 mL</b> bottle	<b>60 mL</b> (1/4 cup)	1 packed level scoop
120 mL bottle	<b>120 mL</b> (1/2 cup)	2 packed level scoops
180 mL bottle	<b>180 mL</b> (3/4 cup)	3 packed level scoops
240 mL bottle	<b>240 mL</b> (1 cup)	4 packed level scoops

**WARNING:** Do not warm the prepared formula beyond preparation temperature of  $40^{\circ}$ C ( $104^{\circ}$ F); this may limit the benefits of LGG culture. Microwaving formula may cause serious burns to baby.

Failure to follow these instructions could result in severe harm. Once prepared, infant formula can spoil quickly. Either feed immediately or cover and store in refrigerator at 2-4°C (35-40°F) for no longer than 24 hours. Do not use prepared formula if it is unrefrigerated for more than a total of 2 hours. Do not freeze prepared formula. After feeding begins, use formula within one hour or discard.

**Powder Storage:** Store cans at room temperature. After opening can, keep tightly covered, store in dry area and use contents within 1 month. Do not freeze powder and avoid excessive heat.



## Nutramigen® A+®

Hypoallergenic, extensively hydrolyzed, iron fortified infant formula for the dietary management of infants with cow's milk protein allergy, including colicky\* infants.

\* Due to cow's milk protein allergy.

Not Kosher

#### INDICATION

Hypoallergenic, extensively hydrolyzed, iron fortified infant formula for the dietary management of infants with cow's milk protein allergy, including colicky\* infants. Contains DHA (a type of Omega-3 fat).

#### **NUTRIENTS**

Analysis	g/100 mL (normal dilution) Nursette® bottle	per 100 mL (normal dilution) powder	per 100 g powder
Energy (kcal/kJ)	68/280	68/280	500/2080
Protein (g)	1.9	1.89	13.9
% of total energy	11	11	11

Source: Casein hydrolyzed, amino acid premix

Carbohydrate (g)	7	7	51
% of total energy	41	41	41

Source (Powder): Corn syrup solids, modified corn starch.

Source (Nursette® Bottle): Corn syrup solids, modified corn starch.

Fat (g)	3.6	3.6	26
% of total energy	48	48	48

Powder Source: Palm olein oil (44%), soy oil (19.5%), coconut oil (19.5%), high oleic sunflower oil (14.5%), ARA & DHA single cell oil blend (2.5%)

Nursette® Source: Palm olein oil (44%), soy oil (19.5%), coconut oil (19.5%), high oleic sunflower oil (14.5%), ARA & DHA single cell oil blend (2.5%)

Linoleic Acid (g)	0.5	0.58	4.3
Linolenic Acid (g)	0.05	0.054	0.4
ARA (mg)	17	23	169
DHA (mg)	11.6	11.5	85



<sup>\*</sup> Due to cow's milk protein allergy.

Analysis	g/100 mL (normal dilution) Nursette® bottle	per 100 mL (normal dilution) powder	per 100 g powder
Minerals			
Ash (g)	0.49		
Calcium (mg) (mMol - Nursette® Bottle)	64 1.59	64 1.59	470 11.7
Phosphorus (mg) (mMol - Nursette® Bottle)	35 1.13	35 1.13	260 8.3
Magnesium (mg) (mMol - Nursette® Bottle)	5.4 0.22	5.4 0.22	40 1.64
Iron (mg)	1.22	1.22	9
Zinc (mg)	0.68	0.68	5
Manganese (mg)	0.017	0.0169	0.124
Copper (mg)	0.051	0.051	0.37
lodine (mg)	0.0102	0.0101	0.075
Selenium (mg)	0.0019	0.00189	0.0139
Sodium (mg) (mMol - Nursette® Bottle)	32 1.38	32 1.38	230 10.2
Potassium (mg) (mMol - Nursette® Bottle)	75 1.9	74 1.9	550 14
Chloride (mg) (mMol - Nursette® Bottle)	58 1.64	58 1.64	430 12
Vitamins			
Vitamin A (IU)	200	200	1490
Vitamin D (IU)	31	34	250
Vitamin E (IU)	1.36	1.35	9.9
Vitamin K (mg)	0.0061	0.0061	0.045
Vitamin C (mg)	8.2	8.1	60
Thiamine (mg)	0.054	0.054	0.4
Riboflavin (mg)	0.061	0.061	0.45

Analysis	g/100 mL (normal dilution) Nursette® bottle	per 100 mL (normal dilution) powder	per 100 g powder
Niacin (mg)	0.68	0.68	5
Pantothenic Acid (mg)	0.34	0.34	2.5
Vitamin B <sub>6</sub> (mg)	0.041	0.041	0.3
Folic Acid (mg)	0.0109	0.0108	0.08
Vitamin B <sub>12</sub> (mg)	0.0002	0.0002	0.00149
Biotin (mg)	0.002	0.002	0.0149
Choline (mg)	16.3	16.2	119
Inositol (mg)	16.3	11.5	85
Carnitine (mg)	1.36	1.35	9.9
Taurine (mg)	4.1	4.1	30
Other Characteristics			
Renal Solute Load (m0	lsm/100 mL)¹		16.9
Osmolality (mOsm/kg I		300 (Powder) 320 (Nursette® Bottle)	
Osmolarity (mOsm/L)		Powder) sette® Bottle)	
Water (g/100 mL)		Powder)	

Powdered formulas are not sterile. "To minimize the risk of infection, powdered formulas are not recommended for use in premature or immunocompromised infants unless no appropriate nutritional alternative is available, and then only with strict medical supervision and careful preparation, storage and use."<sup>2</sup>

89 (Nursette® Bottle)

- 1. Fomon SJ, Ziegler EE. Renal solute load and potential renal solute load in infancy. J Pediatr. 1999;134:11-14.
- 2. International Formula Council. *Infant Feeding: Safety Issues for Health Care Professionals.* Atlanta, GA: International Formula Council; 2004:4.

Nutrient values are subject to change. Consult actual product labels for most current information.

#### INGREDIENTS

#### **POWDER**

Corn Syrup Solids, Hydrolyzed Casein\* (From Milk), Palm Olein Oil, Modified Corn Starch, Soy Oil, Coconut Oil, High Oleic Sunflower Oil, Mortierella Alpina Oil†, L-Cystine, Crypthecodinium Cohnii Oil‡, L-Tyrosine, L-Tryptophan, Taurine, L-Carnitine, Ascorbyl Palmitate, Mixed Tocopherols, (May Contain Potassium Hydroxide), **Minerals** (Calcium Citrate, Calcium Hydroxide, Calcium Phosphate, Cupric Sulfate, Ferrous Sulfate, Magnesium Oxide, Manganese Sulfate, Potassium Chloride, Potassium Citrate, Sodium Citrate, Sodium Selenite And Zinc Sulfate), **Vitamins** (Ascorbic Acid, Inositol, Niacinamide, Pyridoxine Hydrochloride, Riboflavin, Thiamine Hydrochloride, Vitamin A Palmitate, Vitamin B<sub>12</sub>, Vitamin D<sub>3</sub> and Vitamin K<sub>1</sub>).

<sup>\*</sup> Modified to be better tolerated in milk allergic babies.

<sup>†</sup> A source of ARA.

<sup>±</sup> A source of DHA.

#### **NURSETTE® BOTTLE**

Ingredients: Water, Corn Syrup Solids, Extensively Hydrolyzed Casein\* (From Milk), Modified Corn Starch, Palm Olein Oil, Soy Oil, Coconut Oil, High Oleic Sunflower Oil, Carrageenan, Citric Acid, Mortierella Alpina Oil $^\dagger$ , Schizochytrium Sp. Oil $^\dagger$ , L-Cystine, L-Tyrosine, L-Tryptophan, Taurine, L-Carnitine, Mixed Tocopherols, Ascorbyl Palmitate, (May Contain Potassium Hydroxide), Minerals (Calcium Carbonate, Calcium Phosphate, Cupric Sulfate, Ferrous Sulfate, Magnesium Oxide, Manganese Sulfate, Potassium Chloride, Potassium Citrate, Potassium Iodide, Sodium Citrate, Sodium Selenite And Zinc Sulfate), Vitamins (Ascorbic Acid, Biotin, Calcium D-Pantothenate, Choline Chloride, DL- $\alpha$ -Tocophery Acetate, Folic Acid, Inositol, Niacinamide, Pyridoxine Hydrochloride, Riboflavin, Thiamine Hydrochloride, Vitamin A Palmitate, Vitamin B12, Vitamin D3 and Vitamin K1).

- \* Modified to be better tolerated in milk allergic babies.
- † A source of ARA.
- ‡ A source of DHA.

# **PRODUCT FORMS**

Product	Format	Unit Size	Approximate Product Yield at Normal Dilution	Shelf Life	Item Number	Availability
Nutramigen® Δ+®	Powder	385 mL	3300 mL (14 bottles x 8 fl oz)	18 Months	1239201	Hospital & Retail
A	Nursette® Bottle	59 mL	59 mL (2 fl oz)	6 Months	3229972	Hospital only

#### PREPARATION OF FEEDINGS

#### **POWDER**

Your baby's health depends on carefully following the instructions. Proper hygiene, preparation, dilution, use and storage are important when preparing infant formula. Powdered infant formulas are not sterile and should <u>not</u> be fed to premature infants or infants who might have immune problems unless directed and supervised by your baby's doctor. Ask your baby's doctor which formula is appropriate for your baby.



1. Wash hands thoroughly with soap and water before preparing formula. Boil clean bottles, nipples, caps and utensils in water (2 minutes at a rolling boil).



2. Boil fresh water at a rolling boil for 2 minutes. Cool to room temperature prior to mixing.



3. Pour desired amount of the cooled water into the bottle. Add powder. Cap bottle and **SHAKE WELL.** 

Use the chart below for the correct amounts of cooled, boiled water and powder. Use scoop in can to measure powder. Store **DRY** scoop in this can.

To Make	Boiled Water	Powder
60 mL bottle	<b>60 mL</b> (1/4 cup)	1 packed level scoop
120 mL bottle	<b>120 mL</b> (1/2 cup)	2 packed level scoops
180 mL bottle	<b>180 mL</b> (3/4 cup)	3 packed level scoops
240 mL bottle	<b>240 mL</b> (1 cup)	4 packed level scoops

**WARNING:** Do not use a microwave oven to warm formula. Serious burns may result.

Failure to follow these instructions could result in severe harm. Once prepared, infant formula can spoil quickly. Either feed immediately or cover and store in refrigerator at 2-4°C (35-40°F) for no longer than 24 hours. Do not use prepared formula if it is unrefrigerated for more than a total of 2 hours. Do not freeze prepared formula. After feeding begins, use formula within one hour or discard.

**Powder Storage:** Store cans at room temperature. After opening and between uses, keep tightly covered, store in dry area and use contents within 1 month. Do not freeze powder and avoid excessive heat.

#### **NURSETTE® BOTTLES**

Your baby's health depends on carefully following the instructions. Proper hygiene, preparation, use and storage are important when preparing infant formula. Use as directed by your baby's doctor.

Inspect each bottle for signs of damage.



1. Wash hands thoroughly with soap and water before preparing bottle for feeding.



2. **SHAKE BOTTLE WELL** and remove cap.



3. Attach **DISPOSABLE NIPPLE UNIT** (not included) or ask your baby's doctor about the need to boil a clean nipple in water before use.

Failure to follow these instructions could result in severe harm. Opened bottles can spoil quickly. Either feed immediately or replace cap and store in refrigerator at 2-4°C (35-40°F) for no longer than 24 hours. Do not use opened bottle if it is unrefrigerated for more than a total of 2 hours. Do not freeze formula. After feeding begins, use formula within one hour or discard.

**WARNING:** Do not use a microwave oven to warm formula. Serious burns may result.

**Storage:** Store unopened bottles at room temperature. Avoid excessive heat and prolonged exposure to light. Do not freeze.

DO NOT USE IF CAP RING IS BROKEN OR MISSING

USE BY EXPIRATION DATE ON CAN, CARTON AND/OR BOTTLE



# **PURAMINO A+®**

Hypoallergenic, 100% free amino acid-based, iron fortified infant formula for the dietary management of infants and toddlers with severe cow's milk protein and multiple food allergies or other conditions requiring an amino acid-based formulation.

OD (M) IFANCA

# **INDICATION**

Hypoallergenic, 100% free amino acid-based, iron fortified infant formula for the dietary management of infants and toddlers with severe cow's milk protein and multiple food allergies or other conditions requring an amino acid-based formulation. Contains DHA (a type of Omega-3 fat) and MCT Oil.

## **NUTRIENTS**

Analysis	per 100 mL (normal dilution)	per 100 g
Energy (kcal/kJ)	68/280	500/2090
Protein equivalent (g)	1.89	13.9
% of total energy	11	11
Source: Amino acid premix		
Carbohydrate (g)	7.2	53
% of total energy	42	42
Source: Corn syrup solids, modified tap	ioca starch	
Fat (g)	3.6	26
% of total energy	47	47
Source: High oleic sunflower oil (34.5%) DHA single cell oil blend (2.5%)	, MCT oil (33%), soy oi	il (30%), ARA &
Linoleic Acid (g)	0.58	4.3
Linolenic Acid (g)	0.054	0.4
ARA (mg)	23	169
DHA (mg)	11.5	85
Minerals		
Calcium (mg)	64	470
Phosphorus (mg)	35	260
Magnesium (mg)	7.4	55





Analysis	per 100 mL (normal dilution)	per 100 g
Iron (mg)	1.22	9
Zinc (mg)	0.68	5
Manganese (mg)	0.0169	0.124
Copper (mg)	0.051	0.37
lodine (mg)	0.0101	0.075
Selenium (mg)	0.00189	0.0139
Sodium (mg)	32	230
Potassium (mg)	74	550
Chloride (mg)	58	430
Vitamins		
Vitamin A (IU)	200	1490
Vitamin D (IU)	34	250
Vitamin E (IU)	1.35	9.9
Vitamin K (mg)	0.0061	0.045
Vitamin C (mg)	8.1	60
Thiamine (mg)	0.054	0.4
Riboflavin (mg)	0.061	0.45
Niacin (mg)	0.68	5
Pantothenic Acid (mg)	0.34	2.5
Vitamin B <sub>6</sub> (mg)	0.041	0.3
Folic Acid (mg)	0.0108	0.08
Vitamin B <sub>12</sub> (mg)	0.0002	0.00149
Biotin (mg)	0.002	0.0149
Choline (mg)	16.2	119
Inositol (mg)	11.5	85
Carnitine (mg)	1.35	9.9
Taurine (mg)	4.1	30
Other Characteristics		
Renal Solute Load (m0sm/100 mL) <sup>1</sup>		16.9
Osmolality (m0sm/kg H <sub>2</sub> 0)		350
Osmolarity (mOsm/L)		320

Powdered formulas are not sterile. "To minimize the risk of infection, powdered formulas are not recommended for use in premature or immunocompromised infants unless no appropriate nutritional alternative is available, and then only with strict medical supervision and careful preparation, storage and use."  $^2$ 

Water (g/100 mL)

90

- 1. Fomon SJ, Ziegler EE. Renal solute load and potential renal solute load in infancy. J Pediatr. 1999;134:11-14.
- International Formula Council. Infant Feeding: Safety Issues for Health Care Professionals. Atlanta, GA: International Formula Council; 2004:4.

Nutrient values are subject to change. Consult actual product labels for most current information.

## **INGREDIENTS**

Corn Syrup Solids, Amino Acids (L-Aspartic Acid, L-Leucine, L-Lysine Hydrochloride, L-Proline, L-Alanine, L-Valine, Monosodium Glutamate, L-Isoleucine, L-Serine, L-Threonine, L-Tyrosine, L-Arginine, L-Phenylalanine, Glycine, L-Cystine, L-Histidine, L-Tryptophan, L-Methionine), High Oleic Sunflower Oil, Medium Chain Triglycerides (Fractionated Coconut Oil), Soy Oil, Modified Tapioca Starch, Mortierella Alpina Oil\*, Crypthecodinium Cohnii Oil†, Taurine, L-Carnitine, Ascorbyl Palmitate, Mixed Tocopherols, **Minerals** (Calcium Citrate, Calcium Hydroxide, Calcium Phosphate, Cupric Sulfate, Ferrous Sulfate, Magnesium Phosphate, Manganese Sulfate, Potassium Chloride, Potassium Citrate, Sodium Citrate, Sodium Iodide, Sodium Selenite And Zinc Sulfate), **Vitamins** (Ascorbic Acid, Biotin, Calcium D-Pantothenate, Choline Chloride, DL-α-Tocopheryl Acetate, Folic Acid, Inositol, Niacinamide, Pyridoxine Hydrochloride, Riboflavin, Thiamine Hydrochloride, Vitamin A Palmitate, Vitamin B<sub>12</sub>, Vitamin D<sub>3</sub> and Vitamin K<sub>1</sub>).

- \* A source of ARA.
- † A source of DHA.

## PRODUCT FORMS

Product	Format	Unit Size	Approximate Product Yield at Normal Dilution	Shelf Life	Item Number	Availability
PURIMINO A+®	Powder	400 g	2900 mL (12 bottles x 8 fl oz)	15 Months	1233351	Hospital & Retail

#### PREPARATION OF FEEDINGS

#### **POWDER**

#### Your baby's health depends on carefully following the instructions below.

Proper hygiene, preparation, dilution, use and storage are important when preparing infant formula. Powdered infant formulas are not sterile and should <u>not</u> be fed to premature infants or infants who might have immune problems unless directed and supervised by your baby's doctor. Ask your baby's doctor which formula is appropriate for your baby.



1. Wash hands thoroughly with soap and water before preparing formula. Boil clean bottles, nipples, caps and utensils in water (2 minutes at a rolling boil).



 Boil fresh water at a rolling boil for 2 minutes.
 Cool to room temperature prior to mixing.



3. Pour desired amount of the cooled water into the bottle. Add powder. Cap bottle and **SHAKE WELL** 

Use the chart below for the correct amounts of cooled, boiled water and powder. Use scoop in can to measure powder. Store **DRY** scoop in this can.

To Make	Boiled Water	Powder
65 mL bottle	<b>60 mL</b> (1/4 cup)	2 unpacked level scoops
130 mL bottle	<b>120 mL</b> (1/2 cup)	4 unpacked level scoops
195 mL bottle	<b>180 mL</b> (3/4 cup)	6 unpacked level scoops
260 mL bottle	<b>240 mL</b> (1 cup)	8 unpacked level scoops

**WARNING:** Do not use a microwave oven to warm formula. Serious burns may result.

Failure to follow these instructions could result in severe harm. Once prepared, infant formula can spoil quickly. Either feed immediately or cover and store in refrigerator at 2-4°C (35-40°F) for no longer than 24 hours. Do not use prepared formula if it is unrefrigerated for more than a total of 2 hours. Do not freeze prepared formula. After feeding begins, use within one hour or discard.

**Storage/Handling:** Store cans at room temperature. After opening and between uses, keep tightly covered, store in dry area and use contents within 1 month. Do not freeze powder and avoid excessive heat.

#### USE BY EXPIRATION DATE ON BOTTOM OF CAN



# **PURAMINO A<sup>+®</sup> JUNIOR**

Hypoallergenic, 100% free amino acid-based powder for the dietary management of children over 12 months of age with severe cow's milk protein and multiple food allergies or other conditions requiring an amino acid-based formulation.

(U) D (M) IFANCA

# INDICATION

Hypoallergenic, 100% free amino acid-based powder for the dietary management of children over 12 months of age with severe cow's milk protein and multiple food allergies or other conditions requiring an amino acid-based formulation. Contains DHA (a type of Omega-3 fat) and MCT Oil.

# **NUTRIENTS**

Analysis	per 100 mL (normal dilution)	per 100 g
Energy (kcal/kJ)	100/420	490/2050
Protein equivalent (g)	2.8	14
% of total energy	11	11
Source: Amino acid premix		
Carbohydrate (g)	12	59
% of total energy	49	49
Source: Corn syrup solids, modified tap	ioca starch	
Fat (g)	4.5	22
% of total energy	40	40
Source: High oleic sunflower oil (46%), single cell oil (1.3%)	MCT oil (33%), soy oil	(20%), DHA
Linoleic Acid (g)	0.6	2.9
Linolenic Acid (g)	0.06	0.3
ARA (mg)	0	0
DHA (mg)	23	113
Minerals		
Calcium (mg)	100	490
Phosphorus (mg)	60	290





Analysis	per 100 mL (normal dilution)	per 100 g
Magnesium (mg)	16	78
Iron (mg)	1	4.9
Zinc (mg)	0.9	4.4
Manganese (mg)	0.057	0.28
Copper (mg)	0.11	0.54
lodine (mg)	0.015	0.074
Selenium (mg)	0.0032	0.0157
Chromium (mg)	0.003	0.0147
Molybdenum (mg)	0.0045	0.022
Sodium (mg)	40	196
Potassium (mg)	100	490
Chloride (mg)	61	300
Vitamins		
Vitamin A (IU)	200	990
Vitamin D (IU)	56	270
Vitamin E (IU)	1.79	8.8
Vitamin K (mg)	0.004	0.02
Vitamin C (mg)	10	49
Thiamine (mg)	0.075	0.37
Riboflavin (mg)	0.1	0.49
Niacin (mg)	1	4.9
Pantothenic Acid (mg)	0.43	2.1
Vitamin B <sub>6</sub> (mg)	0.1	0.49
Folic Acid (mg)	0.015	0.074
Vitamin B <sub>12</sub> (mg)	0.00025	0.00123
Biotin (mg)	0.0022	0.0108
Choline (mg)	22	108
Inositol (mg)	17	83
Carnitine (mg)	2	9.8
Taurine (mg)	6	29

Other Characteristics	
Renal Solute Load (m0sm/100 mL) <sup>1</sup>	24
Osmolality (mOsm/kg H <sub>2</sub> O)	570
Osmolarity (mOsm/L)	480
Water (g/100 mL)	85

Nutrient values are subject to change. Consult actual product labels for most current information.

# **INGREDIENTS**

Corn Syrup Solids, High Oleic Sunflower Oil, Amino Acids (Potassium Aspartate, L-Leucine, L-Lysine Hydrochloride, L-Proline, L-Alanine, L-Valine, L-Isoleucine, Monosodium Glutamate, L-Threonine, L-Serine, L-Tyrosine, L-Aspartic Acid, L-Arginine, L-Phenylalanine, L-Histidine, L-Cystine, Glycine, L-Tryptophan, L-Methionine), Medium Chain Triglycerides (Fractionated Coconut Oil), Soy Oil, Modified Tapioca Starch, Schizochytrium Sp. Oil\*, Taurine, L-Carnitine, Ascorbyl Palmitate, Mixed Tocopherols, **Minerals** (Calcium Citrate, Calcium Phosphate, Chromic Chloride, Cupric Sulfate, Ferrous Sulfate, Magnesium Phosphate, Manganese Sulfate, Sodium Citrate, Sodium Iodide, Sodium Molybdate, Sodium Selenite And Zinc Sulfate), **Vitamins** (Ascorbic Acid, Biotin, Calcium D-Pantothenate, Choline Chloride, DL- $\alpha$ -Tocopheryl Acetate, Folic Acid, Inositol, Niacinamide, Pyridoxine Hydrochloride, Riboflavin, Thiamine Hydrochloride, Vitamin A Palmitate, Vitamin B12, Vitamin D3 and Vitamin K1).

# **PRODUCT FORMS**

Product	Format	Unit Size	Approximate Product Yield at Normal Dilution	Shelf Life	Item Number	Availability
PURIMINO A+® JUNIOR	Powder	400 g	2900 mL (8 bottles x 8 fl oz)	15 Months	2047834	Hospital & Retail

## PREPARATION OF FEEDINGS

USE ONLY AS DIRECTED BY YOUR MEDICAL PROFESSIONAL. PURAMINO  $A^{+ \circ}$  JUNIOR IS NOT DESIGNED FOR USE AS AN INFANT FORMULA.

<sup>1.</sup> Fomon SJ, Ziegler EE. Renal solute load and potential renal solute load in infancy. J Pediatr. 1999;134:11-14.

<sup>\*</sup> A source of DHA.

# INSTRUCTIONS FOR PREPARATION AND USE FOR ORAL FEEDING:

Your medical professional will provide the correct amount of powder to mix with water for consumption.\* It is important to follow the directions below. Measure the correct amount of water into a suitable container for mixing. Next, add the required amount of PURAMINO A<sup>+®</sup> JUNIOR powder. Mix well until blended.

Failure to follow these instructions could result in severe harm. Once prepared, this product can spoil quickly. Either consume the prepared beverage immediately or cover and store in refrigerator at 2-4°C (35-40°F). Use within 24 hours of preparation. Shake well before each use. Do not use prepared product if it is unrefrigerated for more than a total of 2 hours. Do not freeze prepared product. After feeding begins, consume the prepared beverage within one hour or discard.

\* If instructed to use the scoop in the can to make 30 kcal/30 mL: For every 30 mL (1 US fl. oz.) of water, 1 unpacked level scoop of powder (6.8 g) will make approximately 35 mL (1.2 US fl. oz.) of prepared product.

To make approximately 177 mL (approximately 6 US fl. oz.) of prepared product, add 5 unpacked level scoops (34 g) of powder to 152 mL (5 US fl. oz.) of water. To make approximately 246 mL (approximately 8 US fl. oz.) of prepared product, add 7 unpacked level scoops (47.6 g) of powder to 211 mL (7 US fl. oz.) of water.

This 400 g can of PURAMINO  $A^{+\otimes}$  JUNIOR will make approximately 2 L (2.1 US quarts) of beverage at 30 kcal/30 mL.

**WARNING:** Do not boil product. Do not use a microwave oven to prepare or warm product. Serious burns may result.

**Powder Storage:** Store cans at room temperature. After opening can, keep tightly covered, store in dry area and use contents within 1 month. Store **DRY** scoop in this can. Do not freeze powder and avoid excessive heat.

**WARNING:** Not for parenteral (I.V.) use.

**USE BY EXPIRATION DATE ON BOTTOM OF CAN** 



# Pregestimil® A+®

Hypoallergenic, extensively hydrolyzed, iron fortified infant formula for the dietary management of infants with fat malabsorption and who may also be sensitive to intact proteins.

**Not Kosher** 

# INDICATION

Hypoallergenic, extensively hydrolyzed, iron fortified infant formula for the dietary management of infants with fat malabsorption and who may also be sensitive to intact proteins. Contains DHA (a type of Omega-3 fat).

# **NUTRIENTS**

Analysis	per 100 mL (normal dilution)	per 100 g
Energy (kcal/kJ)	68/280	500/2090
Protein (g)	1.89	14
% of total energy	11	11
Source: Casein hydrolysate, amino ac	id premix	
Carbohydrate (g)	6.9	51
% of total energy	40 (Powder)	40

% of total energy	41 (Nursette® Bottle)†	40
Source (Powder): Corn syrup solids, m Source (Nursette® Bottle): Corn syrup		
Fat (g)	3.8	28
% of total energy	49 (Powder)	49

Source (Powder): MCT (55%), soy (35%), corn (10%), high oleic safflower (7.5%) oils, ARA & DHA single cell oil blend (2.5%)

Source (Nursette® Bottle): MCT (55%), soy (35%), high oleic safflower (7.5%) oils, ARA & DHA single cell oil blend (2.5%)

Linoleic Acid (g)	0.64	4.7
-------------------	------	-----

<sup>†</sup> Per 100 mL of undiluted liquid.





Bottle)†

Analysis	per 100 mL (normal dilution)	per 100 g
Linolenic Acid (g)	0.064 (Powder) 0.081 (Nursette® Bottle)*	0.48
ARA (mg)	23	170
DHA (mg)	11.5	85
Minerals		
Calcium (mg) (mMol - Nursette® Bottle)	64 1.59	470 -
Phosphorus (mg) (mMol - Nursette® Bottle)	35 1.13	260 -
Magnesium (mg) (mMol - Nursette® Bottle)	5.4 0.22	40 -
Iron (mg)	1.22	9
Zinc (mg)	0.68	5
Manganese (mg)	0.0169	0.125
Copper (mg)	0.051	0.38
lodine (mg)	0.0101	0.075
Selenium (mg)	0.00189	0.014
Sodium (mg) (mMol - Nursette® Bottle)	32 1.38	240
Potassium (mg) (mMol - Nursette® Bottle)	74 1.9	550 -
Chloride (mg) (mMol - Nursette® Bottle)	58 1.64	430 -
Vitamins		
Vitamin A (IU)	240	1750
Vitamin D (IU)	34	250
Vitamin E (IU)	2.7	20
Vitamin K (mg)	0.0081	0.06
Vitamin C (mg)	8.1	60
Thiamine (mg)	0.054	0.4
Riboflavin (mg)	0.061	0.45

<sup>\*</sup> Per 100 mL of undiluted liquid.

Analysis	per 100 mL (normal dilution)	per 100 g
Niacin (mg)	0.68	5
Pantothenic Acid (mg)	0.34	2.5
Vitamin B <sub>6</sub> (mg)	0.041	0.3
Folic Acid (mg)	0.0108	0.08
Vitamin B <sub>12</sub> (mg)	0.0002	0.0015
Biotin (mg)	0.002	0.015
Choline (mg)	16.2	120
Inositol (mg)	11.5	85
Carnitine (mg)	1.35	10
Taurine (mg)	4.1	30

Other Characteristics			
Renal Solute Load (m0sm/100 mL) <sup>1</sup> 16.9			
Osmolality (mOsm/kg H <sub>2</sub> O)	320 (Powder) 340 (Nursette® Bottle)*		
Osmolarity (mOsm/L)	290 (Powder) 310 (Nursette® Bottle)*		
Water (g/100 mL)	90 (Powder) 90 (Nursette® Bottle)*		

<sup>\*</sup> Per 100 mL of undiluted liquid.

Powdered formulas are not sterile. "To minimize the risk of infection, powdered formulas are not recommended for use in premature or immunocompromised infants unless no appropriate nutritional alternative is available, and then only with strict medical supervision and careful preparation, storage and use."<sup>2</sup>

- 1. Fomon SJ, Ziegler EE. Renal solute load and potential renal solute load in infancy. J Pediatr. 1999;134:11-14.
- International Formula Council. Infant Feeding: Safety Issues for Health Care Professionals. Atlanta, GA: International Formula Council; 2004:4.

Nutrient values are subject to change. Consult actual product labels for most current information.

## **INGREDIENTS**

#### **POWDER**

Corn Syrup Solids, Hydrolyzed Casein† (From Milk), Medium Chain Triglycerides (Fractionated Coconut Oil), Modified Corn Starch, Soy Oil, Corn Oil, High Oleic Vegetable Oil (Safflower And/Or Sunflower Oil), Mortierella Alpina Oil‡, L-Cystine, Crypthecodinium Cohnii Oil‡, L-Tyrosine, L-Tryptophan, Taurine, L-Carnitine, Ascorbyl Palmitate, Mixed Tocopherols, (May Contain Potassium Hydroxide), Minerals (Calcium Citrate, Calcium Phosphate, Cupric Sulfate, Ferrous Sulfate, Magnesium Oxide, Manganese Sulfate, Potassium Chloride, Potassium Citrate,

Sodium Citrate, Sodium Iodide, Sodium Selenite And Zinc Sulfate), **Vitamins** (Ascorbic Acid, Biotin, Calcium D-Pantothenate, Choline Chloride, DL- $\alpha$ -Tocopheryl Acetate, Folic Acid, Inositol, Niacinamide, Pyridoxine Hydrochloride, Riboflavin, Thiamine Hydrochloride, Vitamin A Palmitate, Vitamin B<sub>12</sub>, Vitamin D<sub>3</sub> and Vitamin K<sub>1</sub>).

- † Modified to be better tolerated in milk allergic babies.
- ‡ A source of ARA.
- ¶ A source of DHA.

#### 59 mL NURSETTE® BOTTLE

Water, Corn Syrup Solids, Hydrolyzed Casein† (From Milk), Medium Chain Triglycerides (Fractionated Coconut Oil), Modified Corn Starch, Soy Oil, High Oleic Vegetable Oil (Safflower And/Or Sunflower Oil), Carrageenan, Mortierella Alpina Oil‡, L-Cystine, Crypthecodinium Cohnii Oil¹, L-Tyrosine, L-Tryptophan, Taurine, L-Carnitine, Ascorbyl Palmitate, Mixed Tocopherols, (May Contain Potassium Hydroxide), **Minerals** (Calcium Carbonate, Calcium Hydroxide, Calcium Phosphate, Cupric Sulfate, Ferrous Sulfate, Magnesium Chloride, Manganese Sulfate, Potassium Chloride, Potassium Citrate, Potassium Phosphate, Sodium Citrate, Sodium Iodide, Sodium Selenite And Zinc Sulfate), **Vitamins** (Ascorbic Acid, Biotin, Calcium D-Pantothenate, Choline Chloride, DL- $\alpha$ -Tocopheryl Acetate, Folic Acid, Inositol, Niacinamide, Pyridoxine Hydrochloride, Riboflavin, Thiamine Hydrochloride, Vitamin A Palmitate, Vitamin B<sub>12</sub>, Vitamin D<sub>3</sub> and Vitamin K<sub>1</sub>).

- † Modified to be better tolerated in milk allergic babies.
- ‡ A source of ARA.
- ¶ A source of DHA.

# **PRODUCT FORMS**

Product	Format	Unit Size	Approximate Product Yield at Normal Dilution	Shelf Life	Item Number	Availability
Pregestimil®	Powder	454 g	3400 mL (14 bottles x 8 fl oz)	21 Months	1238161	Hospital & Online
A+®	Nursette® Bottle	59 mL	59 mL (2 fl oz)	12 Months	1273987	Hospital only

# PREPARATION OF FEEDINGS

#### **POWDER**

Your baby's health depends on carefully following the instructions below.

Use only as directed by a medical professional. Improper hygiene, preparation, dilution, use or storage may result in severe harm. Powdered infant formulas are not sterile and should <u>not</u> be fed to premature infants or infants who might have immune problems unless directed and supervised by your baby's doctor. Ask your baby's doctor which formula is appropriate for your baby.



1. Wash hands thoroughly with soap and water before preparing formula. Boil clean bottles, nipples, caps and utensils in water (2 minutes at a rolling boil).



2. Boil fresh water at a rolling boil for 2 minutes. Cool to room temperature prior to mixing.



3. Pour desired amount of the cooled water into the bottle. Add powder. Cap bottle and **SHAKE WELL.** 

Use the chart below for the correct amounts of cooled, boiled water and powder. Use scoop in can to measure powder. Store **DRY** scoop in this can.

To Make	Boiled Water	Powder
60 mL bottle	<b>60 mL</b> (1/4 cup)	1 packed level scoop
120 mL bottle	<b>120 mL</b> (1/2 cup)	2 packed level scoops
195 mL bottle	<b>180 mL</b> (3/4 cup)	3 packed level scoops
260 mL bottle	<b>240 mL</b> (1 cup)	4 packed level scoops

**WARNING:** Do not use a microwave oven to warm formula. Serious burns may result.

Failure to follow these instructions could result in severe harm. Once prepared, infant formula can spoil quickly. Either feed immediately or cover and store in refrigerator at 2-4°C (35-40°F) for no longer than 24 hours. Do not use prepared formula if it is unrefrigerated for more than a total of 2 hours. Do not freeze prepared formula. After feeding begins, use within one hour or discard.

**Storage/Handling:** Store powder at room temperature. After opening and between uses, keep tightly covered, store in dry area and use contents within 1 month. Do not freeze powder and avoid excessive heat.

#### USE BY EXPIRATION DATE ON BOTTOM OF CAN

#### **NURSETTE® BOTTLES**

Your baby's health depends on carefully following the instructions below.

Proper hygiene, preparation, use and storage are important when preparing infant formula. Use as directed by your baby's doctor.

Inspect each bottle for signs of damage.



1. Wash hands thoroughly with soap and water before preparing bottle for feeding.



2. SHAKE BOTTLE WELL and remove cap.



3. Attach **DISPOSABLE NIPPLE UNIT** (not included) or ask your baby's doctor about the need to boil a clean nipple in water before use.

Failure to follow these instructions could result in severe harm. Opened bottles and prepared bottles can spoil quickly. Either feed immediately or replace cap and store in refrigerator at 2-4°C (35-40°F) for no longer than 24 hours. Do not use opened bottle or prepared bottle if it is unrefrigerated for more than a total of 2 hours. Do not freeze. After feeding begins, use formula within one hour or discard.

**WARNING:** Do not use a microwave oven to warm formula. Serious burns may result.

**Storage:** Store unopened bottles at room temperature. Avoid excessive heat and prolonged exposure to light. Do not freeze.

DO NOT USE IF CAP RING IS BROKEN OR MISSING

**USE BY EXPIRATION DATE ON CARTON AND BOTTLE** 



# Enfamil A+® 2

Cow's milk-based, iron fortified transitional infant formula for infants 6 months of age or older.

**⊕**D

# INDICATION

Cow's milk-based, iron fortified transitional infant formula for infants 6 months of age or older. Contains age appropriate levels of calcium, iron & protein. Also contains DHA (a type of Omega-3 fat) and two dietary fibres (galactooligosaccharides and polydextrose).

# **NUTRIENTS**

Analysis	per 100 mL (normal dilution or RTF)	per 100 g
Energy (kcal/kJ)	68/280	500/2090
Protein (g)	1.76 (Powder) 1.8 (RTF & Conc)	12.9
% of total energy	10	10

Source: Nonfat milk (Powder & Conc): skim milk & milk protein isolate (RTF)

Source: Nonfat milk (Powder & Conc); skim milk & milk protein isolate (RTF)			
Carbohydrate (g)	7.3 (Powder) 7.5 (Conc) 7.1 (RTF)	54	
% of total energy	42 (Powder & RTF) 45 (Conc)	42	
Galactooligosaccharides (g)	0.2	1.49	
Polydextrose (g)	0.2	1.49	
Source: Lactose, corn syrup solids, galactooligosaccharides, polydextrose			
Fat (g)	3.6 (Powder & RTF) 3.4 (Conc)	26	
% of total energy	48 (Powder & RTF) 45 (Conc)	48	

Source: Palm olein (44%), soy (19.5%), coconut (19.5%), high oleic sunflower oils (14.5%), ARA and DHA single cell oil blend (2.5%)

Linoleic Acid (g)	0.54 (Powder)	,
Efficienc Acid (g)	0.5 (RTF & Conc)	4





Analysis	per 100 mL (normal dilution or RTF)	per 100 g
Linolenic Acid (g)	0.052 (Powder) 0.05 (Conc) 0.1 (RTF)	0.38
ARA (mg)	23	169
DHA (mg)	11.5 (Powder & Conc) 11 (RTF)	84
Minerals		
Ash (g)	0.68 (Powder) 0.61 (Conc) 0.47 (RTF)	5
Calcium (mg)	132 (Powder) 81 (Conc & RTF)	970
Phosphorus (mg)	88 (Powder) 54 (Conc & RTF)	650
Magnesium (mg)	5.4	40
Iron (mg)	1.01 (Powder & RTF) 1.02 (Conc)	7.4
Zinc (mg)	0.68	5
Manganese (mg)	0.0101 (Powder & RTF) 0.0102 (Conc)	0.074
Copper (mg)	0.051	0.37
lodine (mg)	0.0068 (Powder & RTF) 0.017 (Conc)	0.05
Selenium (mg)	0.00189 (Powder & RTF) 0.002 (Conc)	0.0139
Sodium (mg)	24	179
Potassium (mg)	88	650
Chloride (mg)	54	400
Vitamins		
Vitamin A (IU)	200	1490
Vitamin D (IU)	41	300
Vitamin E (IU)	1.35 (Powder & RTF) 1.36 (Conc)	9.9
Vitamin K (mg)	0.0061	0.045

Analysis	per 100 mL (normal dilution or RTF)	per 100 g
Vitamin C (mg)	8.1	60
Thiamine (mg)	0.054	0.4
Riboflavin (mg)	0.095	0.7
Niacin (mg)	0.68	5
Pantothenic Acid (mg)	0.34	2.5
Vitamin B <sub>6</sub> (mg)	0.041	0.3
Folic Acid (mg)	0.0108 (Powder & RTF) 0.0109 (Conc)	0.079
Vitamin B <sub>12</sub> (mg)	0.0002	0.00149
Biotin (mg)	0.002	0.0149
Choline (mg)	16.2 (Powder & RTF) 16.3 (Conc)	119
Inositol (mg)	4.1 (Powder & RTF) 12.2 (Conc)	30
Carnitine (mg)	1.35 (Powder & RTF) None added (Conc)	9.9
Taurine (mg)	4.1 (Powder) 3.8 (RTF & Conc)	30

Other Characteristics	
Renal Solute Load (m0sm/100 mL) <sup>1</sup>	17.6 (Powder) 16.9 (Conc) 16.6 (RTF)
Osmolality (m0sm/kg H <sub>2</sub> 0)	270 (Powder) 280 (Conc) 250 (RTF)
Osmolarity (mOsm/L)	240 (Powder) 250 (Conc) 230 (RTF)
Water (g/100 mL)	89 (Powder) 90 (Conc & RTF)

Powdered formulas are not sterile. "To minimize the risk of infection, powdered formulas are not recommended for use in premature or immunocompromised infants unless no appropriate nutritional alternative is available, and then only with strict medical supervision and careful preparation, storage and use."<sup>2</sup>

Nutrient values are subject to change. Consult actual product labels for most current information.

<sup>1.</sup> Fomon SJ, Ziegler EE. Renal solute load and potential renal solute load in infancy. J Pediatr. 1999;134:11-14.

<sup>2.</sup> International Formula Council. Infant Feeding: Safety Issues for Health Care Professionals. Atlanta, GA: International Formula Council; 2004:4.

## **INGREDIENTS**

#### **POWDER**

Milk Ingredients, Corn Syrup Solids, Palm Olein Oil, Lactose, Coconut Oil, Soy Oil, High Oleic Sunflower Oil, Polydextrose, Galactooligosaccharide Solids (GOS), Soy Lecithin, Mortierella Alpina Oil\*, Crypthecodinium Cohnii Oil†, Taurine, L-Carnitine, Ascorbyl Palmitate, Mixed Tocopherols, (May Contain Potassium Hydroxide), **Minerals** (Calcium Phosphate, Cupric Sulfate, Ferrous Sulfate, Manganese Sulfate, Potassium Chloride, Sodium Chloride, Sodium Citrate, Sodium Selenite And Zinc Sulfate), **Vitamins** (Ascorbic Acid, Biotin, Calcium D-Pantothenate, Choline Chloride, DL- $\alpha$ -Tocopheryl Acetate, Folic Acid, Inositol, Niacinamide, Pyridoxine Hydrochloride, Riboflavin, Thiamine Hydrochloride, Vitamin A Palmitate, Vitamin B<sub>12</sub>, Vitamin D<sub>3</sub> and Vitamin K<sub>1</sub>).

- \* A source of ARA.
- † A source of DHA.

#### CONCENTRATE

Water, Modified Milk Ingredients, Corn Syrup Solids, Palm Olein Oil, Lactose, Soy Oil, Coconut Oil, High Oleic Sunflower Oil, Polydextrose, Galactooligosaccharide Solids (GOS), Soy Lecithin, Mortierella Alpina Oil\*, Crypthecodinium Cohnii Oil†, Taurine, Carrageenan, Ascorbyl Palmitate, Mixed Tocopherols, (May Contain Potassium Hydroxide), **Minerals** (Calcium Carbonate, Calcium Phosphate, Cupric Sulfate, Ferrous Sulfate, Magnesium Phosphate, Manganese Sulfate, Potassium Iodide, Potassium Citrate, Sodium Selenite And Zinc Sulfate) **Vitamins** (Ascorbic Acid, Biotin, Calcium D-Pantothenate, Choline Chloride, DL- $\alpha$ -Tocopheryl Acetate, Folic Acid, Inositol, Niacinamide, Pyridoxine Hydrochloride, Riboflavin, Sodium Ascorbate, Thiamine Hydrochloride, Vitamin A Palmitate, Vitamin B<sub>12</sub>, Vitamin D<sub>3</sub> and Vitamin K<sub>1</sub>).

- \* A source of ARA.
- † A source of DHA.

#### **READY TO FEED**

Water, Modified Milk Ingredients, Corn Syrup Solids, Lactose, Palm Olein Oil, Soy Oil, Coconut Oil, High Oleic Sunflower Oil, Galactooligosaccharide Solids (GOS), Polydextrose, Mono- And Diglycerides, Soy Lecithin, Mortierella Alpina Oil\*, Carrageenan, Crypthecodinium Cohnii Oil\*, Taurine, L-Carnitine, Ascorbyl Palmitate, Mixed Tocopherols, (May Contain Potassium Hydroxide), **Minerals** (Calcium Hydroxide, Calcium Phosphate, Cupric Sulfate, Ferrous Sulfate, Manganese Sulfate, Potassium Chloride, Potassium Citrate, Sodium Selenite And Zinc Sulfate), **Vitamins** (Ascorbic Acid, Biotin, Calcium D-Pantothenate, Choline Chloride, DL- $\alpha$ -Tocopheryl Acetate, Folic Acid, Inositol, Niacinamide, Pyridoxine Hydrochloride, Riboflavin, Sodium Ascorbate, Thiamine Hydrochloride, Vitamin A Palmitate, Vitamin B<sub>12</sub>, Vitamin D<sub>3</sub> and Vitamin K<sub>1</sub>).

- \* A source of ARA.
- † A source of DHA.

### PRODUCT FORMS

Product	Format	Unit Size	Approximate Product Yield at Normal Dilution	Shelf Life	Item Number	Availability
	Powder	550 g	4000 mL (17 bottles x 8 fl oz)	18 Months	2014690	Hospital & Retail
Enfamil®	Powder	992 g	7300 mL (31 bottles x 8 fl oz)	18 Months	2014689	Hospital & Retail
A+® 2	Concentrate	385 mL	770 mL (3 bottles x 8 fl oz)	12 Months	2015336	Hospital & Retail
	RTF	237 mL	237 mL (1 bottle x 8 fl oz)	12 Months	2050852	Hospital & Retail

# PREPARATION OF FEEDINGS

#### **POWDER**

Your baby's health depends on carefully following the instructions below.

Proper hygiene, preparation, dilution, use and storage are important when preparing infant formula. Powdered infant formulas are not sterile and should **not** be fed to premature infants or infants who might have immune problems unless directed and supervised by your baby's doctor. Ask your baby's doctor which formula is appropriate for your baby.



1. Wash hands thoroughly with soap and water before preparing formula.



2. Pour desired amount of 3. Cap bottle and SHAKE water into the bottle. Add powder.



WELL.

Use the chart below for the correct amounts of water and powder. Use scoop in tub to measure powder. Store **DRY** scoop in this tub.

To Make	Water	Powder
<b>60 mL</b> bottle	<b>60 mL</b> (1/4 cup)	1 unpacked level scoop
120 mL bottle	<b>120 mL</b> (1/2 cup)	2 unpacked level scoops
180 mL bottle	<b>180 mL</b> (3/4 cup)	3 unpacked level scoops
<b>240 mL</b> bottle	<b>240 mL</b> (1 cup)	4 unpacked level scoops

**WARNING:** Do not use a microwave oven to warm formula. Serious burns may result.

Failure to follow these instructions could result in severe harm. Once prepared, infant formula can spoil quickly. Either feed immediately or cover and store in refrigerator at  $2-4^{\circ}\text{C}$  ( $35-40^{\circ}\text{F}$ ) for no longer than 24 hours. Do not use prepared formula if it is unrefrigerated for more than a total of 2 hours. Do not freeze prepared formula. After feeding begins, use within one hour or discard.

**Storage/Handling:** Store powder at room temperature. After opening and between uses, keep tightly covered, store in a dry area, and use contents within 1 month. Do not freeze powder and avoid excessive heat.

#### CONCENTRATE

Your baby's health depends on carefully following the instructions below. Proper hygiene, preparation, dilution, use and storage are important when preparing infant formula. Use as directed by your baby's doctor.







3. Pour desired amount of water into the bottle. Add an **equal** amount of concentrated liquid. Cap bottle and **SHAKE WELL**.

**WARNING:** Do not use a microwave oven to warm formula. Serious burns may result.

Failure to follow these instructions could result in severe harm. Opened cans and prepared formula can spoil quickly. Either feed immediately or cover and store in refrigerator at  $2-4^{\circ}\text{C}$  ( $35-40^{\circ}\text{F}$ ) for no longer than 48 hours. Do not use opened can and/or prepared formula if they are unrefrigerated for more than a

total of 2 hours. Do not freeze formula. After feeding begins, use within one hour or discard.

**Storage:** Store unopened cans at room temperature. Avoid excessive heat. Do not freeze

#### **READY TO FEED**

Your baby's health depends on carefully following the instructions below. Proper hygiene, preparation, dilution, use and storage are important when preparing infant formula. Use as directed by your baby's doctor. **Inspect** each bottle for signs of damage.



1. Wash hands thoroughly with soap and water before preparing bottle for feeding.



2. SHAKE BOTTLE WELL, remove protective seal around cap, remove cap and foil seal.



3. Attach NIPPLE UNIT (not included) or ask your baby's doctor about the need to boil a clean nipple in water before use.

**WARNING:** Do not use a microwave oven to warm formula. Serious burns may result.

Failure to follow these instructions could result in severe harm. Open bottles and prepared bottles can spoil quickly. Either feed immediately or cover and store in refrigerator at 2-4°C (35-40°F) for no longer than 48 hours. Do not use opened bottle or prepared bottle if it is unrefrigerated for more than a total of 2 hours. Do not freeze formula. After feeding begins, use formula within one hour or discard.

**Storage:** Store unopened bottles at room temperature. Avoid excessive heat and prolonged exposure to light. Do not freeze.



# Enfamil® Sterile Water

Sterile water for oral use only.

O Pareve (manufactured on dairy equipment) (M) IFANCA

# **INDICATION**

Water for oral use only.

# INGREDIENT

Water

# **PRODUCT FORMS**

Product	Format	Unit Size	Approximate Product Yield at Normal Dilution	Shelf Life	ltem Number	Availability
Enfamil® Sterile Water	Nursette® Bottle	59 mL	59 mL (2 fl oz)	12 Months	1215250	Hospital only

Nutrients - N/A

# PREPARATION OF FEEDINGS

Your baby's health depends on carefully following the instructions below.

Proper hygiene, preparation, use and storage are important when preparing infant feedings. Use as directed by your baby's doctor.

Inspect each bottle for signs of damage.





1. Wash hands thoroughly with soap and water before preparing bottle for feeding.



2. Remove cap.



3. Attach **DISPOSABLE NIPPLE UNIT** (not included) or ask your baby's doctor about the need to boil a clean nipple in water before use.

Upon opening, either feed immediately or replace cap and store in refrigerator at 2-4°C (35-40°F) for no longer than 24 hours. After feeding begins, do not refrigerate feeding bottle. You must use within 1 hour or discard.

**WARNING:** Do not use a microwave oven to warm water. Serious burns may result.

WARNING: Not for parenteral (I.V.) use.

This water is ready for use as drinking water or for other oral purposes as directed by your baby's doctor.

**STORAGE:** Store unopened bottles at room temperature. Avoid excessive heat. Do not freeze.

#### DO NOT USE IF CAP RING IS BROKEN OR MISSING

#### **USE BY EXPIRATION DATE ON CARTON AND BOTTLE**



# Enfamil® Sterile Water – 946 mL

Sterile water for oral use only.

Please contact your Hospital Account Manager for product access and contract inclusion.

(I) Pareve (manufactured on dairy equipment) (M) IFANCA

# **INDICATION**

Water for oral use only.

# **INGREDIENT**

Water

# **PRODUCT FORMS**

Product	Format	Unit Size	Approximate Product Yield at Normal Dilution	Shelf Life	ltem Number	Availability
Enfamil® Sterile Water – 946 mL	Liquid	946 mL	946 mL/unit	12 months	3130303	Hospital only

Nutrients - N/A

# PREPARATION OF FEEDINGS

Your baby's health depends on carefully following the instructions below.

Proper hygiene, preparation, use and storage are important when preparing infant feedings. Use as directed by your baby's doctor.

Inspect each bottle for signs of damage.

This product is for use in preparing feeds in centralized preparation rooms.

Upon opening, either feed immediately or replace cap and store in refrigerator at 2-4°C (35-40°F) for no longer than 24 hours. After feeding begins, do not refrigerate feeding bottle. You must use within 1 hour or discard.

**WARNING:** Do not use a microwave oven to warm water. Serious burns may result.



WARNING: Not for parenteral (I.V.) use.

This water is ready for use as drinking water or for other oral purposes as directed by your baby's doctor.

**STORAGE:** Store unopened bottles at room temperature. Avoid excessive heat. Do not freeze.

DO NOT USE IF CAP RING IS BROKEN OR MISSING

**USE BY EXPIRATION DATE ON BOTTLE** 



# Enfamil® 5% Glucose in Water

5% glucose in water. For oral use only.

Pareve (manufactured on dairy equipment) (M) IFANCA

# **INDICATION**

5% glucose in water. For oral use only.

# **INGREDIENTS**

Water, Dextrose

# **NUTRIENTS**

Nutrition Facts	per 1 Nursette® bottle (59 mL)
Calories	10
Fat	0 g
Carbohydrate	3 g
Sugars	3 g
Protein	0 g

Not a significant source of sodium, fibre, vitamin A, vitamin C, calcium or iron

# **PRODUCT FORMS**

Product	Format	Unit Size	Approximate Product Yield at Normal Dilution	Shelf Life	ltem Number	Availability
Enfamil® 5% Glucose in Water	Nursette® Bottle	59 mL	59 mL (2 fl oz)	21 Months	1216351	Hospital only



### PREPARATION OF FEEDINGS

Your baby's health depends on carefully following the instructions below.

Proper hygiene, preparation, use and storage are important when preparing infant feedings. Use as directed by your baby's doctor.

Inspect each bottle for signs of damage.



1. Wash hands thoroughly with soap and water before preparing bottle for feeding.



2. Remove cap.



3. Attach **DISPOSABLE NIPPLE UNIT** (not included) or ask your baby's doctor about the need to boil a clean nipple in water before use.

Upon opening, either feed immediately or replace cap and store in refrigerator at 2-4°C (35-40°F) for no longer than 24 hours. After feeding begins, do not refrigerate feeding bottle. You must use within one hour or discard.

**WARNING:** Do not use a microwave oven to warm formula. Serious burns may result.

**STORAGE:** Store unopened bottles at room temperature. Avoid excessive heat. Do not freeze.

WARNING: Not for parenteral (I.V.) use.

DO NOT USE IF CAP RING IS BROKEN OR MISSING

**USE BY EXPIRATION DATE ON CARTON AND BOTTLE** 



Date published 11/2024 – Please reference Mead Johnson Nutrition's online Product Handbook for the most up to date product details (www.enfamil.ca/medical).



# Enfagrow A<sup>+®</sup>

Cow's milk-based, nutritional supplement designed for children 1-5 years of age.

OD (M) IFANCA

# **INDICATION**

Cow's milk-based, nutritional supplement designed for children 1-5 years of age. Contains 23 vitamins and minerals including age appropriate levels of calcium and iron, as well as DHA (a type of Omega-3 fat) and two dietary fibres (galactooligosaccharides and polydextrose).

# **NUTRIENTS**

Analysis	per 200 mL Serving Reconstituted (36 g or 3 unpacked level scoops of powder)	Amount per serving (Ready to Drink)
Energy (kcal/kJ)	160/670	160/670
Protein (g)	6.4	6.4
Fat (g)	6.3	5.2
Linoleic Acid (g)	0.9	0.8
Linolenic Acid (g)	0.1	0.1
Saturated (g)	2.6	2.1
Docosahexaenoic acid (mg)	26	26
Carbohydrate (g)	20	22
Galactooligosaccharides (g)	0.4	0.4
Polydextrose (g)	0.4	0.4
Sodium (mg)	80	95
Potassium (mg)	300	300
Vitamin A (RE)	250	177
Vitamin D (mg)	0.0009	0.0009
Vitamin E (mg)	1.81	1.81
Vitamin C (mg)	10	10
Thiamine (mg)	0.3	0.3



Analysis	per 200 mL Serving Reconstituted (36 g or 3 unpacked level scoops of powder)	Amount per serving (Ready to Drink)
Riboflavin (mg)	0.36	0.36
Niacin (mg)	5.8	5.4
Pantothenic Acid (mg)	1.13	1.13
Vitamin B <sub>6</sub> (mg)	0.36	0.36
Folacin (mg)	0.057	0.057
Vitamin B <sub>12</sub> (mg)	0.0002	0.0002
Biotin (mg)	0.03	0.03
Calcium (mg)	250	220
Phosphorus (mg)	187	187
Magnesium (mg)	38	38
Iron (mg)	2.5	2.5
Zinc (mg)	2.6	2.6
Manganese (mg)	0.86	0.86
Copper (mg)	0.3	0.3
lodine (mg)	0.015 (milk flavour) 0.024 (vanilla flavour)	0.04
Chloride (mg)	180	180

% Daily Value		
	Children less than 2 years old	Children 2 years of age and older
Calcium	50% (Powder) 40% (RTD)	25% (Powder) 20% (RTD)
Iron	35% (Powder) 40% (RTD)	20%

Nutrient values are subject to change. Consult actual product labels for most current information.

# **INGREDIENTS**

# **VANILLA FLAVOUR POWDER**

Skim Milk Powder, Sugars (Corn Syrup Solids, Sugar), Palm Olein Oil, Coconut Oil, Soy Oil, High Oleic Sunflower Oil, Galactooligosaccharide Solids (GOS), Polydextrose, Microencapsulated Tuna Oil\* (Tuna Oil, Sodium Caseinate, Dried

Glucose Syrup, Dextrose, Sodium Ascorbate, Tocopherols, Soy Lecithin), Natural and Artificial Vanilla Flavour, Soy Lecithin, Ascorbyl Palmitate, (May Contain Potassium Hydroxide), **Minerals** (Calcium Carbonate, Calcium Phosphate, Cupric Sulfate, Ferrous Sulfate, Magnesium Phosphate, Manganese Sulfate and Zinc Sulfate), **Vitamins** (Ascorbic Acid, Biotin, Calcium D-Pantothenate, Dl- $\alpha$ -Tocopheryl Acetate, Folic Acid, Niacinamide, Pyridoxine Hydrochloride, Riboflavin, Sodium Ascorbate, Thiamine Hydrochloride, Vitamin A Palmitate and Vitamin D<sub>3</sub>).

# MILK FLAVOUR POWDER

\* A source of DHA.

Skim Milk Powder, Sugars (Corn Syrup Solids), Palm Olein Oil, Coconut Oil, Soy Oil, High Oleic Sunflower Oil, Galactooligosaccharide Solids (GOS), Polydextrose, Microencapsulated Tuna Oil\* (Tuna Oil, Sodium Caseinate, Dried Glucose Syrup, Dextrose, Sodium Ascorbate, Tocopherols and Soy Lecithin), Soy Lecithin, Natural Flavor, Ascorbyl Palmitate (May Contain Potassium Hydroxide), **Minerals** (Calcium Carbonate, Calcium Phosphate, Cupric Sulfate, Ferrous Sulfate, Magnesium Phosphate, Manganese Sulfate and Zinc Sulfate), **Vitamins** (Ascorbic Acid, Biotin, Calcium D-Pantothenate, Dl- $\alpha$ -Tocopheryl Acetate, Folic Acid, Niacinamide, Pyridoxine Hydrochloride, Riboflavin, Sodium Ascorbate, Thiamine Hydrochloride, Vitamin A Palmitate and Vitamin D3).

\* A source of DHA

#### **VANILLA FLAVOUR READY TO FEED**

Water, Nonfat Milk Powder, Corn Syrup Solids, Palm Olein Oil, Sugar, Soy Oil, Coconut Oil, High Oleic Sunflower Oil, Polydextrose, Galactooligosaccharide Solids (GOS), Soy Lecithin, Mono- And Diglycerides, Natural and Artificial Flavours, Crypthecodinium Cohnii Oil\*, Carrageenan, Cellulose Gum, Cellulose Gel, Ascorbyl Palmitate, Mixed Tocopherols (May Contain Potassium Hydroxide), **Minerals** (Calcium Phosphate, Cupric Sulfate, Ferrous Sulfate, Magnesium Phosphate, Manganese Sulfate, Potassium Chloride And Zinc Sulfate), **Vitamins** (Ascorbic Acid, Biotin, Calcium D-Pantothenate, DL- $\alpha$ -Tocopheryl Acetate, Folic Acid, Niacinamide, Pyridoxine Hydrochloride, Riboflavin, Thiamine Hydrochloride, Vitamin A Palmitate and Vitamin D3).

\* A source of DHA.

#### MILK FLAVOUR READY TO FEED

Water, Nonfat Milk Powder, Corn Syrup Solids, Palm Olein Oil, Soy Oil, Coconut Oil, High Oleic Sunflower Oil, Polydextrose, Galactooligosaccharide Solids (GOS), Soy Lecithin, Natural Flavour, Mono- And Diglycerides, Crypthecodinium Cohnii Oil\*, Carrageenan, Cellulose Gum, Cellulose Gel, Ascorbyl Palmitate, Mixed Tocopherols, (May Contain Potassium Hydroxide), **Minerals** (Calcium Phosphate, Cupric Sulfate, Ferrous Sulfate, Magnesium Phosphate, Manganese Sulfate, Potassium Chloride And Zinc Sulfate), **Vitamins** (Ascorbic Acid, Biotin, Calcium D-Pantothenate, DL- $\alpha$ -Tocopheryl Acetate, Folic Acid, Niacinamide, Pyridoxine Hydrochloride, Riboflavin, Thiamine Hydrochloride, Vitamin A Palmitate and Vitamin D3).

\* A source of DHA.

## **PRODUCT FORMS**

Product	Format	Unit Size	Approximate Product Yield at Normal Dilution	Shelf Life	Item Number	Availability
Enfagrow A <sup>+®</sup> Toddler	Powder	907 g	25 x 200 mL	18 Months	3181752	Retail only
Nutritional Drink, Vanilla Flavour	, Vanilla Ready to 237 L 1 hottle x	12 Months	2053715	Retail only		
Enfagrow A <sup>+®</sup> Toddler	Powder	907 g	26 x 200 mL	18 Months	3181749	Retail only
Nutritional Drink, Milk Flavour	Ready to Drink	237 mL	237 mL (1 bottle x 8 fl oz )	12 Months	2053714	Retail only

## PREPARATION OF FEEDINGS

### **POWDER**

Enfagrow  $A^{+\otimes}$  is for children 1-5 years old and not intended for infants under 1 year of age.



**Storage:** After opening can, store tightly covered at room temperature in a dry area and use contents within 1 month. Store **DRY** scoop in the can.

### **USE BY EXPIRATION DATE ON BOTTOM OF CAN**

### **READY TO DRINK**

Shake well before opening. Remove protective seal around bottle cap, remove bottle cap and foil seal, and serve. Refrigerate after opening and use within 48 hours.

**WARNING:** Do not use a microwave oven to warm formula. Serious burns may result.

**Storage:** Store unopened bottles at room temparature. Avoid excessive heat and prolonged exposure to light. Do not freeze.

DO NOT ACCEPT IF PACKAGE HAS BEEN OPENED. DO NOT USE IF PROTECTIVE SEAL AROUND BOTTLE CAP, OR IF FOIL SEAL UNDER BOTTLE CAP, IS DAMAGED OR MISSING.

**USE BY EXPIRATION DATE ON CARTON AND BOTTLE** 



Date published 11/2024 – Please reference Mead Johnson Nutrition's online Product Handbook for the most up to date product details (www.enfamil.ca/medical).



# Portagen®

A milk-protein based powder with medium-chain triglycerides (MCT) for adults and children.

OD (M) IFANCA

## **INDICATION**

A milk-protein based powder with medium-chain triglycerides (MCT) for adults and children.

Analysis	per 100 mL	per 100 g	Per 946 mL 30 kcal/30 mL
Energy (kcal/kJ)	101/420	470/1970	960/4020
Protein (g)	3.7	17	35
% of total energy	14	14	14
Source: Sodium caseinate			
Carbohydrate (g)	11.6	54	110
% of total energy	46	46	46
Source: Corn syrup solids, sug	ar		
Fat (g)	4.7	22	45
% of total energy	40	40	40
Source: MCT (87%), corn oil (13	3%)		
Linoleic Acid (g)	0.28	1.3	2.7
Minerals			
Calcium (mg)	95	440	900
Phosphorus (mg)	71	330	670
Magnesium (mg)	21	96	200
Iron (mg)	1.89	8.8	17.9
Zinc (mg)	0.95	4.4	9
Manganese (mg)	0.127	0.59	1.2
Copper (mg)	0.159	0.74	1.51
lodine (mg)	0.0071	0.033	0.067





Analysis	per 100 mL	per 100 g	Per 946 mL 30 kcal/30 mL
Selenium (mg)	n/a	n/a	n/a
Sodium (mg)	52	240	490
Potassium (mg)	127	590	1200
Chloride (mg)	86	400	820
Vitamins			
Vitamin A (IU)	340	1560	3200
Vitamin D (IU)	28	130	270
Vitamin E (IU)	1.59	7.4	15.1
Vitamin K (mg)	0.008	0.037	0.075
Vitamin C (mg)	3.9	18	37
Thiamine (mg)	0.08	0.37	0.75
Riboflavin (mg)	0.095	0.44	0.9
Niacin (mg)	1.05	4.9	10
Pantothenic Acid (mg)	0.54	2.5	5.1
Vitamin B <sub>6</sub> (mg)	0.105	0.49	1
Folic Acid (mg)	0.0101	0.047	0.0960
Vitamin B <sub>12</sub> (mg)	0.00032	0.0015	0.0031
Biotin (mg)	0.0039	0.018	0.037
Choline (mg)	13.5	63	128
Inositol (mg)	nd	nd	nd
Carnitine (mg)	2.9	13.3	27
Taurine (mg)	n/a	n/a	n/a
Other Characteristics			

Other Characteristics	
Potential Renal Solute Load (m0sm/100 mL) <sup>1</sup>	31
Osmolality (mOsm/kg H <sub>2</sub> O)	350
Osmolarity (mOsm/L)	300
Water (g/100 mL)	87

n/a = None added

nd = Not declared

Portagen® should be used only under the supervision of a physician or dietitian.

This product is not nutritionally complete. If used long term, supplementation of essential fatty acids and ultra-trace minerals should be considered.

<sup>1.</sup> Fomon SJ, Ziegler EE. Renal solute load and potential renal solute load in infancy. *J Pediatr.* 1999;134:11-14. Nutrient values are subject to change. Consult actual product labels for most current information.

### **INGREDIENTS**

Corn Syrup Solids, Medium Chain Triglycerides (Fractionated Coconut Oil), Sodium Caseinate, Sugar, Corn Oil, Soy Lecithin, L-Carnitine, (May Contain Potassium Hydroxide And/Or Hydrochloric Acid), **Minerals** (Calcium Citrate, Calcium Phosphate, Cupric Sulfate, Ferrous Sulfate, Magnesium Phosphate, Manganese Sulfate, Potassium Chloride, Potassium Citrate, Sodium Citrate, Sodium Iodide And Zinc Sulfate), **Vitamins** (Ascorbic Acid, Biotin, Calcium D-Pantothenate, Choline Chloride, DL- $\alpha$ -Tocopheryl Acetate, Folic Acid, Niacinamide, Pyridoxine Hydrochloride, Riboflavin, Thiamine Hydrochloride, Vitamin A Palmitate, Vitamin B<sub>12</sub>, Vitamin D<sub>3</sub> And Vitamin K<sub>1</sub>).

## PRODUCT FORMS

Product	Format	Unit Size	Approximate Product Yield at Normal Dilution	Shelf Life	Item Number	Availability
Portagen®	Powder	410 g	1.9 L @ 30 kcal/ 30 mL	24 Months	2032499	Hospital

### PREPARATION OF FEEDINGS

USE AS DIRECTED BY YOUR PHYSICIAN OR DIETITIAN.

### PORTAGEN® POWDER IS NOT DESIGNED FOR USE AS AN INFANT FORMULA.

Portagen® powder is a readily-digestible protein, fat and carbohydrate source that can be made into a nutritional beverage to supplement the diet.

Portagen® powder is not intended as a sole source of nutrition. For chronic (long-term) use: Supplementation of essential fatty acids and ultra trace minerals should be considered. To be used only as directed by your doctor.

WARNING: The consumer's health depends on carefully following the instructions below. Use only as directed by a medical professional. Improper hygiene, preparation, dilution, use or storage may result in severe harm. Nutritional powders are not sterile and should <u>not</u> be fed to premature infants or those persons who might have immune problems unless directed and supervised by a medical professional.

- For 946 mL (32 US fl oz) of prepared product, add 1 2/3 packed level measuring cups (204 g) of powder to 790 mL (27 US fl oz) of water.\*
- For approximately 131 mL (4.5 US fl oz) of prepared product, add 3 packed level scoops (28.2 g) to 110 mL (3.5 US fl oz) of water.\*
- Both preparations will result in 30 kcal/30 mL of prepared products. For other volumes prepare as directed by your doctor.
- Pour desired amount of water into a suitable container. Add powder and mix vigorously until thoroughly blended.

This 410 g can of PORTAGEN® will make approximately 1.9 L (2 US quarts) of beverage at 30 kcal/30 mL.\*

Failure to follow these instructions could result in severe harm. Once prepared, product can spoil quickly. Either consume immediately or cover and store in refrigerator at 2-4°C (35-40°F) for no longer than 24 hours. Shake well before each use. Do not use prepared product if it is unrefrigerated for more than a total of 2 hours. Do not freeze prepared product. After feeding begins, do not refrigerate the container. You must use within 1 hour or discard.

**Powder Storage:** Store cans at room temperature. After opening can, keep tightly covered, store in dry area and use contents within 1 month. Do not freeze powder and avoid excessive heat.

WARNING: Not for parenteral (I.V.) use.

\* Using mL measurements are the most accurate for meeting target caloric density; US fl oz values are approximations and are based on US conversion factors.

### **USE BY EXPIRATION DATE ON BOTTOM OF CAN**



# Enfamil® Enfalyte®

## Oral Electrolyte Solution

Oral electrolyte maintenance solution. Light cherry flavour.

(M) Pareve (manufactured on dairy equipment) (M) IFANCA

## **INDICATION**

Oral electrolyte maintenance solution. Helps promote fluid absorption.

Analysis	per 100 mL
Energy (kcal/kJ)	12.6/53
Protein (g)	0
Carbohydrate (g)	3.2
% of total energy	100
Source: Corn syrup solids (94%), c	itrates (6%)
Fat (g)	0

Electrolytes	Per 100 mL	g/litre	mMol/litre
Sodium (mg) (mMol)	115 5	1.15	50
Potassium (mg) (mMol)	98 2.5	0.98	25
Citrate	-	2.10	11
Chloride (mg) (mMol)	160 4.5	1.60	45

Other Characteristics	
Potential Renal Solute Load (m0sm/100 mL) <sup>1</sup>	12
Osmolality (mOsm/kg H <sub>2</sub> O)	200
Osmolarity (mOsm/L)	200
Water content (g/100 mL)	98

<sup>1.</sup> Fomon SJ, Ziegler EE. Renal solute load and potential renal solute load in infancy. *J Pediatr.* 1999;134:11-14. Nutrient values are subject to change. Consult actual product labels for most current information.





## **INGREDIENTS**

**Medicinal Ingredients:** Potassium Citrate, Sodium Chloride, Citric Acid, Sodium Citrate.

Non-Medicinal Ingredients: Water, Corn Syrup Solids and Natural Flavour.

## **PRODUCT FORMS**

Product	Format	Unit Size	Approximate Product Yield at Normal Dilution	Shelf Life	Item Number	Availability
Enfamil® Enfalyte® Oral Electrolyte Solution	Bottle	177 mL	177 mL	12 Months	2021478	Hospital only

Your baby's health depends on carefully following the instructions below. Proper hygiene, preparation, use and storage are important when preparing infant feedings. Use as directed by your baby's doctor. Inspect each bottle for signs of damage.

## PREPARATION OF FEEDINGS

#### **RTF**



Caution: Do not use a microwave oven to warm formula. Serious burns may result.

Failure to follow these instructions could result in severe harm. Opened bottles and prepared bottles can spoil quickly. Upon opening, either feed immediately or replace cap and store in refrigerator at 2-4°C (35-40°F) for no longer than 48 hours. After feeding begins, do not refrigerate opened bottle. You must use within 1 hour or discard.

**Storage:** Store unopened bottles at room temperature. Avoid excessive heat. Do not freeze.

**WARNING:** Do not mix with infant formula, milk, fruit juices or other electrolytecontaining liquids. Not for parenteral (I.V.) use.

### DO NOT USE IF CAP RING IS BROKEN OR MISSING

### **USE BY EXPIRATION DATE ON BOTTLE**

Oral feedings of Enfamil® Enfalyte® may be used for maintenance of water and electrolytes lost during diarrhea and vomiting in infants, children and adults.

**NOTE:** ENFALYTE ORAL ELECTROLYTE SOLUTION HAS AN OSMOLALITY OF APPROXIMATELY 200 m0sm/kg OF WATER AND CONTAINS 31.5 g OF CORN SYRUP SOLIDS PER LITER. EACH 100 mL OF ENFALYTE SUPPLIES ABOUT 12 CALORIES; EACH FL OZ, ABOUT 3.5 CALORIES.

THIS PRODUCT SHOULD BE USED ONLY AS DIRECTED BY YOUR BABY'S DOCTOR. IF VOMITING OR FEVER IS PRESENT, OR DIARRHEA CONTINUES BEYOND 24 HOURS, CONSULT YOUR BABY'S DOCTOR.





Date published 11/2024 – Please reference Mead Johnson Nutrition's online Product Handbook for the most up to date product details (www.enfamil.ca/medical).



# Phenyl-Free® 1

Phenylalanine-free, iron fortified infant formula with DHA and ARA for infants and young children with documented phenylketonuria (PKU).

O D (M) IFANCA

## **INDICATION**

Phenylalanine-free, iron fortified infant formula with DHA and ARA for infants and young children with documented phenylketonuria (PKU).

Analysis	per 100 mL	per 100 g			
Energy (kcal/kJ)	68/280	500/2100			
Protein equivalent (g)	2.2	16.2			
% of total energy	13	13			
Source: Amino acid premix					
Carbohydrate (g)	6.9	51			
% of total energy	40	40			
Source: Corn syrup solids, modified corn	starch, sugar				
Fat (g)	3.5	26			
% of total energy	47	47			
Source: Palm olein oil (44%), soy oil (19.5%), coconut oil (19.5%), high oleic sunflower oil (14.5%), ARA & DHA single cell oil blend (2.5%)					
Linoleic Acid (g)	0.54	4			
Linolenic Acid (g)	0.046	0.34			
ARA (mg)	24	175			
DHA (mg)	11.9	88			
Minerals					
Calcium (mg)	89	660			
(mMol)	2.2	16.5			
Phosphorus (mg)	59	440			
(mMol)	1.92	14.2			
Magnesium (mg)	8.9	66			
(mMol)	0.37	2.7			





Analysis	per 100 mL	per 100 g
Iron (mg)	1.3	9.6
Zinc (mg)	1.16	8.6
Manganese (mg)	0.051	0.38
Copper (mg)	0.116	0.86
lodine (mg)	0.0103	0.076
Selenium (mg)	0.0019	0.0141
Sodium (mg) (mMol)	32 1.41	240 10.4
Potassium (mg) (mMol)	76 1.93	560 14.3
Chloride (mg) (mMol)	58 1.64	430 12.1
Vitamins		
Vitamin A (IU)	210	1520
Vitamin D (IU)	51	380
Vitamin E (IU)	1.35	10
Vitamin K (mg)	0.0059	0.044
Vitamin C (mg)	8.1	60
Thiamine (mg)	0.135	1
Riboflavin (mg)	0.135	1
Niacin (mg)	1.35	10
Pantothenic Acid (mg)	0.51	3.8
Vitamin B <sub>6</sub> (mg)	0.135	1
Folic Acid (mg)	0.0135	0.1
Vitamin B <sub>12</sub> (mg)	0.00027	0.002
Biotin (mg)	0.0051	0.038
Choline (mg)	16.7	124
Inositol (mg)	11.6	86
Carnitine (mg)	6.9	51
Taurine (mg)	4.1	30

Other Characteristics	
Potential Renal Solute Load (m0sm/100 mL) <sup>1</sup>	19.5
Osmolality (mOsm/kg H <sub>2</sub> O)*	350
Osmolarity (mOsm/L)*	320
Water (g/100 mL)	90

All metabolic products are for use under the direct and continuing supervision of a doctor or metabolic dietitian.

This product is nutritionally incomplete. Care must be taken to provide enough phenylalanine to support growth, using other foods with this amino acid as required.

Nutritional powders are not sterile.

- \* 100 mL values and Other Characteristics information is based on this product mixed with water to a caloric density of 0.68 kcal/mL.
- 1. Fomon SJ, Ziegler EE. Renal solute load and potential renal solute load in infancy. J Pediatr. 1999;134:11-14.

Nutrient values are subject to change. Consult actual product labels for most current information.

### **INGREDIENTS**

Corn Syrup Solids, Vegetable Oil (Palm Olein Oil, Coconut Oil, Soy Oil, High Oleic Sunflower Oil), Amino Acids (L-Glutamine, L-Leucine, Potassium Aspartate, L-Lysine Hydrochloride, L-Tyrosine, L-Proline, L-Valine, L-Isoleucine, L-Alanine, L-Arginine, L-Threonine, L-Serine, Glycine, L-Histidine, L-Methionine, L-Tryptophan, L-Cystine), Modified Corn Starch, Sugar, Mortierella Alpina Oil†, Crypthecodinium Cohnii Oil‡, Ethyl Vanillin, L-Carnitine, Taurine, Ascorbyl Palmitate, Mixed Tocopherols (May Contain Potassium Hydroxide), Minerals (Calcium Phosphate, Choline Chloride, Cupric Sulfate, Ferrous Sulfate, Inositol, Magnesium Oxide, Manganese Sulfate, Potassium Chloride, Potassium Citrate, Sodium Citrate, Sodium Iodide, Sodium Selenite And Zinc Sulfate), Vitamins (Ascorbic Acid, Biotin, Calcium Pantothenate, Folic Acid, Niacinamide, Riboflavin, Thiamine Hydrochloride, Vitamin A Palmitate, Vitamin B<sub>12</sub>, Vitamin B<sub>6</sub> Hydrochloride, Vitamin D<sub>3</sub>, Vitamin E Acetate And Vitamin K<sub>1</sub>).

- † A source of Arachidonic Acid (ARA).
- ‡ A source of Docosahexaenoic Acid (DHA).

### PRODUCT FORMS

Product	Format	Unit Size	Approximate Product Yield at Normal Dilution	Shelf Life	Item Number	Availability
Phenyl-Free® 1	Powder	454 g	Yield Not Applicable to Metabolics	18 Months	2001922	Hospital & Retail Pharmacy (on prescription)

### PREPARATION OF FEEDINGS

Mead Johnson Phenyl-Free® 1 is to be used only in the dietary management of infants and young children with documented phenylketonuria under direct and continuing medical supervision.

WARNING: This product is nutritionally incomplete. Care must be taken to provide enough phenylalanine to support growth, using other foods with phenylalanine as required. Medical professionals must carefully and constantly supervise the use of Phenyl-Free® 1 with other foods and liquids and adjust the diet based on frequent blood tests.

### Your baby's health depends on carefully following the instructions below.

Proper hygiene, preparation, dilution, use and storage are important when preparing formula. Powdered products are not sterile and should **not** be fed to premature infants or those persons who might have immune problems unless directed and supervised by your baby's doctor.



1. Wash hands thoroughly with soap and water before preparing formula. Boil clean bottles, nipples, caps, scoop and utensils in water (2 minutes at a rolling boil).



2. Boil fresh water at a rolling boil for 2 minutes. Cool to room temperature prior to mixing.



3. Pour desired amount of the cooled water into the bottle. Add powder. Cap bottle and **SHAKE WELL**.

Your baby's doctor will provide instructions for the correct amount of cooled, boiled water and powder. If instructed to use the scoop in the can, each scoop (unpacked and level) delivers approximately 4.5 g of powder. Store the DRY scoop in this can.

Failure to follow these instructions could result in severe harm. Once prepared, formula can spoil quickly. Either feed immediately or cover and store in refrigerator at 2-4°C (35-40°F) for no longer than 24 hours. Do not use prepared formula if it is unrefrigerated for more than a total of 2 hours. Do not freeze prepared formula. After feeding begins, use within 1 hour or discard.

**WARNING:** Do not use a microwave oven to warm formula. Serious burns may result.

**Powder Storage:** Store cans at room temperature. After opening can, keep tightly covered, store in dry area and use contents within 1 month. Do not freeze powder and avoid excessive heat. **Use by date on bottom of can.** 



## Phenyl-Free® 2

Phenylalanine-free powder for dietary management of children and adults (3 years and older) with phenylketonuria (PKU).

(I) D (M) IFANCA

## **INDICATION**

Phenylalanine-free powder for dietary management of children and adults (3 years and older) with phenylketonuria (PKU).

Analysis	per 100 mL	per 100 g
Energy (kcal/kJ)	85/360	410/1720
Protein equivalent (g)	4.6	22
% of total energy	22	22
Source: Amino Acid Premix		
Carbohydrate (g)	12.4	60
% of total energy	59	59
Source: Sugar, corn syrup solids, modifie	ed corn starch	
Fat (g)	1.78	8.6
% of total energy	19	19
Source: Soy oil		
Linoleic Acid (g)	0.87	4.2
Linolenic Acid (g)	0.108	0.52
Minerals		
Calcium (mg) (mMol)	151 3.8	730 18.2
Phosphorus (mg) (mMol)	151 4.9	730 24
Magnesium (mg) (mMol)	34 1.39	163 6.7
Iron (mg)	2.5	12.2
Zinc (mg)	2.5	12.2





Analysis	per 100 mL	per 100 g
Manganese (mg)	0.27	1.31
Copper (mg)	0.25	1.22
lodine (mg)	0.013	0.063
Selenium (mg)	0.006	0.029
Sodium (mg) (mMol)	126 5.5	610 27
Potassium (mg) (mMol)	230 5.8	1100 28
Chloride (mg) (mMol)	178 5	860 24
Vitamins		
Vitamin A (IU)	300	1430
Vitamin D (IU)	60	290
Vitamin E (IU)	2	9.8
Vitamin K (mg)	0.0085	0.041
Vitamin C (mg)	10.1	49
Thiamine (mg)	0.25	1.22
Riboflavin (mg)	0.2	0.98
Niacin (mg)	4.6	22
Pantothenic Acid (mg)	1.01	4.9
Vitamin B <sub>6</sub> (mg)	0.2	0.98
Folic Acid (mg)	0.072	0.35
Vitamin B <sub>12</sub> (mg)	0.0005	0.0024
Biotin (mg)	0.0101	0.049
Choline (mg)	20	98
Inositol (mg)	10.1	49
Carnitine (mg)	10.1	49
Taurine (mg)	10.1	49
Other Characteristics		
Potential Renal Solute Load (m0sm/100	47	
Osmolality (mOsm/kg H <sub>2</sub> O)*	920	
Osmolarity (mOsm/L)*		790
Water (g/100 mL)	86	

Use only as directed by a medical professional.

Powder medical foods are not sterile.

WARNING: This protein is incomplete since it does not contain the essential amino acid phenylalanine.

- \* 100 mL values and Other Characteristics information is based on this product mixed with water to a caloric density of 0.85 kcal/mL.
- 1. Fomon SJ, Ziegler EE. Renal solute load and potential renal solute load in infancy. J Pediatr. 1999;134:11-14.

Nutrient values are subject to change. Consult actual product labels for most current information.

### **INGREDIENTS**

Sugar, Amino Acids (L-Glutamine, L-Leucine, Potassium Aspartate, L-Lysine Hydrochloride, L-Tyrosine, L-Proline, L-Valine, L-Isoleucine, L-Alanine, Larginine, L-Threonine, L-Serine, Glycine, L-Histidine, L-Methionine, L-Tryptophan, L-Cystine), Corn Syrup Solids, Soy Oil, Modified Corn Starch, Calcium Phosphate, Sodium Citrate, Magnesium Phosphate, Potassium Chloride, And Less Than 1%: Sodium Phosphate, Potassium Citrate, Choline Chloride, Ascorbic Acid, Taurine, Inositol, Ferrous Sulfate, L-Carnitine, Zinc Sulfate, Niacinamide, Vitamin E Acetate, Vitamin A Palmitate, Calcium Pantothenate, Cupric Sulfate, Manganese Sulfate, Vitamin B<sub>12</sub>, Thiamin Hydrochloride, Vitamin B<sub>6</sub> Hydrochloride, Riboflavin, Folic Acid, Vitamin D<sub>3</sub>, Chromic Chloride, Sodium Molybdate, Sodium Iodide, Sodium Selenite, Biotin, Vitamin K<sub>1</sub>, Ethyl Vanillin.

## **PRODUCT FORMS**

Product	Format	Unit Size	Approximate Product Yield at Normal Dilution	Shelf Life	Item Number	Availability
Phenyl-Free® 2	Powder	454 g	Yield Not Applicable to Metabolics	24 Months	1281335	Hospital & Retail Pharmacy (on prescription)

### PREPARATION OF FEEDINGS

**Mead Johnson Phenyl-Free® 2** is to be used only in the dietary management of children and adults with documented Phenylketonuria while under direct and continuing medical supervision.

**WARNING:** This product is nutritionally incomplete. Continuing medical supervision and frequent blood tests are required.

**Phenyl-Free® 2** must be supplemented using other food sources of protein and fluid to provide enough phenylalanine to support dietary requirements of children and adults.

### INSTRUCTIONS FOR PREPARATION AND USE

Your medical professional will provide the correct amount of powder to mix with water for consumption.\*

It is important to follow the directions below.

Measure the correct amount of water into a suitable container for mixing. Then add the required amount of **Phenyl-Free® 2. Mix well until blended.** Consume the prepared beverage immediately or cover and refrigerate. Use within 48 hours of preparation. Mix before drinking.

\* If instructed to use the scoop in the can, each unpacked, level scoop delivers **approximately 14.4 g** of powder. Store the **DRY** scoop in this can.

**Powder Storage:** Store cans at room temperature. After opening can, keep tightly covered, store in a dry area and use. **Use by date on bottom of can.** 



# Phenyl-Free® 2 HP

High protein phenylalanine-free powder for dietary management of children and adults (3 years and older) with phenylketonuria (PKU).

(U) D (M) IFANCA

## INDICATION

High protein phenylalanine-free powder for dietary management of children and adults (3 years and older) with phenylketonuria (PKU).

Analysis	per 100 mL	per 100 g
Energy (kcal/kJ)	85/360	390/1630
Protein equivalent (g)	8.6	40
% of total energy	41	41
Source: Amino acid premix		
Carbohydrate (g)	9.5	44
% of total energy	44	44
Source: Sugar, corn syrup solids, amino	acid premix, modifi	ed corn starch
Fat (g)	1.36	6.3
% of total energy	15	15
Source: Soy oil		
Linoleic Acid (g)	0.65	3
Linolenic Acid (g)	0.082	0.38
Minerals		
Calcium (mg)	210	980
(mMol)	5.3	24
Phosphorus (mg)	210	980
(mMol)	6.8	32
Magnesium (mg)	63	290
(mMol)	2.6	11.9
Iron (mg)	3.4	15.7
Zinc (mg)	3.4	15.7
Manganese (mg)	0.34	1.57
Copper (mg)	0.34	1.57





Analysis	per 100 mL	per 100 g
lodine (mg)	0.0108	0.05
Selenium (mg)	0.0078	0.036
Sodium (mg)	89	410
(mMol)	3.9	17.8
Potassium (mg) (mMol)	250 6.5	1180 30
Chloride (mg) (mMol)	210 6	980 28
Vitamins		
Vitamin A (IU)	420	1960
Vitamin D (IU)	84	390
Vitamin E (IU)	2.5	11.8
Vitamin K (mg)	0.011	0.051
Vitamin C (mg)	13.6	63
Thiamine (mg)	0.34	1.57
Riboflavin (mg)	0.28	1.29
Niacin (mg)	6.3	29
Pantothenic Acid (mg)	1.36	6.3
Vitamin B <sub>6</sub> (mg)	0.28	1.29
Folic Acid (mg)	0.102	0.47
Vitamin B <sub>12</sub> (mg)	0.00067	0.0031
Biotin (mg)	0.0136	0.063
Choline (mg)	14.5	67
Inositol (mg)	13.6	63
Carnitine (mg)	7.8	36
Taurine (mg)	13.6	63
Other Characteristics		
Potential Renal Solute Load (m0sm/100	72	
Osmolality (m0sm/kg H <sub>2</sub> 0)	1240	
Osmolarity (m0sm/L)		1070
Water (g/100 mL)	86	

<sup>1.</sup> Fomon SJ, Ziegler EE. Renal solute load and potential renal solute load in infancy. J Pediatr. 1999;134:11-14.

### **INGREDIENTS**

Amino Acids (L-Glutamine, L-Leucine, Potassium Aspartate, L-Lysine Hydrochloride, L-Tyrosine, L-Proline, L-Valine, L-Isoleucine, L-Alanine, L-Arginine, L-Threonine, L-Serine, Glycine, L-Histidine, L-Methionine, L-Tryptophan, L-Cystine), Sugar, Corn Syrup Solids, Soy Oil, Modified Corn Starch, Calcium Phosphate, Magnesium Phosphate, And Less Than 1%: Sodium Citrate, Potassium Chloride, Sodium Phosphate, Potassium Citrate, Choline Chloride, Ascorbic Acid, Taurine, Inositol, Ferrous Sulfate, Zinc Sulfate, Niacinamide, L-Carnitine, Vitamin E Acetate, Vitamin A Palmitate, Calcium Pantothenate, Cupric Sulfate, Manganese Sulfate, Vitamin B<sub>12</sub>, Thiamin Hydrochloride, Vitamin B<sub>6</sub> Hydrochloride, Riboflavin, Folic Acid, Vitamin D<sub>3</sub>, Chromic Chloride, Sodium Molybdate, Sodium Selenite, Biotin, Vitamin K<sub>1</sub>, Sodium Iodide, Ethyl Vanillin.

## **PRODUCT FORMS**

Product	Format	Unit Size	Approximate Product Yield at Normal Dilution	Shelf Life	Item Number	Availability
Phenyl-Free® 2 HP	Powder	454 g	Yield Not Applicable to Metabolics	24 Months	1281334	Hospital & Retail Pharmacy (on prescription)

#### PREPARATION OF FEEDINGS

**Mead Johnson Phenyl-Free® 2 HP** is to be used only in the dietary management of children and adults with documented Phenylketonuria while under direct and continuing medical supervision.

**WARNING:** This product is nutritionally incomplete. Continuing medical supervision and frequent blood tests are required. **Phenyl-Free® 2 HP** must be supplemented using other food sources of protein and fluid to provide enough phenylalanine to support dietary requirements of children and adults.

### INSTRUCTIONS FOR PREPARATION AND USE

Your medical professional will provide the correct amount of powder to mix with water for consumption.\*

It is important to follow the directions below.

Measure the correct amount of water into a suitable container for mixing. Then add the required amount of **Phenyl-Free® 2 HP. Mix well until blended.** Consume the prepared beverage immediately or cover and refrigerate. Use within 48 hours of preparation. Mix before drinking.

<sup>\*</sup> If instructed to use the scoop in the can, each unpacked, level scoop delivers approximately 15.1 g of powder.

Store the DRY scoop in this can.

**Powder Storage:** Store cans at room temperature. After opening can, keep tightly covered, store in a dry area and use contents within 1 month.

Do not freeze powder and avoid excessive heat. Use by date on bottom of can.



## TYROS 1

Tyrosine-and phenylalanine-free, iron fortified infant formula with DHA and ARA for infants and young children with documented tyrosinemia type I and type II.

(I) D (M) IFANCA

## **INDICATION**

Tyrosine-and phenylalanine-free, iron fortified infant formula with DHA and ARA for infants and young children with documented tyrosinemia type I and type II.

Analysis	per 100 mL	per 100 g					
Energy (kcal/kJ)	68/280	500/2100					
Protein equivalent (g)	2.3	16.7					
% of total energy	13	13					
Source: Amino acid premix							
Carbohydrate (g)	6.9	51					
% of total energy	40	40					
Source: Corn syrup solids, modified corn starch, sugar							
Fat (g)	3.5	26					
% of total energy	47	47					
Source: Palm olein oil (44%), soy oil (19.5 sunflower oil (14.5%), ARA & DHA single							
Linoleic Acid (g)	0.54	4					
Linolenic Acid (g)	0.046	0.34					
ARA (mg)	24	175					
DHA (mg)	11.9	88					
Minerals							
Calcium (mg)	89	660					
(mMol)	2.2	16.5					
Phosphorus (mg)	59	440					
(mMol)	1.92	14.2					





Analysis	per 100 mL	per 100 g
Magnesium (mg)	8.9	66
(mMol)	0.37	2.7
Iron (mg)	1.3	9.6
Zinc (mg)	1.16	8.6
Manganese (mg)	0.051	0.38
Copper (mg)	0.116	0.86
lodine (mg)	0.0103	0.076
Selenium (mg)	0.0019	0.0141
Sodium (mg) (mMol)	32 1.41	240 10.4
Potassium (mg) (mMol)	82 2.1	610 15.6
Chloride (mg) (mMol)	58 1.64	430 12.1
Vitamins		
Vitamin A (IU)	210	1520
Vitamin D (IU)	51	380
Vitamin E (IU)	1.35	10
Vitamin K (mg)	0.0054	0.044
Vitamin C (mg)	8.1	60
Thiamine (mg)	0.135	1
Riboflavin (mg)	0.135	1
Niacin (mg)	1.35	10
Pantothenic Acid (mg)	0.51	3.8
Vitamin B <sub>6</sub> (mg)	0.135	1
Folic Acid (mg)	0.0135	0.1
Vitamin B <sub>12</sub> (mg)	0.00027	0.002
Biotin (mg)	0.0051	0.038
Choline (mg)	16.7	124
Inositol (mg)	11.6	86
Carnitine (mg)	6.8	50
Taurine (mg)	4.1	30

Other Characteristics	
Potential Renal Solute Load (m0sm/100 mL) <sup>1</sup>	20
Osmolality (m0sm/kg H <sub>2</sub> 0)*	360
Osmolarity (mOsm/L)*	320
Water (g/100 mL)	90

All metabolic products are for use under the direct and continuing supervision of a doctor or metabolic dietitian.

This product is nutritionally incomplete. Care must be taken to provide enough phenylalanine and tyrosine to support growth, using other foods with these amino acids as required. Nutritional powders are not sterile.

- \* 100 mL values and Other Characteristics information is based on this product mixed with water to a caloric density of 0.68 kcal/mL.
- 1. Fomon SJ, Ziegler EE. Renal solute load and potential renal solute load in infancy. *J Pediatr*. 1999;134:11-14. Nutrient values are subject to change. Consult actual product labels for most current information.

### **INGREDIENTS**

Corn Syrup Solids, Vegetable Oil (Palm Olein Oil, Coconut Oil, Soy Oil, High Oleic Sunflower Oil), Amino Acids (L-Glutamine, L-Leucine, Potassium Aspartate, L-Lysine Hydrochloride, L-Proline, L-Valine, L-Alanine, L-Isoleucine, L-Arginine, L-Threonine, L-Serine, Glycine, L-Histidine, L-Methionine, L-Tryptophan, L-Cystine), Modified Corn Starch, Sugar, Calcium Phosphate, Mortierella Alpine Oil, Crypthecodinium Cohnii Oil<sup>†</sup>, Ethyl Vanillin, L-Carnitine, Taurine, Ascorbyl Palmitate, Mixed Tocopherols (May Contain Potassium Hydroxide), **Minerals** (Choline Chloride, Cupric Sulfate, Ferrous Sulfate, Inositol, Magnesium Oxide, Manganese Sulfate, Potassium Chloride, Potassium Citrate, Sodium Citrate, Sodium Iodide, Sodium Selenite And Zinc Sulfate), **Vitamins** (Ascorbic Acid, Biotin, Calcium Pantothenate, Folic Acid, Niacinamide, Riboflavin, Thiamine Hydrochloride, Vitamin A Palmitate, Vitamin B<sub>12</sub>, Vitamin B<sub>6</sub> Hydrochloride, Vitamin E Acetate And Vitamin K<sub>1</sub>).

### PRODUCT FORMS

Product	Format	Unit Size	Approximate Product Yield at Normal Dilution	Shelf Life	Item Number	Availability
TYROS 1	Powder	454 g	Yield Not Applicable to Metabolics	18 Months	2001957	Hospital & Retail Pharmacy (on prescription)

### PREPARATION OF FEEDINGS

**Mead Johnson TYROS 1** is to be used only in the dietary management of infants and young children with documented tyrosinemia Type I and Type II while under direct and continuing medical supervision.

<sup>†</sup> A source of Docosahexaenoic Acid (DHA).

WARNING: This product is nutritionally incomplete. Care must be taken to provide enough phenylalanine and tyrosine to support growth, using other foods with these amino acids as required. Medical professionals must carefully and constantly supervise use of TYROS 1 with other foods and liquids and adjust the diet based on frequent blood tests.

#### INSTRUCTIONS FOR PREPARATION AND USE

### Your baby's health depends on carefully following the instructions below.

Proper hygiene, preparation, dilution, use and storage are important when preparing formula. Powdered formula products are not sterile and should **not** be fed to premature infants or those persons who might have immune problems unless directed and supervised by your baby's doctor.



1. Wash hands thoroughly with soap and water before preparing formula. Boil clean bottles, nipples, caps and utensils in water (2 minutes at a rolling boil).



2. Boil fresh water at a rolling boil for 2 minutes. Cool to room temperature prior to mixing.



3. Pour desired amount of the cooled water into the bottle. Add powder. Cap bottle and **SHAKE WELL.** 

Your baby's doctor will provide instructions for the correct amount of cooled, boiled water and powder. If instructed to use the scoop in the can, each scoop (unpacked and level) delivers approximately 4.5 g of powder. Store the DRY scoop in this can.

Failure to follow these instructions could result in severe harm. Once prepared, formula can spoil quickly. Either feed immediately or cover and store in refrigerator at 2-4°C (35-40°F) for no longer than 24 hours. Do not use prepared formula if it is unrefrigerated for more than a total of 2 hours. Do not freeze prepared formula. After feeding begins, use product within 1 hour or discard.

**WARNING:** Do not use a microwave oven to warm formula. Serious burns may result.

**STORAGE:** Store cans at room temperature. After opening can, keep tightly covered, store in dry area and use contents within one month. Do not freeze powder and avoid excessive heat. **Use by date on bottom of can.** 



## TYROS 2

Phenylalanine and tyrosine-free powder for dietary management of children and adults (3 years and older) with inborn errors of tyrosine metabolism including tyrosinemia type II.

O D (M) IFANCA

## **INDICATION**

Phenylalanine and tyrosine-free powder for dietary management of children and adults (3 years and older) with inborn errors of tyrosine metabolism including tyrosinemia type II.

Analysis	per 100 mL	per 100 g				
Energy (kcal/kJ)	85/360	410/1720				
Protein equivalent (g)	4.6	22				
% of total energy	22	22				
Source: Amino acid premix						
Carbohydrate (g)	12.5	60				
% of total energy	59	59				
Source: Corn syrup solids, sugar, modified corn starch						
Fat (g)	1.77	8.5				
% of total energy	19	19				
Source: Soy oil						
Linoleic Acid (g)	0.87	4.2				
Linolenic Acid (g)	0.108	0.52				
Minerals						
Calcium (mg) (mMol)	152 3.8	730 18.2				
Phosphorus (mg) (mMol)	152 4.9	730 24				
Magnesium (mg) (mMol)	34 1.4	163 6.7				
Iron (mg)	2.5	12.2				





Analysis	per 100 mL	per 100 g
Zinc (mg)	2.5	12.2
Manganese (mg)	0.27	1.3
Copper (mg)	0.25	1.22
lodine (mg)	0.0131	0.063
Selenium (mg)	0.0058	0.028
Sodium (mg) (mMol)	127 5.5	610 27
Potassium (mg) (mMol)	230 5.9	1100 28
Chloride (mg) (mMol)	177 5.0	850 24
Vitamins		
Vitamin A (IU)	300	1420
Vitamin D (IU)	58	280
Vitamin E (IU)	2	9.8
Vitamin K (mg)	0.0085	0.041
Vitamin C (mg)	10.2	49
Thiamine (mg)	0.25	1.22
Riboflavin (mg)	0.2	0.98
Niacin (mg)	4.6	22
Pantothenic Acid (mg)	1.02	4.9
Vitamin B <sub>6</sub> (mg)	0.2	0.98
Folic Acid (mg)	0.073	0.35
Vitamin B <sub>12</sub> (mg)	0.0005	0.0024
Biotin (mg)	0.0102	0.049
Choline (mg)	20	98
Inositol (mg)	10.2	49
Carnitine (mg)	10.2	49
Taurine (mg)	10.2	49

Other Characteristics	
Potential Renal Solute Load (m0sm/100 mL) <sup>1</sup>	48
Osmolality (mOsm/kg H <sub>2</sub> O)*	830
Osmolarity (mOsm/L)*	710
Water (g/100 mL)	86

Use only as directed by a medical professional.

Powder medical foods are not sterile.

**WARNING:** This protein is incomplete since it does not contain the essential amino acids tyrosine and phenylalanine.

- \* 100 mL values and Other Characteristics information is based on this product mixed with water to a caloric density of 0.85 kcal/mL.
- 1. Fomon SJ, Ziegler EE. Renal solute load and potential renal solute load in infancy. *J Pediatr*. 1999;134:11-14. Nutrient values are subject to change. Consult actual product labels for most current information.

### **INGREDIENTS**

Corn Syrup Solids, Amino Acids (L-Glutamine, L-Leucine, Potassium Aspartate, L-Lysine Hydrochloride, L-Proline, L-Valine, L-Isoleucine, L-Alanine, Larginine, L-Threonine, L-Serine, Glycine, L-Histidine, L-Methionine, L-Tryptophan, L-Cystine), Sugar, Soy Oil, Modified Corn Starch, Calcium Phosphate, Sodium Citrate, Magnesium Phosphate, Potassium Chloride, And Less Than 1%: Sodium Phosphate, Potassium Citrate, Choline Chloride, Ascorbic Acid, Taurine, Inositol, Ferrous Sulfate, L-Carnitine, Zinc Sulfate, Niacinamide, Vitamin E Acetate, Vitamin A Palmitate, Calcium Pantothenate, Cupric Sulfate, Manganese Sulfate, Vitamin B<sub>12</sub>, Thiamin Hydrochloride, Vitamin B<sub>6</sub> Hydrochloride, Riboflavin, Folic Acid, Vitamin D<sub>3</sub>, Chromic Chloride, Sodium Molybdate, Sodium Iodide, Sodium Selenite, Biotin, Vitamin K<sub>1</sub>, Ethyl Vanillin.

## **PRODUCT FORMS**

Product	Format	Unit Size	Approximate Product Yield at Normal Dilution	Shelf Life	Item Number	Availability
TYROS 2	Powder	454 g	Yield Not Applicable to Metabolics	24 Months	1281396	Hospital & Retail Pharmacy (on prescription)

### PREPARATION OF FEEDINGS

**Mead Johnson TYROS 2** is to be used only in the dietary management of children and adults with documented inborn errors of tyrosine metabolism including tyrosinemia type II while under direct and continuing medical supervision.

**WARNING:** This product is nutritionally incomplete. Continuing medical supervision and frequent blood tests are required. **TYROS 2** must be supplemented using other food sources of protein and fluid to provide enough tyrosine and phenylalanine to support dietary requirements of children and adults.

#### INSTRUCTIONS FOR PREPARATION AND USE

Your medical professional will provide the correct amount of powder to mix with water for consumption.\*

It is important to follow the directions below.

Measure the correct amount of water into a suitable container for mixing. Then add the required amount of **TYROS 2. Mix well until blended.** Consume the prepared beverage immediately or cover and refrigerate. Use within 48 hours of preparation. Mix before drinking.

\* If instructed to use the scoop in the can, each unpacked, level scoop delivers approximately 14.5 g of powder. Store the DRY scoop in this can.

**Powder Storage:** Store cans at room temperature. After opening can, keep tightly covered, store in a dry area and use contents within 1 month. Do not freeze powder and avoid excessive heat. **Use by date on bottom of can.** 



## **BCAD 1**

An isoleucine-, leucine-, and valine-free, iron fortified infant formula with DHA and ARA for the dietary management of infants and young children with documented Maple Syrup Urine Disease (MSUD).

(U) D (M) IFANCA

## INDICATION

An isoleucine-, leucine-, and valine-free, iron fortified infant formula with DHA and ARA for the dietary management of infants and young children with documented Maple Syrup Urine Disease (MSUD).

Analysis	per 100 mL	per 100 g			
Energy (kcal/kJ)	68/280	500/2100			
Protein equivalent (g)	2.2	16.2			
% of total energy	13	13			
Source: Amino acid premix					
Carbohydrate (g)	6.9	51			
% of total energy	40	40			
Source: Corn syrup solids, modified corn	starch, sugar				
Fat (g)	3.5	26			
% of total energy	47	47			
Source: Palm olein oil (44%), soy oil (19.5%), coconut oil (19.5%), high oleic sunflower oil (14.5%), ARA & DHA single cell oil blend (2.5%)					
Linoleic Acid (g)	0.54	4			
Linolenic Acid (g)	0.046	0.34			
ARA (mg)	24	175			
DHA (mg)	11.9	88			
Minerals					
Calcium (mg)	89	660			
(mMol)	2.2	16.5			
Phosphorus (mg)	59	440			
(mMol)	1.92	14.2			





Analysis	per 100 mL	per 100 g
Magnesium (mg)	8.9	66
(mMol)	0.37	2.7
Iron (mg)	1.3	9.6
Zinc (mg)	1.16	8.6
Manganese (mg)	0.051	0.38
Copper (mg)	0.116	0.86
lodine (mg)	0.0103	0.076
Selenium (mg)	0.0019	0.0141
Sodium (mg) (mMol)	32 1.41	240 10.4
Potassium (mg) (mMol)	81 2.1	600 15.3
Chloride (mg) (mMol)	68 1.9	500 14.1
Vitamins		
Vitamin A (IU)	210	1520
Vitamin D (IU)	51	380
Vitamin E (IU)	1.35	10
Vitamin K (mg)	0.0059	0.044
Vitamin C (mg)	8.1	60
Thiamine (mg)	0.135	1
Riboflavin (mg)	0.135	1
Niacin (mg)	1.35	10
Pantothenic Acid (mg)	0.51	3.8
Vitamin B <sub>6</sub> (mg)	0.135	1
Folic Acid (mg)	0.0135	0.1
Vitamin B <sub>12</sub> (mg)	0.00027	0.002
Biotin (mg)	0.0051	0.038
Choline (mg)	16.7	124
Inositol (mg)	11.6	86
Carnitine (mg)	6.8	50
Taurine (mg)	4.1	30

Other Characteristics				
Potential Renal Solute Load (m0sm/100 mL) <sup>1</sup>	20			
Osmolality (m0sm/kg H <sub>2</sub> 0)*	330			
Osmolarity (m0sm/L)*	300			
Water (g/100 mL)	90			

All metabolic products are for use under the direct and continuing supervision of a doctor or metabolic dietitian.

This product is nutritionally incomplete. Care must be taken to provide enough isoleucine, leucine and valine to support growth, using other foods with these amino acids as required. Nutritional powders are not sterile.

- \* 100 mL values and Other Characteristics information is based on this product mixed with water to a caloric density of 0.68 kcal/mL.
- 1. Fomon SJ, Ziegler EE. Renal solute load and potential renal solute load in infancy. *J Pediatr*. 1999;134:11-14. Nutrient values are subject to change. Consult actual product labels for most current information.

### **INGREDIENTS**

Corn Syrup Solids, Vegetable Oil (Palm Olein Oil, Coconut Oil, Soy Oil, High Oleic Sunflower Oil), Amino Acids (L-Glutamine, L-Lysine Hydrochloride, Potassium Aspartate, L-Proline, L-Alanine, L-Arginine, L-Phenylalanine, L-Tyrosine, L-Serine, L-Threonine, Glycine, L-Histidine, L-Methionine, L-Tryptophan, L-Cystine), Modified Corn Starch, Sugar, Mortierella Alpina Oil<sup>†</sup>, Crypthecodinium Cohnii Oil<sup>‡</sup>, Ethyl Vanillin, L-Carnitine, Taurine, Ascorbyl Palmitate, Mixed Tocopherols (May Contain Potassium Hydroxide), Minerals (Calcium Phosphate, Choline Chloride, Cupric Sulfate, Ferrous Sulfate, Inositol, Magnesium Oxide, Manganese Sulfate, Potassium Chloride, Potassium Citrate, Sodium Iodide, Sodium Selenite And Zinc Sulfate), Vitamins (Ascorbic Acid, Biotin, Calcium Pantothenate, Folic Acid, Niacinamide, Riboflavin, Thiamine Hydrochloride, Vitamin A Palmitate, Vitamin B<sub>12</sub>, Vitamin B<sub>6</sub> Hydrochloride, Vitamin E Acetate And Vitamin K<sub>1</sub>).

- † A source of Arachidonic Acid (ARA).
- ‡ A source of Docosahexaenoic Acid (DHA).

### PRODUCT FORMS

Product	Format	Unit Size	Approximate Product Yield at Normal Dilution	Shelf Life	Item Number	Availability
BCAD 1	Powder	454 g	Yield Not Applicable to Metabolics	18 Months	2001952	Hospital & Retail Pharmacy (on prescription)

## PREPARATION OF FEEDINGS

**Mead Johnson BCAD 1** is to be used only in the dietary management of infants and young children with documented Maple Syrup Urine Disease while under direct and continuing medical supervision.

WARNING: This product is nutritionally incomplete. Care must be taken to provide enough isoleucine, leucine and valine to support growth, using other foods with these amino acids as required. Medical professionals must carefully and constantly supervise use of BCAD 1 with other foods and liquids and adjust the diet based on frequent blood tests.

#### INSTRUCTIONS FOR PREPARATION AND USE

Your baby's health depends on carefully following the instructions below.

Proper hygiene, preparation, dilution, use and storage are important when preparing formula. Powdered formulas are not sterile and should <u>not</u> be fed to premature infants or those persons who might have immune problems unless directed and supervised by your baby's doctor.



1. Wash hands thoroughly. Boil clean bottles, nipples, caps and utensils in water (2 minutes at a rolling boil).



2. Boil fresh water at a rolling boil for 2 minutes. Cool to room temperature prior to mixing.



3. Pour desired amount of the cooled water into the bottle. Add powder. Cap bottle and shake vigorously.

Your baby's doctor will provide instructions for the correct amount of cooled, boiled water and powder. Each scoop (unpacked and level) delivers approximately 4.5 g of powder. Store the DRY scoop in this can.

Failure to follow these instructions could result in severe harm. Once prepared, formula can spoil quickly. Either feed immediately or cover and store in refrigerator at 2-4°C (35-40°F) for no longer than 24 hours. Do not use prepared formula if it is unrefrigerated for more than a total of 2 hours. Do not freeze prepared formula. After feeding begins, do not refrigerate feeding bottle. You must use within 1 hour or discard.

**WARNING:** Do not use a microwave oven to warm formula. Serious burns may result.

**STORAGE:** Store cans at room temperature. After opening can, keep tightly covered, store in dry area and use contents within one month. Do not freeze powder and avoid excessive heat. **Use by date on bottom of can.** 



## BCAD 2

Isoleucine-, leucine- and valine-free powder for dietary management of children and adults (3 years and older) with Maple Syrup Urine Disease (MSUD) or other inborn errors of branched chain amino acid metabolism.

OD (M) IFANCA

## **INDICATION**

Isoleucine-, leucine- and valine-free powder for dietary management of children and adults (3 years and older) with Maple Syrup Urine Disease (MSUD) or other inborn errors of branched chain amino acid metabolism.

Analysis	per 100 mL	per 100 g			
Energy (kcal/kJ)	85/360	410/1720			
Protein equivalent (g)	5	24			
% of total energy	24	24			
Source: Amino acid premix					
Carbohydrate (g)	11.9	57			
% of total energy	57	57			
Source: Corn syrup solids, sugar, modifie	ed corn starch				
Fat (g)	1.77	8.5			
% of total energy	19	19			
Source: Soy oil					
Linoleic Acid (g)	0.87	4.2			
Linolenic Acid (g)	0.108	0.52			
Minerals					
Calcium (mg) (mMol)	152 3.8	730 18.2			
Phosphorus (mg) (mMol)	152 4.9	730 24			
Magnesium (mg) (mMol)	34 1.4	163 6.7			
Iron (mg)	2.5	12.2			





Analysis	per 100 mL	per 100 g
Zinc (mg)	2.5	12.2
Manganese (mg)	0.27	1.3
Copper (mg)	0.25	1.22
lodine (mg)	0.0131	0.063
Selenium (mg)	0.0058	0.028
Sodium (mg) (mMol)	127 5.5	610 27
Potassium (mg) (mMol)	250 6.5	1220 31
Chloride (mg) (mMol)	210 6	1020 29
Vitamins		
Vitamin A (IU)	360	1730
Vitamin D (IU)	73	350
Vitamin E (IU)	2.4	11.4
Vitamin K (mg)	0.0098	0.047
Vitamin C (mg)	11.9	57
Thiamine (mg)	0.3	1.42
Riboflavin (mg)	0.24	1.14
Niacin (mg)	5.4	26
Pantothenic Acid (mg)	1.19	5.7
Vitamin B <sub>6</sub> (mg)	0.24	1.14
Folic Acid (mg)	0.085	0.41
Vitamin B <sub>12</sub> (mg)	0.00058	0.0028
Biotin (mg)	0.0119	0.057
Choline (mg)	20	98
Inositol (mg)	11.9	57
Carnitine (mg)	10.2	49
Taurine (mg)	11.9	57

Other Characteristics	
Potential Renal Solute Load (m0sm/100 mL) <sup>1</sup>	51
Osmolality (mOsm/kg H <sub>2</sub> O)*	900
Osmolarity (mOsm/L)*	780
Water (g/100 mL)	86

Use only as directed by a medical professional.

Powder medical foods are not sterile.

WARNING: This protein is incomplete since it does not contain the essential amino acids isoleucine, leucine and valine.

- \* 100 mL values and Other Characteristics information is based on this product mixed with water to a caloric density of 0.85 kcal/mL.
- Fomon SJ, Ziegler EE. Renal solute load and potential renal solute load in infancy. J Pediatr. 1999;134:11-14.
   Nutrient values are subject to change. Consult actual product labels for most current information.

#### **INGREDIENTS**

Corn Syrup Solids, Amino Acids (L-Glutamine, L-Lysine Hydrochloride, Potassium Aspartate, L-Proline, L-Alanine, L-Arginine, L-Phenylalanine, L-Tyrosine, L-Threonine, L-Serine, Glycine, L-Histidine, L-Methionine, L-Tryptophan, L-Cystine), Sugar, Soy Oil, Modified Corn Starch, Calcium Phosphate, And Less Than 1%: Sodium Citrate, Magnesium Phosphate, Potassium Chloride, Sodium Phosphate, Potassium Citrate, Choline Chloride, Ascorbic Acid, Taurine, Inositol, Ferrous Sulfate, L-Carnitine, Zinc Sulfate, Niacinamide, Vitamin E Acetate, Vitamin A Palmitate, Calcium Pantothenate, Cupric Sulfate, Manganese Sulfate, Vitamin B<sub>12</sub>, Thiamin Hydrochloride, Vitamin B<sub>6</sub> Hydrochloride, Riboflavin, Folic Acid, Vitamin D<sub>3</sub>, Chromic Chloride, Sodium Molybdate, Sodium Iodide, Biotin, Sodium Selenite, Vitamin K<sub>1</sub>, Ethyl Vanillin.

#### **PRODUCT FORMS**

Product	Format	Unit Size	Approximate Product Yield at Normal Dilution	Shelf Life	Item Number	Availability
BCAD 2	Powder	454 g	Yield Not Applicable to Metabolics	24 Months	1281386	Hospital & Retail Pharmacy (on prescription)

#### PREPARATION OF FEEDINGS

**Mead Johnson BCAD 2** is to be used only in the dietary management of children and adults with documented maple syrup urine disease or other inborn errors of branched chain amino acid metabolism while under direct and continuing medical supervision.

**WARNING:** This product is nutritionally incomplete. Continuing medical supervision and frequent blood tests are required. **BCAD 2** must be supplemented using other food sources of protein and fluid to provide enough isoleucine, leucine and valine to support dietary requirements of children and adults.

#### INSTRUCTIONS FOR PREPARATION AND USE

Your medical professional will provide the correct amount of powder to mix with water for consumption.

It is important to follow the directions below.

**Powder Storage:** Store cans at room temperature. After opening can, keep tightly covered, store in a dry area and use contents within 1 month. Do not freeze powder and avoid excessive heat. **Use by date on bottom of can.** 

**WARNING:** Not for parenteral (I.V.) use.



## PFD<sub>1</sub>

Protein-and amino acid-free infant formula\* with DHA and ARA for infants and young children with documented inborn errors of amino acid metabolism.

\* This product does contain taurine, a non-protein building amino acid.

(U) D (M) IFANCA

#### INDICATION

Protein-and amino acid-free infant formula\* with DHA and ARA for infants and young children with documented inborn errors of amino acid metabolism.

### **NUTRIENTS**

Analysis	per 100 mL	per 100 g			
Energy (kcal/kJ)	69/290	530/2220			
Protein equivalent (g)	0	0			
% of total energy	0	0			
Source: None added					
Carbohydrate (g)	7.8	60			
% of total energy	46	46			
Source: Corn syrup solids, modified corn	starch, sugar				
Fat (g)	4.2	32			
% of total energy	54	54			
Source: Palm olein oil (44%), soy oil (19.5%), coconut oil (19.5%), high oleic sunflower oil (14.5%), ARA & DHA single cell oil blend (2.5%)					
Linoleic Acid (g)	0.64	4.9			
Linolenic Acid (g)	0.052	0.4			
ARA (mg)	25	190			
DHA (mg)	12.4	95			
Minerals					
Calcium (mg) (mMol)	103 2.6	790 20			
Phosphorus (mg) (mMol)	69 2.2	530 17.1			





<sup>\*</sup> This product does contain taurine, a non-protein building amino acid.

Analysis	per 100 mL	per 100 g
Magnesium (mg)	10.3	79
(mMol)	0.42	3.3
Iron (mg)	1.37	10.5
Zinc (mg)	1.37	10.5
Manganese (mg)	0.061	0.47
Copper (mg)	0.137	1.05
lodine (mg)	0.0117	0.09
Selenium (mg)	0.0021	0.0158
Sodium (mg) (mMol)	36 1.58	280 12.2
Potassium (mg) (mMol)	90 2.3	690 17.6
Chloride (mg) (mMol)	69 1.94	530 14.9
Vitamins		
Vitamin A (IU)	240	1840
Vitamin D (IU)	59	450
Vitamin E (IU)	1.72	13.2
Vitamin K (mg)	0.0069	0.053
Vitamin C (mg)	9.6	74
Thiamine (mg)	0.172	1.32
Riboflavin (mg)	0.157	1.21
Niacin (mg)	1.57	12.1
Pantothenic Acid (mg)	0.6	4.6
Vitamin B <sub>6</sub> (mg)	0.157	1.21
Folic Acid (mg)	0.0172	1.32
Vitamin B <sub>12</sub> (mg)	0.00034	0.0026
Biotin (mg)	0.0061	0.047
Choline (mg)	17.6	135
Inositol (mg)	14.2	109
Carnitine (mg)	8.2	63
Taurine (mg)	4.8	37

Other Characteristics	
Potential Renal Solute Load (m0sm/100 mL) <sup>1</sup>	8
Osmolality (mOsm/kg H <sub>2</sub> O)*	171
Osmolarity (mOsm/L)*	154
Water (g/100 mL)	90

All metabolic products are for use under the direct and continuing supervision of a doctor or metabolic dietitian

This product is nutritionally incomplete. Care must be taken to provide enough protein or amino acids to support growth, using other foods with protein as required. Nutritional powders are not sterile.

- \* 100 mL values and Other Characteristics information is based on this product mixed with water to a caloric density of 0.69 kcal/mL.
- 1. Fomon SJ, Ziegler EE. Renal solute load and potential renal solute load in infancy. *J Pediatr*. 1999;134:11-14. Nutrient values are subject to change. Consult actual product labels for most current information.

#### INGREDIENTS

Corn Syrup Solids, Vegetable Oil (Palm Olein Oil, Coconut Oil, Soy Oil, High Oleic Sunflower Oil), Modified Corn Starch, Sugar, Mortierella Alpina Oil<sup>†</sup>, Crypthecodinium Cohnii Oil<sup>‡</sup>, L-Carnitine, Taurine, Ethyl Vanillin, Ascorbyl Palmitate, Mixed Tocopherols (May Contain Potassium Hydroxide), Minerals (Calcium Phosphate, Choline Chloride, Cupric Sulfate, Ferrous Sulfate, Inositol, Magnesium Oxide, Manganese Sulfate, Potassium Chloride, Potassium Citrate, Sodium Citrate, Sodium Iodide, Sodium Selenite And Zinc Sulfate), Vitamins (Ascorbic Acid, Biotin, Calcium Pantothenate, Folic Acid, Niacinamide, Riboflavin, Thiamine Hydrochloride, Vitamin A Palmitate, Vitamin B<sub>12</sub>, Vitamin B<sub>6</sub> Hydrochloride, Vitamin D<sub>3</sub>, Vitamin E Acetate And Vitamin K<sub>1</sub>).

- † A source of Arachidonic Acid (ARA).
- ‡ A source of Docosahexaenoic Acid (DHA).

#### PRODUCT FORMS

Product	Format	Unit Size	Approximate Product Yield at Normal Dilution	Shelf Life	Item Number	Availability
PFD 1	Powder	400 g	Yield Not Applicable to Metabolics	18 Months	3144090	Hospital & Retail Pharmacy (on prescription)

#### PREPARATION OF FEEDINGS

**Mead Johnson PFD 1** is to be used only in the dietary management of infants and young children with documented inborn errors of amino acid metabolism under direct and continuing medical supervision.

WARNING: This product is nutritionally incomplete. Care must be taken to provide enough protein or amino acids to support growth, using other foods with this product as required. Medical professionals must carefully and constantly supervise use of PFD 1 with other foods and liquids and adjust the diet based on frequent blood tests.

#### INSTRUCTIONS FOR PREPARATION AND USE

Your baby's health depends on carefully following the instructions below. Proper hygiene, preparation, dilution, use and storage are important when preparing formula. Powdered formulas are not sterile and should <u>not</u> be fed to premature infants or those persons who might have immune problems unless directed and supervised by your baby's doctor.



1. Wash hands thoroughly with soap and water before preparing formula. Boil clean bottles, nipples, caps and utensils in water (2 minutes at a rolling boil).



2. Boil fresh water at a rolling boil for 2 minutes. Cool to room temperature prior to mixing.



3. Pour desired amount of the cooled water into the bottle. Add powder. Cap bottle and SHAKE WELL.

Your baby's doctor will provide instructions for the correct amount of cooled, boiled water and powder. If instructed to use the scoop in the can, each scoop (unpacked and level) delivers approximately 4.5 g of powder. Store the DRY scoop in this can.

Failure to follow these instructions could result in severe harm. Once prepared, formula can spoil quickly. Either feed immediately or cover and store in refrigerator at 2-4°C (35-40°F) for no longer than 24 hours. Do not use prepared formula if it is unrefrigerated for more than a total of 2 hours. Do not freeze prepared formula. After feeding begins, use product within 1 hour or discard.

**WARNING:** Do not use a microwave oven to warm formula. Serious burns may result.

**STORAGE:** Store cans at room temperature. After opening can, keep tightly covered, store in dry area and use contents within one month. Do not freeze powder and avoid excessive heat. **Use by date on bottom of can.** 

**WARNING:** Not for parenteral (I.V.) use.



## PFD<sub>2</sub>

Protein- and amino acid-free powder\* for dietary management of children and adults (3 years and older) with documented amino acid metabolic disorders.

\* This product does contain taurine, a non-protein building amino acid.

OD (M) IFANCA

#### **INDICATION**

Protein-and amino acid-free powder\*for dietary management of children and adults (3 years and older) with documented amino acid metabolic disorders.

### **NUTRIENTS**

Analysis	per 100 mL	per 100 g
Energy (kcal/kJ)	85/360	400/1670
Protein equivalent (g)	0	0
% of total energy	0	0
Source: None added		
Carbohydrate (g)	18.7	88
% of total energy	89	89
Source: Corn syrup solids, sugar, modifie	ed corn starch	
Fat (g)	1.02	4.8
% of total energy	11	11
Source: Soy oil		
Linoleic Acid (g)	0.49	2.3
Linolenic Acid (g)	0.06	0.28
Minerals		
Calcium (mg) (mMol)	85 2.1	400 10
Phosphorus (mg) (mMol)	85 2.7	400 12.9
Magnesium (mg) (mMol)	19.4 0.8	91 3.7





<sup>\*</sup> This product does contain taurine, a non-protein building amino acid.

Analysis	per 100 mL	per 100 g
Iron (mg)	1.34	6.3
Zinc (mg)	1.34	6.3
Manganese (mg)	0.147	0.69
Copper (mg)	0.134	0.63
lodine (mg)	0.0075	0.035
Selenium (mg)	0.003	0.0142
Sodium (mg) (mMol)	77 3.3	360 15.7
Potassium (mg) (mMol)	72 1.85	340 8.7
Chloride (mg) (mMol)	68 1.92	320 9
Vitamins		
Vitamin A (IU)	168	790
Vitamin D (IU)	33	154
Vitamin E (IU)	1.09	5.1
Vitamin K (mg)	0.0047	0.022
Vitamin C (mg)	5.5	26
Thiamine (mg)	0.134	0.63
Riboflavin (mg)	0.109	0.51
Niacin (mg)	2.4	11.5
Pantothenic Acid (mg)	0.55	2.6
Vitamin B <sub>6</sub> (mg)	0.109	0.51
Folic Acid (mg)	0.038	0.178
Vitamin B <sub>12</sub> (mg)	0.00025	0.00119
Biotin (mg)	0.0055	0.026
Choline (mg)	10.9	51
Inositol (mg)	5.5	26
Carnitine (mg)	5.5	26
Taurine (mg)	5.5	26

Other Characteristics	
Potential Renal Solute Load (m0sm/100 mL) <sup>1</sup>	9.8
Osmolality (mOsm/kg H <sub>2</sub> O)*	450
Osmolarity (mOsm/L)*	380
Water (g/100 mL)	86

Use only as directed by a medical professional.

Powder medical foods are not sterile.

WARNING: This product is incomplete since it does not contain any protein or amino acids.

- \*100 mL values and Other Characteristics information is based on this product mixed with water to a caloric density of 0.85 kcal/mL
- Fomon SJ, Ziegler EE. Renal solute load and potential renal solute load in infancy. J Pediatr. 1999;134:11-14.
   Nutrient values are subject to change. Consult actual product labels for most current information.

#### **INGREDIENTS**

Corn Syrup Solids, Sugar, Soy Oil, Modified Corn Starch, Calcium Phosphate, And Less Than 1%: Sodium Citrate, Magnesium Phosphate, Potassium Chloride, Sodium Phosphate, Potassium Citrate, Choline Chloride, Ascorbic Acid, Taurine, Inositol, Ferrous Sulfate, L-Carnitine, Zinc Sulfate, Niacinamide, Vitamin E Acetate, Calcium Pantothenate, Maltodextrin, Cupric Sulfate, Manganese Sulfate, Vitamin A Palmitate, Thiamin Hydrochloride, Vitamin B<sub>6</sub> Hydrochloride, Riboflavin, Folic Acid, Vitamin D<sub>3</sub>, Chromic Chloride, Sodium Molybdate, Sodium Iodide, Sodium Selenite, Biotin, Vitamin K<sub>1</sub>, Vitamin B<sub>12</sub>, Ethyl Vanillin.

#### PRODUCT FORMS

Product	Format	Unit Size	Approximate Product Yield at Normal Dilution	Shelf Life	Item Number	Availability
PFD 2	Powder	454 g	Yield Not Applicable to Metabolics	24 Months	1281389	Hospital & Retail Pharmacy (on prescription)

#### PREPARATION OF FEEDINGS

**Mead Johnson PFD 2** is to be used only in the dietary management of children and adults with documented amino acid metabolic disorders while under direct and continuing medical supervision.

WARNING: This product is nutritionally incomplete. Continuing medical supervision and frequent blood tests are required. PFD 2 must be supplemented using other food sources of protein and fluid to support dietary requirements of children and adults.

#### INSTRUCTIONS FOR PREPARATION AND USE

Your medical professional will provide the correct amount of powder to mix with water for consumption.\*

It is important to follow the directions below.

Measure the correct amount of water into a suitable container for mixing. Then add the required amount of **PFD 2. Mix well until blended.** Consume the prepared beverage immediately or cover and refrigerate. Use within 48 hours of preparation. Mix before drinking.

\* If instructed to use the scoop in the can, each unpacked, level scoop delivers **approximately 14.5 g** of powder. Store the **DRY** scoop in this can.

**Powder Storage:** Store cans at room temperature. After opening can, keep tightly covered, store in a dry area and use contents within 1 month. Do not freeze powder and avoid excessive heat. **Use by date on bottom of can.** 

WARNING: Not for parenteral (I.V.) use.



Date published 11/2024 – Please reference Mead Johnson Nutrition's online Product Handbook for the most up to date product details (www.enfamil.ca/medical).



# Enfamil® D-Vi-Sol® (Liquid)

## Vitamin D Supplement

Liquid vitamin D supplement for all full-term breastfed infants.

ФD

NPN 00762881

#### **INDICATION**

Liquid vitamin D supplement for all full-term breastfed infants. Enfamil® D-Vi-Sol® provides breastfed infants with vitamin D which helps in the normal development and maintenance of bones and teeth. Helps to prevent vitamin D deficiency. A joint statement of Health Canada, Canadian Paediatric Society, Dietitians of Canada, and Breastfeeding Committee for Canada states that a daily vitamin D supplement of 10 mcg (400 IU) is recommended for exclusively and partially breastfed infants, from birth to one year of age.

#### RATIONALE AND SPECIAL CHARACTERISTICS

Enfamil® D-Vi-Sol® liquid provides 400 IU per mL of vitamin D to meet the recommended daily intake of vitamin D. It is sugar-free, gluten-free, and lactose-free.

#### DOSAGE AND ADMINISTRATION

Daily Dosage for all full-term breastfed infants: 1 mL (fill provided syringe to 1.0 mL line). Dispense slowly into child's mouth, toward inner cheek.

If preferred, the liquid may be slowly mixed with formula, juice, cereal, or other food and fed within one hour.

Medicinal Ingredients (per 1 mL)		
Vitamin D <sub>3</sub>	(cholecalciferol)	400 IU (10 mcg)

#### NON-MEDICINAL INGREDIENTS

Glycerin, polysorbate 80, caramel, citric acid anhydrous, sodium citrate dihydrate, sodium hydroxide, artificial flavour and purified water. Store at room temperature 15-30°C (59-86°F).

#### **PRECAUTIONS**

Keep this and all medications out of the reach of children. Safety sealed bottle. Do not use if seal is broken.



#### **REFERENCE**

 Health Canada, Canadian Pediatric Society, Dietitians of Canada and Breastfeeding Committee for Canada. Nutrition for Healthy Term Infants: Recommendations from Birth to Six Months, 2012.

## **PRODUCT FORMS**

Product	Format	Unit Size	Approximate Product Yield at Normal Dilution	Shelf Life	Item Number	Availability
Enfamil® D-Vi-Sol®	Liquid	50 mL	Yield Not Applicable to Vitamin	18 Months	1015277	Hospital & Retail



# Enfamil® Tri-Vi-Sol® (Liquid)

# Multi-Vitamin Supplement of Vitamins A, D, and C

Liquid vitamin A, D, and C supplement for infants and children under 3 years of age.

For the maintenance of good health for growing infants and toddlers.

**O**D

NPN 00762903

#### INDICATION

Liquid vitamin A, D, and C supplement for infants and children under 3 years of age. For the maintenance of good health for growing infants and toddlers.

### RATIONALE AND SPECIAL CHARACTERISTICS

Enfamil® Tri-Vi-Sol® vitamin liquid provides 3 frequently prescribed vitamins in amounts suitable for daily administration to infants and toddlers. Enfamil® Tri-Vi-Sol® vitamin liquid is lactose-free, gluten-free, and free of artificial sweeteners.

#### DOSAGE AND ADMINISTRATION

Daily Dosage: 1 mL (fill provided syringe to 1.0 mL line). Dispense slowly into child's mouth, toward inner cheek. If preferred, the liquid may be slowly mixed with formula, juice, cereal, or other food and fed within one hour.

Medicinal Ingredients (per 1 mL)	
Vitamin C (ascorbic acid)	30 mg
Vitamin A (vitamin A palmitate)	750 IU (225 mcg RAE)
Vitamin D <sub>3</sub> (cholecalciferol)	400 IU (10 mcg)

#### **NON-MEDICINAL INGREDIENTS**

Glycerin, purified water, polysorbate 80, sodium hydroxide, artificial flavour and caramel colour.

Store at room temperature 15-30°C (59-86°F).

#### **PRECAUTIONS**

Keep this and all medications out of the reach of children. Safety sealed bottle. Do not use if seal is broken.





## **PRODUCT FORMS**

Product	Format	Unit Size	Approximate Product Yield at Normal Dilution	Shelf Life	Item Number	Availability
Enfamil® Tri-Vi-Sol®	Liquid	50 mL	Yield Not Applicable to Vitamin	15 Months	1015245	Hospital & Retail



# Enfamil® Poly-Vi-Sol® (Liquid)

## Multi-Vitamin Supplement

Liquid multivitamin supplementation for infants and children under 3 years of age. For the maintenance of good health for growing infants and toddlers.

**(**) D

NPN 00762946

#### INDICATION

Liquid multivitamin supplementation for infants and children under 3 years of age. For the maintenance of good health for growing infants and toddlers.

#### RATIONALE AND SPECIAL CHARACTERISTICS

Enfamil® Poly-Vi-Sol® multivitamin supplement liquid provides 6 important vitamins and is a supplement for an infant's and toddler's transition to solid foods, during growth spurts, and for the picky eater. It is fruit-flavored, lactose-free, gluten-free, and free of artificial sweeteners.

#### **DOSAGE AND ADMINISTRATION**

Daily Dosage: 1 mL (fill provided syringe to 1.0 mL line). Dispense slowly into child's mouth, toward inner cheek. If preferred, the liquid may be slowly mixed with formula, juice, cereal, or other food and fed within one hour.

Medicinal Ingredients (per 1 mL)	
Vitamin C (ascorbic acid)	30 mg
Niacinamide (niacinamide)	4 mg
Riboflavin (riboflavin 5'-phosphate sodium)	0.6 mg
Thiamine (thiamine hydrochloride)	0.5 mg
Vitamin A (vitamin A palmitate)	750 IU (225 mcg RAE)
Vitamin D <sub>3</sub> (cholecalciferol)	400 IU (10 mcg)

#### NON-MEDICINAL INGREDIENTS

Glycerin, purified water, polysorbate 80, caramel colour and mixed fruit flavour. Store at room temperature 15-30°C (59-86°F).

#### **PRECAUTIONS**

Keep this and all medications out of the reach of children. Safety sealed bottle. Do not use if seal is broken.





## **PRODUCT FORMS**

Product	Format	Unit Size	Approximate Product Yield at Normal Dilution	Shelf Life	Item Number	Availability
Enfamil® Poly-Vi-Sol®	Liquid	50 mL	Yield Not Applicable to Vitamin	15 Months	1015280	Hospital & Retail



# Enfamil® Fer-In-Sol® (Liquid)

Ferrous Sulfate Oral Solution U.S.P.

Helps to prevent iron deficiency.

(U) D

NPN 00762954

### **INDICATION**

Helps to prevent iron deficiency.

#### **NUTRIENTS**

Medicinal Ingredients (per 1 mL)						
Iron (ferrous sulfate heptahydrate)	75 mg					
Equivalent to: Elemental Iron	15 mg					

#### **NON-MEDICINAL INGREDIENTS**

Natural flavouring, citric acid anhydrous, ethyl alcohol, sodium bisulfate, sorbital solution, sugar and purified water.

#### **DOSAGE AND ADMINISTRATION**

Use as directed by your physician. For oral use.

**Dosage:** For children 0-2 years. Take with food. Take a few hours before or after taking other medications. Dispense slowly into child's mouth, toward inner cheek.

Supplementation dose: 0.5 to 1 mL daily or as directed by your physician.

Therapeutic dose: 2.5 mL daily or as directed by your physician. Divide into three (3) doses and take three (3) times daily.

#### **Cautions and Warnings:**

At therapeutic dose, some people may experience constipation, diarrhea and/ or vomiting. Temporary discoloration of the teeth due to this iron liquid can be minimized by thorough brushing.

Normal appearance of Enfamil® Fer-in-Sol® liquid is colourless, but the colour may vary from blue-green to gold. Any colour within this range does not affect the potency of the product.

Store at room temperature 15-30°C (59-86°F).





#### SIDE EFFECTS

While taking iron, stools may appear darker in colour. This is normal and no cause for concern. When iron liquid is given to babies, some darkening of teeth may occur. This is not serious and may be removed by rubbing the teeth with baking soda.

#### **PRECAUTIONS**

Use as directed by your physician. For oral use only. Keep this and all medications out of the reach of children. There is enough drug in the package to seriously harm a child. Safety sealed bottle. Do not use if seal is broken.

#### ADVERSE REACTION

Some people may experience constipation, diarrhea and/or vomiting.

#### **ACCIDENTAL OVERDOSAGE OR INTAKE**

In case of accidental overdose, the physician, Poison Control Center, or hospital emergency should be notified immediately.

#### PRODUCT FORMS

Product	Format	Unit Size	Approximate Product Yield at Normal Dilution	Shelf Life	Item Number	Availability
Enfamil® Fer-Vi-Sol®	Liquid	50 mL	Yield Not Applicable to Vitamin	24 Months	1015154	Hospital & Retail



# Enfamil® Fer-In-Sol® (Syrup)

Ferrous Sulfate Oral Solution U.S.P.

Helps to prevent iron deficiency.

**(**)D

NPN 00017884

### INDICATION

Helps to prevent iron deficiency. Enfamil® Fer-in-Sol® syrup is fruit-flavoured, alcohol-free, lactose-free and gluten-free.

### **NUTRIENTS**

Fer-In-Sol syrup contains	Per 5.0 mL
Iron (ferrous sulfate heptahydrate)	150 mg
Equivalent to: Elemental Iron	30 mg

#### **NON-MEDICINAL INGREDIENTS**

Sucrose, sorbitol solution, natural and artificial flavour, citric acid anhydrous, sodium chloride, sodium bisulfite, sulphuric acid and purified water.

Store at room temperature 15-30°C (59-86°F).

#### DOSAGE AND ADMINISTRATION

Take a few hours before or after taking other medication. Mix with water or fruit juice and take with food.

Supplementation dose:						
Administer once daily or as directed by your physician.						
0-6 years	1.25 mL					
6-12 years	2.5 mL					
Adults	5 mL					



Therapeutic dose:						
Divide into three (3) doses and administer three (3) times per day or as directed by your physician.						
0-2 years	2.5 to 5 mL					
2-6 years 5 mL						
6-12 years 5 mL to 12 mL						
Adults	10 to 15 mL					

#### **PRECAUTIONS**

Keep this and all medication out of the reach of children. There is enough drug in the package to seriously harm a child. Safety sealed bottle. Do not use if seal is broken.

#### ADVERSE REACTION

Some people may experience constipation, diarrhea and/or vomiting. Temporary discolouration of the teeth due to this iron syrup can be minimized by thorough brushing.

#### **ACCIDENTAL OVERDOSAGE OR INTAKE**

In case of accidental overdose, the physician, Poison Control Center, or hospital emergency should be notified immediately.

## **PRODUCT FORMS**

Product	Format	Unit Size	Approximate Product Yield at Normal Dilution	Shelf Life	Item Number	Availability
Enfamil® Fer-In-Sol®	Syrup	250 mL	Yield Not Applicable to Vitamin	24 Months	1015278	Hospital & Retail





Date published 11/2024 – Please reference Mead Johnson Nutrition's online Product Handbook for the most up to date product details (www.enfamil.ca/medical).

# Enfamil® Nipples

Enfamil® Nipples are designed to be used with Nursette® bottles. Three different nipples are available, to satisfy the various preferences of infants, parents and healthcare professionals. All are ready to use and disposable. Nipples are packaged 240 per case. Enfamil® Nipples are made of surgical-grade plastic and are latex free.

#### SPECIAL FEATURES OF THE NEW NIPPLES

- Ready-to-use: Individually wrapped and disposable for safety and convenience
- Made with TPE: Not made with latex, BPA, PVC, phthalate or DEHP
- Reinforced: The nipples are now reinforced preventing complete nipple collapse while feeding
- Optimal ring design: New ring design with grooves supports grip
- New teat design: Indentation in the teat offers better strength/bite resistance and prevents blocking
- Needle technology for nipple apertures: Consistent flow rate and lesser standard deviation
- Vented: Allows airflow during feeding



## Enfamil® Standard-Flow Soft Nipple Item # 3196175 - 240 per case

A standard-flow single hole (0.018"), soft nipple designed for routine formula feeding.

Product Composition	Latex	BPA	PVC	Phthalates or
	(Y/N)	(Y/N)	(Y/N)	DEHP
Surgical-grade plastic (TPE)	N	N	N	N



# Enfamil® Slow-Flow Soft Nipple Item # 3196177 - 240 per case

A slow-flow, single hole (0.015"), soft nipple for when a slower flow rate is desired. Generally used by premature infants or those with special feeding needs

Product Composition	Latex	BPA	PVC	Phthalates or
	(Y/N)	(Y/N)	(Y/N)	DEHP
Surgical-grade plastic (TPE)	N	N	N	N



## Enfamil® Extra Slow-Flow Soft Nipple Item # 3196178 - 240 per case

A slower flow, single hole (0.011"), soft nipple for when a slower flow rate than the Enfamil® Slow-Flow Soft Nipple is desired. Generally used by premature infants or those with special feeding needs.

Product Composition	Latex	BPA	PVC	Phthalates or
	(Y/N)	(Y/N)	(Y/N)	DEHP
Surgical-grade plastic (TPE)	N	N	N	N



## NUK® Orthodontic Slow-Flow Nipple Item # 3221215 - 70 per case

NUK® Orthodontic Slow-Flow Nipple is made out of TPE (thermoplastic elastomer) and is latex free. This safe product is designed to simulate the shape of a mother's nipple during breastfeeding. The product also has an Anti-Colic Air System which

prevents the nipple from collapsing and reduces risk of swallowing air\*. The nipples are individually packed, sterilized, ready to use and are recommended for single use only. They are also compatible with all Mead Johnson Nutrition Nursette® and ready-to-feed products.

Product Composition	Latex	BPA	PVC	Phthalates or
	(Y/N)	(Y/N)	(Y/N)	DEHP
Surgical-grade plastic (TPE)	N	N	N	N

<sup>\*</sup> These features are based on the product guide provided by Medic Pro MAPA Clinical Support Document for – NUK® Classic (Standard-Neck) Disposable Teats TPE, Mat.-Nr. 10.520.208 Rev. 418, 03/2019.

## **MAM Pacifier**



#### MAM Comfort™ Pacifier<sup>†</sup>

#### Item # 3122589 - 144 per case

Please contact your Hospital Account Manager for product access and contract inclusion.

- Meets hospital and AAP requirements.
- Designed for infants between 1400 gm 6500 gm, to comfort, promote proper oral

development and support natural suckling behavior.

- 100% silicone, BPA/BPS free, one-piece pacifier is extra soft and specially designed for newborns.
- It is 32% lighter<sup>†</sup> than competitive one-piece pacifiers allowing the pacifier to stay in the baby's mouth more easily.
- The Orthodontic SkinSoft™ silicone nipple ensures the pacifier is easily accepted by babies.
- Shield is perfectly sized for newborns and shaped for minimal skin contact to protect baby's sensitive skin.

Product Composition	Latex	BPA	BPS	PVC	Phthalates or
	(Y/N)	(Y/N)	(Y/N)	(Y/N)	DEHP
One-piece 100% silicone construction	N	N	N	N	N

<sup>†</sup> Scientific reports from the WILD Hi-Precision Institute (Austria) confirm that the MAM Comfort is on average 32% lighter than other one-piece silicone pacifiers. Measured results: 8.2 g - 16.6 g.

# Feeding Bottles and Specialty Nursers

Mead Johnson distributes plastic bottles, Grad-U-Feed® Nursers, and Cleft Lip/ Palate Nursers. All are ready to use and disposable. All are latex free and BPA free.



## Ready-To-Use Plastic Bottles Item # 3170018 - 48 per case

Convenient plastic bottles hold 240 mL (8 fl oz) and come complete with an attached cap. The bottles are made of polypropylene, are latex free and are compatible with all Enfamil® Nipples giving clinicians the flexibility to use the bottle for storage and feeding purposes.

Product Composition	Latex	BPA	PVC	Phthalates or
	(Y/N)	(Y/N)	(Y/N)	DEHP
Polypropylene	N	N	N	N



## Grad-U-Feed® Nurser Item # 2051330 - 100 per case

The Grad-U-Feed® Nurser is sterile and can hold 60 mL [2 fl oz] of feeding. It is used to feed small amounts to infants in neonatal intensive care nurseries. The easy-to-read graduated nurser can collect and store breast milk and infant formula. The ready-to-use sterile nurser can withstand freezing temperatures as low as -24°C and have been drop tested for product integrity. The Grad-U-Feed® Nurser is made of

polypropylene and is latex free. Grad-U-Feed® Caps are sold separately. Nipple units are not provided with Grad-U-Feed® Nursers. Enfamil® Nipples may be used.

Product Composition	Latex	BPA	PVC	Phthalates or
	(Y/N)	(Y/N)	(Y/N)	DEHP
Polypropylene	N	N	N	N





## Grad-U-Feed® Nurser with Cap Item # 3118902 - 100 per case

The Grad-U-Feed® Nurser with Cap is sterile, individually packaged together and ready to use. The easy-to-read graduated nurser can can hold 60 mL (2 fl oz) of feeding. It is used to feed small amounts to infants in neonatal intensive care nurseries. The Grad-U-Feed® Nurser with Cap can collect and store breast milk and infant formula. The ready-to-use sterile nurser with cap can withstand freezing temperatures as low as -24°C and have been drop tested for product

integrity. The Grad-U-Feed® Nurser with Cap is made of polypropylene and is latex free. Nipple units are not provided with Grad-U-Feed® Nursers. Enfamil® Nipples may be used.

Product Composition	Latex	BPA	PVC	Phthalates or
	(Y/N)	(Y/N)	(Y/N)	DEHP
Polypropylene	N	N	N	N



## Grad-U-Feed® Nurser & Cap Dual Pack Item # 2051329 - 100 per case (50x2)

The Grad-U-Feed® Nurser & Cap Dual Pack constitutes of two units of individually packaged, sterile & ready-to-use nursers and caps. This format his highly recommended for usage in hospitals with centralized preparation rooms. The easy-to-read graduated nurser can can hold 60 mL (2 fl oz) of feeding. It is used to feed small amounts to infants in neonatal intensive care nurseries. The Grad-U-Feed® Nurser with cap can collect and store breast milk and infant formula. The ready-to-use sterile nurser with cap can withstand freezing

temperatures as low as  $-24^{\circ}\text{C}$  and have been drop tested for product integrity. The Grad-U-Feed® Nurser with cap is made of polypropylene and is latex free. Nipple units are not provided with Grad-U-Feed® Nursers. Enfamil® Nipples may be used.

Product Composition	Latex	BPA	PVC	Phthalates or
	(Y/N)	(Y/N)	(Y/N)	DEHP
Polypropylene	N	N	N	N



## Grad-U-Feed® Caps Item # 2051331 - 200 per case

These white plastic caps are designed to fit the new Grad-U-Feed® Nursers.

Product Composition	Latex	BPA	PVC	Phthalates or
	(Y/N)	(Y/N)	(Y/N)	DEHP
Polypropylene	N	N	N	N



## Cleft Lip/Palate Nurser Item # 1015466 - 72 per case

The Cleft Lip/Palate Nurser is a squeezable bottle that holds 180 mL (6 fl oz). It comes with an elongated nipple designed for babies with a cleft lip or cleft palate. The nurser is made of low-density polyethylene and is latex free. The nipple is made of surgical-grade plastic and is latex free.

Product Composition	Latex	BPA	PVC	Phthalates or
	(Y/N)	(Y/N)	(Y/N)	DEHP
Polypropylene	N	N	N	N



## Sterile Breast Milk Feeding and Storage (30 mL) Available for orders from Q1 2025 Item # 3298204 - 100/case

Please contact your Hospital Account Manager for product access and contract inclusion.

- Sterile, individually packaged, to reduce chances of cross contamination.
- Designed for breast milk storage and feeding.
- Simplifies the collection, storage and feeding of expressed breast milk, even in low quantities.
- It eliminates the need to transfer precious breast

milk between containers and thus lowering the chance of contamination and risk of infection.

- Compatible with all Enfamil® Nipples.
- Recommended for bed side prep or usage in centralized preparation rooms.
- Recommended for single use only.
- Freezable.

Product Composition	Latex	BPA	PVC	Phthalates or
	(Y/N)	(Y/N)	(Y/N)	DEHP
Polypropylene	N	N	N	N



# Sterile Breast Milk Storage and Feeding Bottle (118 mL)

## Available for orders from Q1 2025

Item # 3296536 - 100/case

Please contact your Hospital Account Manager for product access and contract inclusion.

- Sterile, individually packaged, to reduce chances of cross contamination.
- Designed for breast milk storage and feeding.
- Compatible with all Enfamil® Nipples.
- Recommended for bed side prep or usage in centralized preparation rooms.
- Recommended for single use only.
- Freezable.

Product Composition	Latex	BPA	PVC	Phthalates or
	(Y/N)	(Y/N)	(Y/N)	DEHP
Polypropylene	N	N	N	N



## Sterile Colostrum Collector with Cap (15 mL) Available for orders from Q1 2025

#### Item # 3296505 - 100/case


Please contact your Hospital Account Manager for product access and contract inclusion.

Enfamil<sup>®</sup> Colostrum Collector is a specialized low volume container, designed to collect, transport, store and administer colostrum safely and securely.

The container (15 mL) is sterile, individually packaged, to reduce chances of cross contamination. The container also has graduations in 5 mL increments to measure the amount of colostrum collected and fed to the infant.

- Compatible with all Enfamil® Nipples.
- Recommended for single use only.
- Freezable.

Product Composition	Latex	BPA	PVC	Phthalates or
	(Y/N)	(Y/N)	(Y/N)	DEHP
Polypropylene	N	N	N	N



Date published 11/2024 – Please reference Mead Johnson Nutrition's online Product Handbook for the most up to date product details (www.enfamil.ca/medical).

## **Powder Infant Formula Dilutions**

Note: Powdered products are not commercially sterile and should not be used for immunocompromised patients, unless clinically required, and then under strict medical supervision of preparation and use.

These dilutions can be used with the following Mead Johnson powder formulas:

## Enfamil A+® NeuroPro™ Infant Per 1 Scoop

kcal Desired		1 Scoop Powder* Added to Indicated Volume of Water,	Formula Yield, mL (fl oz)
per fl oz†	per mL	mL (fl oz)	
20	0.68	60 (2.0)	66 (2.2)
22	0.74	54 (1.8)	60 (2.0)
24	0.81	49 (1.6)	55 (1.9)
26	0.88	44 (1.5)	51 (1.7)
27	0.91	42 (1.4)	49 (1.7)
28	0.95	41 (1.4)	47 (1.6)
30	1.01	37 (1.3)	44 (1.5)

<sup>\*</sup> Enfamil A<sup>+®</sup> NeuroPro™ Infant powder should be measured with unpacked, level scoops, using the scoop enclosed in the can.

One-half fl oz of water can be measured using one tablespoon (1/2 fl oz water = 1 Tbsp water).

When accuracy is essential, water should be measured in mL, and powder should be weighed rather than measured.

Powders mix best when added on top of water. Add powder, then close bottle and shake for about 5 seconds. The number of scoops and the amount of water to add can be doubled or tripled to make a larger volume.

One unpacked scoop of Enfamil  $A^{+\otimes}$  NeuroPro $^{\mathbb{M}}$  Infant powder (8.8 g) provides 44 Calories (5.1 kcal/g).

One unpacked scoop of Enfamil  $A^{+\otimes}$  NeuroPro<sup>M</sup> Infant powder displaces about 0.2 fl oz of water (displacement for Enfamil  $A^{+\otimes}$  NeuroPro<sup>M</sup> Infant is 0.77 mL/g).

One U.S. fl oz = 29.57 mL





<sup>†</sup> Fluid ounce measures are provided for convenience when larger variance in prepared dilution is acceptable. Fluid ounce measures in above table are rounded to the nearest 0.1 fluid ounce.

#### Enfamil®

#### Per 1 Scoop

kcal Desired		1 Scoop Powder* Added to Indicated Volume of Water,	Formula Yield, mL (fl oz)
per fl oz†	per mL	mL (fl oz)	
20	0.68	59 (2.0)	65 (2.2)
22	0.74	53 (1.8)	59 (2.0)
24	0.81	48 (1.6)	54 (1.8)
26	0.88	44 (1.5)	50 (1.7)
27	0.91	42 (1.4)	48 (1.6)
28	0.95	41 (1.4)	47 (1.6)
30	1.01	37 (1.3)	44 (1.5)

<sup>\*</sup> Enfamil® should be measured with unpacked, level scoops.

Powders mix best when added on **top** of water. Then close the bottle and shake for about 5 seconds. The number of scoops and the amount of water to add can be doubled or tripled to make a larger volume.

One scoop of Enfamil® provides 44 kcal. One scoop of powder displaces about 6 mL or 0.2 fl oz. One U.S. fl oz = 29.57 mL 1 Tbsp water = 15 mL water

Enfamil A<sup>+®</sup> EnfaCare<sup>®</sup> Per 1 Scoop

kcal Desired		1 Scoop Powder <sup>‡</sup> Added to Indicated Volume of Water,	Formula Yield, mL (fl oz)
per fl oz†	per mL	mL (fl oz)	
22	0.74	57 (1.9)	65 (2.2)
24	0.81	52 (1.7)	59 (2.0)
26	0.88	47 (1.6)	55 (1.8)
27	0.91	45 (1.5)	53 (1.8)
28	0.95	43 (1.5)	51 (1.7)
30	1.01	40 (1.3)	47 (1.6)

 $<sup>^{\</sup>dagger}$  Fluid ounce measures in above table are rounded to the nearest 0.1 fl oz.

<sup>†</sup> Fluid ounce measures in above table are rounded to the nearest 0.1 fl oz.

<sup>‡</sup> Enfamil A<sup>+®</sup> EnfaCare<sup>®</sup> powder should be measured with **unpacked**, level scoops.

Powders mix best when added on **top** of water. Then close the bottle and shake for about 5 seconds. The number of scoops and the amount of water to add can be doubled or tripled to make a larger volume.

One scoop of Enfamil A $^{+}$ ® EnfaCare $^{\odot}$  powder provides 48 kcal. One scoop of powder displaces about 7 mL or 0.3 fl oz water One U.S. fl oz = 29.57 mL 1 Tbsp water = 15 mL water

Enfamil A+® 2 Per 1 Scoop

kcal Desired		1 Scoop Powder <sup>§</sup> Added to Indicated Volume of Water,	Formula Yield, mL (fl oz)
per fl oz†	per mL	mL (fl oz)	1112 (11 02)
20	0.68	60 (2.0)	67 (2.3)
22	0.74	54 (1.8)	60 (2.0)
24	0.81	49 (1.6)	55 (1.9)
26	0.88	44 (1.5)	51 (1.7)
27	0.91	42 (1.4)	49 (1.7)
28	0.95	41 (1.4)	48 (1.6)
30	1.01	37 (1.3)	44 (1.5)

 $<sup>^{\</sup>dagger}$  Fluid ounce measures in above table are rounded to the nearest 0.1 fl oz.

Powders mix best when added on **top** of water. Then close the bottle and shake for about 5 seconds. The number of scoops and the amount of water to add can be doubled or tripled to make a larger volume.

One scoop of Enfamil A $^{+}$ ° 2 powder provides 45 kcal. One scoop of powder displaces about 7 mL or 0.2 fl oz water One U.S. fl oz = 29.57 mL

Enfamil® 2 Per 1 Scoop

kcal Desired		1 Scoop Powder* Added to Indicated Volume of Water,	Formula Yield, mL (fl oz)
per fl oz†	per mL	mL (fl oz)	ML (II 02)
20	0.68	60 (2.0)	66 (2.2)
22	0.74	54 (1.8)	60 (2.0)

<sup>§</sup> Enfamil A+® 2 should be measured with unpacked, level scoops.

<sup>1</sup> Tbsp water = 15 mL water

kcal Desired		1 Scoop Powder* Added to Indicated Volume of Water,	Formula Yield, mL (fl oz)
per fl oz†	per mL	mL (fl oz)	ML (II 02)
24	0.81	49 (1.6)	55 (1.9)
26	0.88	44 (1.5)	51 (1.7)
27	0.91	43 (1.4)	49 (1.7)
28	0.95	41 (1.4)	47 (1.6)
30	1.01	38 (1.3)	44 (1.5)

<sup>\*</sup> Enfamil® 2 should be measured with unpacked, level scoops.

Powders mix best when added on **top** of water. Then close the bottle and shake for about 5 seconds. The number of scoops and the amount of water to add can be doubled or tripled to make a larger volume.

One scoop of Enfamil® 2 powder provides 45 kcal One scoop of powder displaces about 6 mL or 0.2 fl oz water One U.S. fl oz = 29.57 mL 1 Tbsp water = 15 mL water

Enfamil A<sup>+®</sup> NeuroPro™ Gentlease<sup>®</sup> Per 1 Scoop

kcal Desired		1 Scoop Powder‡ Added to Indicated Volume of Water,	Formula Yield, mL (fl oz)
per fl oz†	per mL	mL (fl oz)	IIIL (II 02)
20	0.68	59 (2.0)	66 (2.2)
22	0.74	53 (1.8)	60 (2.0)
24	0.81	48 (1.6)	55 (1.9)
26	0.88	44 (1.5)	51 (1.7)
27	0.91	42 (1.4)	49 (1.7)
28	0.95	40 (1.4)	47 (1.6)
30	1.01	37 (1.3)	44 (1.6)

 $<sup>\</sup>ensuremath{^{\dagger}}$  Fluid ounce measures in above table are rounded to the nearest 0.1 fl oz.

Powders mix best when added on **top** of water. Then close the bottle and shake for about 5 seconds. The number of scoops and the amount of water to add can be doubled or tripled to make a larger volume.

<sup>&</sup>lt;sup>†</sup> Fluid ounce measures in above table are rounded to the nearest 0.1 fl oz.

<sup>‡</sup> Enfamil A<sup>+</sup>® NeuroPro™ Gentlease® should be measured with **unpacked**, level scoops.

One scoop of Enfamil  $A^{+\, \odot}$  NeuroPro $^{\mathbb{M}}$  Gentlease $^{\otimes}$  powder provides 45 kcal. One scoop of powder displaces about 7 mL or 0.2 fl oz water O.e. U.S. fl oz = 29.57 mL

1 Tbsp water = 15 mL water

## Enfamil A<sup>+®</sup> For feeding babies who frequently Spit Up<sup>§</sup> Per 1 Scoop

kcal De	sired	1 Scoop Powder# Added to Indicated Volume of Water,	Formula Yield, mL (fl oz)
per fl oz†	per mL	mL (fl oz)	ML (II 02)
20	0.68	59 (2.0)	66 (2.2)
22	0.74	53 (1.8)	60 (2.0)
24	0.81	48 (1.6)	55 (1.9)

 $<sup>\</sup>dagger$  Fluid ounce measures in above table are rounded to the nearest 0.1 fl oz.

Enfamil A<sup>+®</sup> For feeding babies who frequently Spit Up\* should not be diluted to caloric concentrations higher than 24 kcal/30 mL because of increased viscosity.

Powders mix best when added on **top** of water. Then close the bottle and shake for about 5 seconds. The number of scoops and the amount of water to add can be doubled or tripled to make a larger volume.

One scoop of Enfamil A+ $^{\circ}$  For feeding babies who frequently Spit Up\* powder provides 45 kcal.

One scoop of powder displaces about 7 mL or 0.2 fl oz water.

One U.S. fl oz = 29.57 mL

1 Tbsp water = 15 mL water

## Nutramigen® A+® with LGG® Per 1 Scoop

kcal De	sired	1 Scoop Powder® Added to Indicated Volume of Water,	Formula Yield, mL (fl oz)		
per fl oz†	per mL	mL (fl oz)	ML (It 02)		
20	0.68	60 (2.0)	67 (2.3)		
22	0.74	54 (1.8)	60 (2.0)		
24	0.81	49 (1.6)	55 (1.9)		
26	0.88	44 (1.5)	51 (1.7)		
27	0.91	42 (1.4)	49 (1.7)		

<sup>§</sup> For babies who spit up more than 4 times per day.

<sup>#</sup> Enfamil A<sup>+®</sup> For feeding babies who frequently Spit Up\* should be measured with **unpacked**, level scoops.

kcal Desired		1 Scoop Powder® Added to Indicated Volume of Water,	Formula Yield, mL (fl oz)	
per fl oz†	per mL	mL (fl oz)	ML (It 02)	
28	0.95	41 (1.4)	48 (1.6)	
30	1.01	38 (1.3)	44 (1.5)	

LGG® is a registered trademark of Chr. Hansen A/S.

Powders mix best when added on **top** of water. Then close the bottle and shake for about 5 seconds. The number of scoops and the amount of water to add can be doubled or tripled to make a larger volume.

One scoop of Nutramigen® A+® with LGG® powder provides 45 kcal One scoop of powder displaces about 7 mL or 0.2 fl oz water One U.S. fl oz = 29.57 mL

### Nutramigen® A+®

#### Per 1 Scoop

kcal Desired		1 Scoop Powder* Added to Indicated Volume of Water,	Formula Yield, mL (fl oz)	
per fl oz†	per mL	mL (fl oz)	IIIL (II 02)	
20	0.68	60 (2.0)	67 (2.3)	
22	0.74	54 (1.8)	60 (2.0)	
24	0.81	49 (1.6)	55 (1.9)	
26	0.88	44 (1.5)	51 (1.7)	
27	0.91	42 (1.4)	49 (1.7)	
28	0.95	41 (1.4)	48 (1.6)	
30	1.01	37 (1.3)	44 (1.5)	

<sup>\*</sup> Nutramigen® A+® powders should be measured with **packed**, level scoops.

Powders mix best when added on **top** of water. Then close the bottle and shake for about 5 seconds. The number of scoops and the amount of water to add can be doubled or tripled to make a larger volume.

One scoop of Nutramigen®  $A^{+}$ ® powder provides 45 kcal One scoop of powder displaces about 7 mL or 0.2 fl oz water One U.S. fl oz = 29.57 mL

<sup>†</sup> Fluid ounce measures in above table are rounded to the nearest 0.1 fl oz.

<sup>¶</sup> Nutramigen® A<sup>+®</sup> with LGG® powders should be measured with **packed**, level scoops.

<sup>1</sup> Tbsp water = 15 mL water

<sup>†</sup> Fluid ounce measures in above table are rounded to the nearest 0.1 fl oz.

<sup>1</sup> Tbsp water = 15 mL water

## PURAMINO A<sup>+®</sup> Per 1 Scoop

kcal De	sired	1 Scoop Powder‡ Added to Indicated Volume of Water,	Formula Yield, mL (fl oz)
per fl oz†	per mL	mL (fl oz)	ML (IL 02)
20	0.68	30 (1.0)	33 (1.1)
22	0.74	27 (0.9)	30 (1.0)
24	0.81	24 (0.8)	28 (0.9)
26	0.88	22 (0.7)	26 (0.9)
27	0.91	21 (0.7)	25 (0.8)
28	0.95	20 (0.7)	24 (0.8)
30	1.01	19 (0.6)	22 (0.8)

<sup>†</sup> Fluid ounce measures in above table are rounded to the nearest 0.1 fl oz.

Powders mix best when added on **top** of water. Then close the bottle and shake for about 5 seconds. The number of scoops and the amount of water to add can be doubled or tripled to make a larger volume.

One scoop of PURAMINO A+ $^{\circ}$  powder provides 22 kcal One scoop of powder displaces about 3 mL or 0.1 fl oz water One U.S. fl oz = 29.57 mL

## PURAMINO A+® JUNIOR Per 1 Scoop

kcal De	sired	1 Scoop Powder§ Added to Indicated Volume of Water,	Formula Yield, mL (fl oz)		
per fl oz†	per mL	mL (fl oz)	1112 (11 02)		
20	0.68	44 (1.5)	49 (1.7)		
22	0.74	40 (1.3)	45 (1.5)		
24	0.81	36 (1.2)	41 (1.4)		
26	0.88	33 (1.1)	38 (1.3)		
27	0.91	31 (1.1)	36 (1.2)		
28	0.95	30 (1.0)	35 (1.2)		
30	1.01	28 (1.0)	33 (1.1)		

<sup>†</sup> Fluid ounce measures in above table are rounded to the nearest 0.1 fl oz.

<sup>‡</sup> PURAMINO A+® should be measured with unpacked, level scoops.

<sup>1</sup> Tbsp water = 15 mL water

<sup>§</sup> PURAMINO A<sup>+®</sup> JUNIOR should be measured with **unpacked**, level scoops.

Powders mix best when added on top of water. Then close the bottle and shake for about 5 seconds. The number of scoops and the amount of water to add can be doubled or tripled to make a larger volume.

One scoop of PURAMINO A+® JUNIOR powder provides 33 kcal One U.S. fl oz = 29.57 mL

1 Tbsp water = 15 mL water

## Pregestimil® A+® Per 1 Scoop

#### 1 Scoop Powder† Added to kcal Desired Formula Yield, Indicated Volume of Water, mL (fl oz) mL (fl oz) per fl oz\* per mL 20 0.68 59 (2.0) 66 [2.2] 0.74 53 (1.8) 60 [2.0] 22 24 0.81 48 [1.6] 55 (1.9) 26 0.88 44 [1.5] 51 [1.7] 27 0.91 42 (1.4) 49 [1.6] 40 (1.4) 47 [1.6] 28 በ 95 30 1.01 37 (1.2) 44 (1.5)

Powders mix best when added on top of water. Then close the bottle and shake for about 5 seconds. The number of scoops and the amount of water to add can be doubled or tripled to make a larger volume.

One scoop of Pregestimil® A+® powder provides 45 kcal One scoop of powder displaces about 7 mL or 0.2 fl oz water One U.S. fl oz = 29.57 mL

1 Tbsp water = 15 mL water

<sup>\*</sup> Fluid ounce measures in above table are rounded to the nearest 0.1 fl oz.

<sup>†</sup> Pregestimil® A+® should be measured with packed, level scoops.

## **Scoop Information For Powder Products**

Powders	Scoop Weight (grams)	kcal/ scoop	Displacement (mL/g powder)	kcal/g	Protein (g/100 g)
Enfamil A+® NeuroPro™ Infant	8.8	44	0.77	5.1	11
Enfamil®	8.7	45	0.77	5.1	10.8
Enfamil A+®	8.8	44	0.77	5.1	10.1
Nutramigen® A+® with LGG®	9 <sup>+</sup>	45	0.77	5.0	13.9
Nutramigen® A+®	9 <sup>+</sup>	45	0.77	5.0	13.9
PURAMINO A <sup>+®</sup>	4.5	22	0.77	5.0	13.9‡
PURAMINO A <sup>+®</sup> JUNIOR	6.8	33	0.75	4.9	14 <sup>‡</sup>
Pregestimil® A <sup>+®</sup>	8.9 <sup>+</sup>	45	0.78	5.0	14
Enfamil A <sup>+®</sup> For feeding babies who frequently Spit Up	8.9	45	0.76	5.0	12.4
Enfamil A <sup>+®</sup> EnfaCare <sup>®</sup>	9.8	48	0.76	4.9	13.8
Enfamil A+® NeuroPro™ Gentlease®	8.7	45	0.77	5.1	12
Enfamil A+® Serenity	8.7	44	0.76	5.1	11.7
Enfamil® 2	8.9	45	0.72	5.0	13
Enfamil A+® 2	9	45	0.77	5.0	12.9
Enfagrow A+® Vanilla and Milk Flavour	12	53	0.74	4.4	17.8
Phenyl-Free® 1	4.5	23	0.76	5.0	16.2 <sup>‡</sup>
Phenyl-Free® 2	14.4	59	0.67	4.1	22 <sup>‡</sup>

LGG® is a registered trademark of Chr. Hansen A/S.





Powders	Scoop Weight (grams)	kcal/ scoop	Displacement (mL/g powder)	kcal/g	Protein (g/100 g)
Phenyl-Free® 2 HP	15.1	59	0.67	3.9	40 <sup>‡</sup>
BCAD 1	4.5	23	0.77	5.0	16.2‡
BCAD 2	14.5	59	0.67	4.1	24 <sup>‡</sup>
PFD 1	4.5	24	0.78	5.3	0
PFD 2	14.9	59	0.65	4.0	0
TYROS 1	4.5	23	0.77	5.0	16.7‡
TYROS 2	14.5	59	0.67	4.1	22 <sup>‡</sup>
Portagen®	9.4†	44	0.75	4.7	17

<sup>†</sup> Packed level measure - all others are unpacked level measures.

When accuracy is essential, gram measurements should be used. Note that scoop sizes do vary among the different products. Use scoop included with the product to ensure the desired caloric density is achieved.

Powdered formulas are not sterile. "To minimize the risk of infection, powdered formulas are not recommended for use in premature or immunocompromised infants unless no appropriate alternative is available, and then only with strict medical supervision and careful preparation, storage and use."

<sup>‡</sup> Protein equivalents.

International Formula Council. Infant Feeding: Safety Issues for Health Care Professionals. Atlanta, GA. International Formula Council; 2004:4.

## Concentrated Liquid Infant Formula Dilutions

These dilutions can be used with the following Mead Johnson concentrate formulas:

Enfamil A+®, Enfamil A+® Lactose Free, Enfamil A+® Soy and Enfamil A+® 2

kcal De	l Desired mL (fl oz) of water to add to 385 mL can of concentrated		Formula Yield, mL (fl oz)		
per fl oz	per mL	liquid	(		
20	0.68	384 (13)	769 (26)		
22	0.74	311 (10.5)	695 (23.5)		
24	0.81	251 (8.5)	636 (21.5)		
26	0.88	207 (7)	591 (20)		
27	0.91	177 (6)	562 (19)		
28	0.95	163 (5.5)	547 (18.5)		
30	1.01	133 (4.5)	518 (17.5)		

Each can of concentrated liquid provides 520 kcal.

30 mL of undiluted concentrated liquid provides 40 kcal.

1 U.S. fl oz = 29.57 mL

1 Tbsp water = 15 mL water

# Amino Acids Profile Infant Formula Amino Acid Compositions ESSENTIAL AMINO ACIDS

Product	Histidine	Isoleucine	Leucine	Lysine	Methionine
Enfamil A <sup>+®</sup> NeuroPro™ Infant mg/100 kcal mg/100 g*	53 280	120 630	220 1170	200 1020	53 280
Enfamil A <sup>+®</sup> NeuroPro™ Gentlease® mg/100 kcal mg/100 g <sup>†</sup>	60 310	131 680	240 1240	200 1030	51 (58 liq) 260
Enfamil A <sup>+®</sup> Concentrate mg/100 kcal mg/100 g	50 250	116 590	210 1080	180 910	46 230
Enfamil A+® For feeding babies who frequently Spit Up mg/100 kcal mg/100 g	73 350	130 620	250 1210	210 1010	63 300
Enfamil A <sup>+®</sup> EnfaCare <sup>®</sup> mg/100 kcal mg/100 g*	70 350	162 800	300 1480	250 1240	64 320
Pregestimil® A*® mg/100 kcal mg/100 g*	84 420	162 810	270 1370	240 1180	73 360
Nutramigen® A+® with LGG® mg/100 kcal mg/100 g*	84 420	162 810	270 1370	240 1180	73 360
Nutramigen® A+® mg/100 kcal mg/100 g*	84 420	162 810	270 1360	240 1170	73 360
PURAMINO A <sup>+®</sup> mg/100 kcal mg/100 g	73 360	190 950	340 1710	220 1110	64 320
PURAMINO A <sup>+®</sup> JUNIOR mg/100 kcal mg/100 g	73 360	190 950	340 1720	220 1120	64 320
Enfamil A+® 2 mg/100 kcal mg/100 g*	75 370	135 670	260 1300	220 1080	65 320
Enfagrow A*® mg/100 kcal mg/100 g*	116 520	210 930	400 1800	340 1500	100 450

LGG® is a registered trademark of Chr. Hansen A/S.

<sup>†</sup> At 20 Cal/fl oz (0.68 Cal/mL).



<sup>\*</sup> Powder only.

Product	Cysteine	Phenyl- alanine	Tyrosine	Threonine	Tryptophan	Valine
Enfamil A+® NeuroPro™ Infant mg/100 kcal mg/100 g*	38 200	90 470	67 (80 liq) 350	124 650	29 153	130 680
Enfamil A⁺® NeuroPro™ Gentlease® mg/100 kcal mg/100 g⁺	32 (34 liq) 168	97 500	83 (87 liq) 430	133 700	29 (30 liq) 149	147 770
Enfamil A <sup>+®</sup> Concentrate mg/100 kcal	36	84	80	122	35	122
Enfamil A <sup>+®</sup> For feeding babies who frequently Spit Up mg/100 kcal mg/100 g	23 108	125 600	110 530	115 550	33 156	158 760
Enfamil A <sup>+®</sup> EnfaCare <sup>®</sup> mg/100 kcal mg/100 g*	51 250	118 580	112 550	171 840	49 240	171 840
Pregestimil® A+® mg/100 kcal mg/100 g*	48 240	137 690	59 290	129 640	46 230	200 1010
Nutramigen® A+® with LGG® mg/100 kcal mg/100 g*	48 240	137 690	59 290	129 640	46 230	200 1010
Nutramigen® A+® mg/100 kcal mg/100 g*	48 240	137 680	59 290	129 640	46 230	200 1000
PURAMINO A <sup>+®</sup> mg/100 kcal mg/100 g	73 360	140 700	151 750	171 850	67 330	210 1060
PURAMINO A+® JUNIOR mg/100 kcal mg/100 g	73 360	140 700	151 760	171 850	67 340	210 1060
Enfamil A <sup>+®</sup> 2 mg/100 kcal mg/100 g*	23 116	130 650	130 650	120 590	37 182	164 810
Enfagrow A+® mg/100 kcal mg/100 g*	36 160	200 890	176 780	184 820	52 230	250 1120

 $LGG^{\otimes}$  is a registered trademark of Chr. Hansen A/S. \* Powder only. † At 20 Cal/fl oz [0.68 Cal/mL].

## **NONESSENTIAL AMINO ACIDS**

Product	Arginine	Alanine	Aspartic Acid	Glutamic Acid
Enfamil A <sup>+®</sup> NeuroPro™ Infant mg/100 kcal mg/100 g*	71 370	101 530	200 1030	400 2100
Enfamil A⁺® NeuroPro™ Gentlease® mg/100 kcal mg/100 g†	81 420	106 550	220 1130	460 2400
Enfamil A <sup>+®</sup> Concentrate mg/100 kcal	60	88	200	400
Enfamil A <sup>+®</sup> For feeding babies who frequently Spit Up mg/100 kcal mg/100 g	88 420	85 410	200 960	550 2600
Enfamil A+® EnfaCare® mg/100 kcal mg/100 g*	84 410	123 610	270 1350	560 2800
Pregestimil® A+® mg/100 kcal mg/100 g*	112 560	95 480	210 1060	640 3200
Nutramigen® A+® with LGG® mg/100 kcal mg/100 g*	112 560	95 480	210 1060	640 3200
Nutramigen® A*® mg/100 kcal mg/100 g*	112 560	95 470	210 1060	640 3200
PURAMINO A⁺® mg/100 kcal mg/100 g	148 740	270 1360	560 2800	168 840
PURAMINO A+® JUNIOR mg/100 kcal mg/100 g	148 740	270 1360	560 2800	168 840
Enfamil A <sup>+®</sup> 2 mg/100 kcal mg/100 g*	91 450	88 440	210 1030	570 2800
Enfagrow A <sup>+®</sup> mg/100 kcal mg/100 g*	140 620	136 610	320 1420	880 3900

LGG° is a registered trademark of Chr. Hansen A/S. \* Powder only.  $\dagger$  At 20 Cal/fl oz (0.68 Cal/mL).

Product	Glycine	Proline	Serine
Enfamil A <sup>+®</sup> NeuroPro™ Infant mg/100 kcal mg/100 g*	42 220	162 850	116 610
Enfamil A⁺® NeuroPro™ Gentlease® mg/100 kcal mg/100 g†	45 230	189 980	127 660
Enfamil A <sup>+®</sup> Concentrate mg/100 kcal	40	160	114
Enfamil A*® For feeding babies who frequently Spit Up mg/100 kcal mg/100 g	50 240	250 1190	145 700
Enfamil A+® EnfaCare® mg/100 kcal mg/100 g*	56 280	220 1100	160 790
Pregestimil <sup>®</sup> A* <sup>®</sup> mg/100 kcal mg/100 g*	59 290	290 1470	162 810
Nutramigen® A⁺® with LGG® mg/100 kcal mg/100 g*	59 290	290 1470	162 810
Nutramigen® A+® mg/100 kcal mg/100 g*	59 290	290 1460	162 810
PURAMINO A <sup>+®</sup> mg/100 kcal mg/100 g	73 360	280 1390	171 850
PURAMINO A+® JUNIOR mg/100 kcal mg/100 g	73 360	280 1400	171 850
Enfamil A+® 2 mg/100 kcal mg/100 g*	52 260	260 1280	151 750
Enfagrow A <sup>+®</sup> mg/100 kcal mg/100 g*	80 360	400 1760	230 1030

 $LGG^{\odot}$  is a registered trademark of Chr. Hansen A/S. \* Powder only. † At 20 Cal/fl oz (0.68 Cal/mL).

## Metabolic Formula Amino Acid Compositions ESSENTIAL AMINO ACIDS

Product	Histidine	Isoleucine	Leucine	Lysine	Methionine
BCAD 1 mg/100 kcal mg/100 g	109 550	0	0	350 1770	93 470
BCAD 2 mg/100 kcal mg/100 g	220 890	0	0	700 2900	189 770
PFD 1 mg/100 kcal mg/100 g	0	0	0	0	0
PFD 2 mg/100 kcal mg/100 g	0	0	0	0	0
Phenyl-Free® 1 mg/100 kcal mg/100 g	80 410	230 1150	420 2100	260 1300	70 360
Phenyl-Free® 2 mg/100 kcal mg/100 g	135 550	380 1560	700 2900	430 1760	119 480
Phenyl-Free® 2 HP mg/100 kcal mg/100 g	260 1000	720 2800	1330 5200	820 3200	220 880
TYROS 1 mg/100 kcal mg/100 g	83 420	240 1200	440 2200	270 1370	73 370
TYROS 2 mg/100 kcal mg/100 g	135 550	390 1580	710 2900	440 1800	119 480

Product	Cysteine	Phenyl- alanine	Tyrosine	Threonine	Tryptophan	Valine
BCAD 1 mg/100 kcal mg/100 g	67 340	182 920	182 920	150 760	80 410	0
BCAD 2 mg/100 kcal mg/100 g	136 550	370 1490	370 1490	300 1220	159 650	0 0

Product	Cysteine	Phenyl- alanine	Tyrosine	Threonine	Tryptophan	Valine
PFD 1 mg/100 kcal mg/100 g	0	0	0	0	0	0
PFD 2 mg/100 kcal mg/100 g	0	0	0	0	0 0	0
Phenyl-Free® 1 mg/100 kcal mg/100 g	48 240	0	320 1600	147 750	58 290	250 1250
Phenyl-Free® 2 mg/100 kcal mg/100 g	81 330	0	530 2200	250 1010	97 400	420 1690
Phenyl-Free® 2 HP mg/100 kcal mg/100 g	153 600	0	1010 4000	470 1840	184 720	790 3100
TYROS 1 mg/100 kcal mg/100 g	53 270	0	0	155 780	63 320	260 1320
TYROS 2 mg/100 kcal mg/100 g	86 350	0	0	250 1030	103 420	430 1740

## **NONESSENTIAL AMINO ACIDS**

Product	Arginine	Alanine	Aspartic Acid	Glutamic Acid
BCAD 1 mg/100 kcal mg/100 g	230 1180	250 1280	310 1560	610 3100
BCAD 2 mg/100 kcal mg/100 g	420 1730	460 1870	550 2300	1100 4500
PFD 1 mg/100 kcal mg/100 g	0	0	0	0
PFD 2 mg/100 kcal mg/100 g	0	0	0	0
Phenyl-Free® 1 mg/100 kcal mg/100 g	192 970	210 1070	250 1280	510 2600
Phenyl-Free® 2 mg/100 kcal mg/100 g	320 1320	360 1450	430 1740	850 3500
Phenyl-Free® 2 HP mg/100 kcal mg/100 g	610 2400	670 2600	810 3200	1610 6300
TYROS 1 mg/100 kcal mg/100 g	220 1100	240 1200	290 1450	570 2900
TYROS 2 mg/100 kcal mg/100 g	360 1450	390 1580	470 1910	930 3800

Product	Glycine	Proline	Serine
BCAD 1 mg/100 kcal mg/100 g	141 710	310 1560	154 780
BCAD 2 mg/100 kcal mg/100 g	250 1030	550 2300	280 1130
PFD 1 mg/100 kcal mg/100 g	0	0	0
PFD 2 mg/100 kcal mg/100 g	0	0	0
Phenyl-Free® 1 mg/100 kcal mg/100 g	115 580	250 1280	125 630
Phenyl-Free® 2 mg/100 kcal mg/100 g	194 790	430 1740	210 860
Phenyl-Free® 2 HP mg/100 kcal mg/100 g	370 1440	810 3200	400 1560
TYROS 1 mg/100 kcal mg/100 g	129 650	290 1450	142 720
TYROS 2 mg/100 kcal mg/100 g	210 860	470 1910	230 950

## Portagen® Amino Acid Composition ESSENTIAL AMINO ACIDS

Product	Histidine	Isoleucine	Leucine	Lysine	Methionine
mg/100 kcal	104	184	350	290	101
mg/100 g	490	870	1630	1380	480

Product	Cysteine	Phenyl- alanine	Tyrosine	Threonine	Tryptophan	Valine
mg/100 kcal	15.5	191	210	158	56	240
mg/100 g	73	900	990	750	270	1120

## **NONESSENTIAL AMINO ACIDS**

Product	Arginine	Alanine	Aspartic Acid	Glutamic Acid
mg/100 kcal	133	115	260	830
mg/100 g	630	540	1240	3900

Product	Glycine	Proline	Serine
mg/100 kcal	68	390	210
mg/100 g	320	1850	1000

## Premature Formula Amino Acid Compositions ESSENTIAL AMINO ACIDS

Product	Histidine	Isoleucine	Leucine	Lysine	Methionine
Enfamil A+® Premature 20 kcal (59 mL) mg/100 kcal mg/100 mL	75 50	174 116	320 210	270 180	69 46
Enfamil A+® Premature 24 kcal (59 mL) mg/100 kcal mg/100 mL	75 60	174 139	320 260	270 220	69 55
Enfamil A <sup>+®</sup> Premature High Protein (59 mL) mg/100 kcal mg/100 mL	88 70	200 162	370 300	320 250	81 64
Enfamil A <sup>+®</sup> EnfaCare <sup>®</sup> (59 mL) mg/100 kcal mg/100 mL	70 53	162 122	300 220	250 189	64 48

Product	Cysteine	Phenyl- alanine	Tyrosine	Threonine	Tryptophan	Valine
Enfamil A+® Premature 20 kcal (59 mL) mg/100 kcal mg/100 mL	55 36	126 84	120 80	183 122	53 35	183 122
Enfamil A+® Premature 24 kcal (59 mL) mg/100 kcal mg/100 mL	55 44	126 101	120 96	183 146	53 42	183 146

Product	Cysteine	Phenyl- alanine	Tyrosine	Threonine	Tryptophan	Valine
Enfamil A+® Premature High Protein (59 mL) mg/100 kcal mg/100 mL	64 51	147 118	140 112	210 171	61 49	210 171
Enfamil A <sup>+®</sup> EnfaCare <sup>®</sup> (59 mL) mg/100 kcal mg/100 mL	51 38	118 88	112 84	171 128	49 37	171 128

## **NONESSENTIAL AMINO ACIDS**

Product	Arginine	Alanine	Aspartic Acid	Glutamic Acid
Enfamil A <sup>+®</sup> Premature 20 kcal (59 mL) mg/100 kcal mg/100 mL	90 60	132 88	290 200	600 400
Enfamil A+® Premature 24 kcal (59 mL) mg/100 kcal mg/100 mL	90 72	132 106	290 240	600 480
Enfamil A <sup>+®</sup> Premature High Protein (59 mL) mg/100 kcal mg/100 mL	105 84	154 123	340 270	700 560
Enfamil A <sup>+®</sup> EnfaCare <sup>®</sup> (59 mL) mg/100 kcal mg/100 mL	84 63	123 92	270 210	560 420

Product	Glycine	Proline	Serine
Enfamil A <sup>+®</sup> Premature 20 kcal (59 mL) mg/100 kcal mg/100 mL	60 40	240 160	171 114
Enfamil A <sup>+®</sup> Premature 24 kcal (59 mL) mg/100 kcal mg/100 mL	60 48	240 192	171 137
Enfamil A <sup>+®</sup> Premature High Protein (59 mL) mg/100 kcal mg/100 mL	70 56	280 220	200 160
Enfamil A <sup>+®</sup> EnfaCare <sup>®</sup> (59 mL) mg/100 kcal mg/100 mL	56 42	220 168	160 120

## Human Milk Fortifiers Amino Acid Compositions

## **ESSENTIAL AMINO ACIDS**

Product	Histidine	Isoleucine	Leucine	Lysine	Methionine
Enfamil® Human Milk Fortifier Powder Per 1 packet (0.71g) mg	7.3	15.4	34	28	7
Enfamil® Liquid Human Milk Fortifier Standard Protein (Without Preterm Human Milk) Per 5 mL mg	9.2	25	59	48	10.1
Enfamil® Liquid Human Milk Fortifier High Protein (Without Preterm Human Milk) Per 5 mL mg	12.3	33	78	64	13.4

Product	Cysteine	Phenyl- alanine	Tyrosine	Threonine	Tryptophan	Valine
Enfamil® Human Milk Fortifier Powder Per 1 packet (0.71 g) mg	6.4	12.3	12.6	14	5.6	16.2
Enfamil® Liquid Human Milk Fortifier Standard Protein (Without Preterm Human Milk) Per 5 mL	15.1	16	16.8	23	10.5	23
Enfamil® Liquid Human Milk Fortifier High Protein (Without Preterm Human Milk) Per 5 mL	20	21	22	30	14	30

## **NONESSENTIAL AMINO ACIDS**

Product	Arginine	Alanine	Aspartic Acid	Glutamic Acid
Enfamil® Human Milk Fortifier Powder Per 1 packet (0.71 g) mg	8.4	12.6	29	56
Enfamil® Liquid Human Milk Fortifier Standard Protein (Without Preterm Human Milk) Per 5 mL mg	10.5	24	53	76
Enfamil® Liquid Human Milk Fortifier High Protein (Without Preterm Human Milk) Per 5 mL mg	14	31	71	101

Product	Glycine	Proline	Serine
Enfamil® Human Milk Fortifier Powder Per 1 packet (0.71 g) mg	5.3	20	14
Enfamil® Liquid Human Milk Fortifier Standard Protein (Without Preterm Human Milk) Per 5 mL mg	7.7	18.9	18.1
Enfamil® Liquid Human Milk Fortifier High Protein (Without Preterm Human Milk) Per 5 mL mg	10.3	25	24

## Kosher Enfamil® Products

Kosher means the food or product has been prepared according to the requirements of Jewish Law.

Some of our products are manufactured under the supervision of the Kashruth Division of the Union of Orthodox Jewish Congregations of America.

O.U. on a product signifies it is kosher and is authorized by the Orthodox Union.

Our kosher-certified products will have a "U" within a circle on the product label.



#### Pareve

Pareve is a Hebrew term that describes food without any meat or dairy ingredients.

## Halal Enfamil® Products

Halal food means food permitted under Islamic Law. It encompasses the food source, how animals are slaughtered, and the preparation, processing, packaging, transportation, and storage of foods. The foremost unlawful foods under Islamic Law are pork products and alcoholic beverages and ingredients derived from these items.

The organization that certifies our products halal is the Islamic Food and Nutrition Council of America or IFANCA®.

#### Alcohol

We do not add alcohol to any of our infant formulas, toddler drinks or medical foods.

Synthetic alcohol is used during the drying process of the nucleotides in our infant formulas; however, the amount of alcohol remaining in the nucleotides is very small (200 ppm [parts per million] or less).

#### Nucleotides

The nucleotides we use in our formulas are certified Halal by the Islamic Food and Nutrition Council of America and Kosher by the Orthodox Union.

The nucleotide ingredients may be certified Halal, but our products are not Halal-certified at this time, with the exception of PURAMINO  $A^{+ \circ}$  and PURAMINO  $A^{+ \circ}$  JUNIOR.

#### Pork

The casein protein in our Nutramigen®  $A^{+}$ ®, Nutramigen®  $A^{+}$ ® with LGG® and Pregestimil®  $A^{+}$ ® are hydrolyzed or broken down by an enzyme process using pork pancreatic enzyme. Very little of the enzyme remains in the final product formulation; in fact, pork contributes an insignificant percentage of the prepared formula.

Portfolio Category	SKU Number	Product Name	Halal Status	Kosher Status
	3221214	Enfamil® Human Milk Fortifier (Powder)	Halal Certified	Kosher Dairy
	3279585	Enfamil® Liquid Human Milk Fortifier Standard Protein 163 mL Flex Pro Bottle (6x4 Bottles Per Case)	Halal Certified	Kosher Dairy
	3279712	Enfamil® Liquid Human Milk Fortifier High Protein 163 mL Flex Pro Bottle (6x4 Bottles Per Case)	Halal Certified	Kosher Dairy
	3180545	Enfamil® Liquid Human Milk Fortifier Standard Protein 5 mL sachet	Halal Certified	Kosher Dairy
Premature	3180315	Enfamil® Liquid Human Milk Fortifier High Protein 5 mL sachet	Halal Certified	Kosher Dairy
	1250747	Enfamil A+® Premature 20 kcal Nursette® Bottle	Halal Certified	Kosher Dairy
	1215248	Enfamil A+® Premature 24 kcal Nursette® Bottle	Halal Certified	Kosher Dairy
	1281250	Enfamil A+® Premature High Protein Nursette® Bottle	Halal Certified	Kosher Dairy
	1171320	Enfamil A+® EnfaCare® Powder	Halal Certified	Kosher Dairy
	1215387	Enfamil A+® EnfaCare® Nursette® Bottle	Halal Certified	Kosher Dairy
	1210915	Enfamil A+® Hypercaloric	Halal Certified	Kosher Dairy
	1215250/ 3130303	Enfamil® Water	Halal Certified	Kosher Pareve (Manufactured on dairy equipment)
	1216351	Enfamil® 5% Glucose in Water	Halal Certified	Kosher Pareve (Manufactured on dairy equipment)
Term & Specialty	1271716	Enfamil A+® Soy Nursette® Bottle	Halal Certified	Kosher Pareve
	3276942	Enfamil A+® Soy Concentrate	Halal Certified	Kosher Pareve
	1271623	Enfamil A+® Lactose Free Nursette® Bottle	Halal Certified	Kosher Dairy
	3278010	Enfamil A+® Lactose Free Concentrate	Halal Certified	Kosher Dairy

Portfolio Category	SKU Number	Product Name	Halal Status	Kosher Status
	3290179	Enfamil A+® NeuroPro™ Nursette® Bottle (Only Hospital)	Halal Certified	Kosher Dairy
	3276799	Enfamil A+® NeuroPro™ Powder	Halal Certified	Kosher Dairy
	3285831	Enfamil A+® NeuroPro™ 237 mL RTF (24 bottles per case)	Halal Certified	Kosher Dairy
	3285832	Enfamil A+® NeuroPro™ 237 mL RTF (18 bottles per case)	Halal Certified	Kosher Dairy
	3236031	Enfamil A+® NeuroPro™ Nursette® Bottle (Only Hospital)	Halal Certified	Kosher Dairy
	3290179	Enfamil A+® NeuroPro™ Hospital Nursette® (48 per case)	Halal Certified	Kosher Dairy
	3276799	Enfamil A+® NeuroPro™ Powder - 587 gms (6 tubs per case)	Halal Certified	Kosher Dairy
	3285832	Enfamil A+® NeuroPro™ Ready to Use (RTU) (18 bottles per case)	Halal Certified	Kosher Dairy
Term & Specialty	3285831	Enfamil A+® NeuroPro™ Ready to Use (RTU) (24 bottles per case)	Halal Certified	Kosher Dairy
	3286052	Enfamil A+® NeuroPro™ Gentlease® Powder – 553 gms (6 tubs per case)	Halal Certified	Kosher Dairy
	3276297	Enfamil A+® NeuroPro™ Gentlease® Ready to Use (RTU) (24 bottles per case)	Halal Certified	Kosher Dairy
	3232162	Enfamil A+® Concentrate	Halal Certified	Kosher Dairy
	1233351	Puramino A+® Powder	Halal Certified	Kosher Dairy
	2021478	Enfamil® Enfalyte® Oral Electrolyte Solution	Halal Certified	Kosher Pareve (Manufactured on dairy equipment)
	1239201	Nutramigen® A+® Powder	_	Not Kosher
	3230626	Nutramigen® A+® with LGG®	_	Not Kosher
	3229971	Nutramigen® A+® Nursette® Bottle	_	Not Kosher
	1238161	Pregestimil® A+® Powder	-	Not Kosher

 $LGG^{\circledcirc}$  is a registered trademark of Chr. Hansen A/S.

Portfolio Category	SKU Number	Product Name	Halal Status	Kosher Status
	1273987	Pregestimil® A+® Nursette® Bottle	_	Not Kosher
Term & Specialty	3271466	Enfamil A+® For feeding babies who frequently Spit Up*	Halal Certified	Kosher Dairy
	Not provided to hospitals	Enfamil A+® Serenity	Halal Certified	Kosher Dairy
	2047834	Puramino A+® JUNIOR	Halal Certified	Kosher Dairy
	3181752	Enfagrow A+® Toddler Nutritional Drink, Vanilla Flavour Powder	Halal Certified	Kosher Dairy
Toddler	3181749	Enfagrow A+® Toddler Nutritional Drink, Milk Flavour Powder	Halal Certified	Kosher Dairy
roducer	2053715	Enfagrow A+® Toddler Nutritional Drink, Vanilla Flavour Ready to Drink Bottles	Halal Certified	Kosher Dairy
	2053714	Enfagrow A+® Toddler Nutritional Drink, Milk Flavour Ready to Drink Bottles	Halal Certified	Kosher Dairy
	2032499	Portagen®	Halal Certified	Kosher Dairy
	2001922	Phenyl-Free® 1 (Powder)	Halal Certified	Kosher Dairy
	1281334	Phenyl-Free® 2 HP (Powder)	Halal Certified	Kosher Dairy
	1281335	Phenyl-Free® 2 (Powder)	Halal Certified	Kosher Dairy
Metabolics	2001957	TYROS 1 (Powder)	Halal Certified	Kosher Dairy
	1281396	TYROS 2 (Powder)	Halal Certified	Kosher Dairy
	2001952	BCAD 1	Halal Certified	Kosher Dairy
	1281386	BCAD 2	Halal Certified	Kosher Dairy
	3144090	PFD 1	Halal Certified	Kosher Dairy
	1281389	PFD 2	Halal Certified	Kosher Dairy
Vitamins	1015277	Enfamil® D-Vi-Sol® (Liquid)	_	Kosher Dairy
* (tallilli)	1015245	Enfamil® Tri-Vi-Sol® (Liquid)	-	Kosher Dairy

 $<sup>^{\</sup>star}$  For babies who spit up more than 4 times per day.

Portfolio Category	SKU Number	Product Name	Halal Status	Kosher Status
	1015280	Enfamil® Poly-Vi-Sol® (Liquid)	_	Kosher Dairy
Vitamins 101515	1015154	Enfamil® Fer-In-Sol® (Liquid)	_	Kosher Dairy
	1015278	Enfamil® Fer-In-Sol® (Syrup)	_	Kosher Dairy

## **Hospital Instructions**

## FOR SAFE INFANT FORMULA PREPARATION, STORAGE, AND USE.

Powdered formulas are not sterile. "To minimize the risk of infection, powdered formulas are not recommended for use in premature or immunocompromised infants unless no appropriate nutritional alternative is available, and then only with strict medical supervision and careful preparation, storage and use."

### Preparation and Use Instructions<sup>1,2</sup>

- Follow hospital procedures carefully
- Check label against individual patient's feeding order (confirm product identity and dilution)
- Wash and dry hands and clean work area thoroughly with an antibacterial sanitizing solution that is appropriate for food contact surfaces
- Ensure container integrity
- Use only sterilized containers and measuring equipment
- Avoid touch contamination of formula at each step
- Visually inspect the formula (if anything looks unusual, set it aside and contact your Mead Johnson representative)
- When preparing powdered or concentrated liquid formulas, chilled, sterilized water is recommended
- Do not use a blender to reconstitute powdered formula
- Do not store in formula warmer or use microwave oven to warm formula

### Storage Instructions for Prepared Formula

### Refrigeration

- Dedicated refrigerators recommended
- Store at 2-4°C (35-40°F) no longer than 24 hours

### Room Temperature

- Should be fed immediately, but hold no longer than a total of 2 hours before feeding
- If bottle is warmed, discard after 1 hour

#### After Feeding Begins

- · Feed within 1 hour or discard
- Do not refrigerate for later feedings



Tube-feeding Hangtimes*† for Prepared Formula From:	Neonates or Immunocompromised Infants/Children	Infants and Children with Healthy Immune Systems
Ready-To-Feed unaltered infant formula (commercially sterile)	4 hrs	8 hrs
Concentrated liquid formulas (commercially sterile) Powdered formulas (not sterile)	4 hrs	4 hrs
Powder added to liquid formula or expressed breast milk <sup>3</sup>	4 hrs	4 hrs
Liquid (commercially sterile) added to expressed breast milk	4 hrs	4 hrs

#### Failure to follow these instructions could result in severe harm.

- International Formula Council. Infant Feeding: Safety Issues for Health Care Professionals. Atlanta, GA: International Formula Council; 2004:4.
- 2. Robbins ST, Meyers R, eds. Infant Feedings: Guidelines for Preparation of Human Milk and Formula in Health Care Facilities. Chicago, Ill: American Dietetic Association; 2011.
- Telang S, Berseth CL, Ferguson PW, et al. Fortifying fresh human milk with commercial powdered human milk fortifiers does not affect bacterial growth during 6 hours at room temperature. J Am Diet Assoc. 2005;105:1567-1572.

The American Dietetic Association is now known as the Academy of Nutrition and Dietetics.

<sup>\*</sup> Adapted from: Robbins and Meyers, p.100.2

<sup>&</sup>lt;sup>†</sup> For reservoir and tube-change guidelines, refer to the ADA publication, Infant Feedings: Guidelines for Preparation of Formula and Breastmilk in Health Care Facilities. 2004.

## **Miscellaneous Nutrient Information**

#### **Calculations and Conversions**

Dietary Reference Intakes - Unit Conversion Factors

Folate	1 DFE = 1 µg food folate		
Niacin	1 NE = 1 mg niacin 1mg NE = 60 mg tryptophan		
Vitamin A	1 RAE = 1 μg retinol = 3.33 IU retinol For preformed vitamin A, 1 RE = 1RAE		
Vitamin D	1 μg = 40 IU		
Vitamin E	1 IU natural or synthetic Vitamin E = 0.67 mg $\alpha$ -tocopherol equivalent ( $\alpha$ -TE) 1 IU synthetic Vitamin E and esters = 0.45 mg $\alpha$ -tocopherol 1 IU natural Vitamin E and esters = 0.67 mg $\alpha$ -tocopherol		



Date published 11/2024 – Please reference Mead Johnson Nutrition's online Product Handbook for the most up to date product details (www.enfamil.ca/medical).

## **Alberta Reimbursement List**

The following Mead Johnson products are listed on the Alberta Drug Benefit Supplement.

PIN	Mead Johnson Product Code	Product	Format	Benefit Category
00999817	2015524	Nutramigen® A+® with LGG®	561 g container	Regular Benefit
00999520	1239201	Nutramigen® A+® Powder	454 g container	Regular Benefit
00999543	1233351	Puramino A+® Powder	400 g container	Special Authorization
00999876	2047834	Puramino A+® Junior Powder	400 g container	Special Authorization
00999564	1171320	Enfamil A+® EnfaCare® Powder	363 g container	Regular Benefit
00762881	1015277	Enfamil® D-Vi-Sol® Liquid	50 mL bottle	Restricted Benefit
00762954	1015154	Enfamil® Fer-In-Sol® Liquid	50 mL bottle	Restricted Benefit
00017884	1015278	Enfamil® Fer-In-Sol® Syrup	50 mL bottle	Restricted Benefit
00762946	1015280	Enfamil® Poly-Vi-Sol® Liquid	50 mL bottle	Restricted Benefit
00762903	1015245	Enfamil® Tri-Vi-Sol® 50 mL bottle		Restricted Benefit

LGG® is a registered trademark of Chr. Hansen A/S.

For more information, please visit:

https://www.alberta.ca/drug-benefit-list-and-drug-review-process.aspx





## **Ontario Reimbursement List**

The following Mead Johnson products are listed on the Ontario Drug Benefit Formulary / Comparative Drug Index.

PIN	Mead Johnson Product Code	Product	Format	Benefit Category
09857565	2015524	Nutramigen® A+® with LGG®	561 g container	Nutrition Product
09857345	1239201	Nutramigen® A+® Powder	454 g container	Nutrition Product
09857369	1233351	Puramino A <sup>+®</sup> Powder	400 g container	Nutrition Product
09854401	2032499	Portagen® Powder	410 g container	Nutrition Product
09857172	1171320	Enfamil A+® EnfaCare® Powder	363 g container	Nutrition Product
00762954	1015154	Enfamil® Fer-In-Sol® Liquid	50 mL bottle	General Benefit

LGG® is a registered trademark of Chr. Hansen A/S.

For more information, please visit:

https://www.health.gov.on.ca/en/public/programs/drugs/programs/odb/opdp\_ nutrition.aspx

## **Quebec Reimbursement List**

The following Mead Johnson products are listed on the Régie de l'assurance maladie du Québec (RAMQ) List of Medications.

PIN	Mead Johnson Product Code	Product	Format	Benefit Category
99101338	2015524	Nutramigen® A+® with LGG®	561 g container	Nutritional Formula - Casein Hydrolysate (Infants and Children)
99100532	1239201	Nutramigen® A+® Powder	454 g container	Nutritional Formula - Casein Hydrolysate (Infants and Children)
99100533	1238161	Pregestimil® A+® Powder	454 g container	Nutritional Formula - Casein Hydrolysate (Infants and Children)
99100715	1233351	Puramino A <sup>+®</sup> Powder	400 g container	Nutritional Formula - Monomeric With Iron (Infants Or Children)
99101278	2047834	Puramino A <sup>+®</sup> Junior Powder	400 g container	Nutritional Formula - Monomeric With Iron (Infants or Children)
00881201	2032499	Portagen® Powder	410 g container	Nutritional Formula - Skim Milk / Coconut Oil
99100122	1171320	Enfamil A <sup>+®</sup> EnfaCare® Powder	363 g container	Nutritional Formula - Post- Discharge Preterm Formula (Infants)
00762881	1015277	Enfamil® D-Vi-Sol® Liquid	50 mL bottle	Vitamin D
00762954	1015154	Enfamil® Fer-In-Sol® Liquid	50 mL bottle	Iron Preparations
00017884	1015278	Enfamil® Fer-In-Sol® Syrup	250 mL bottle	Iron Preparations
00762903	1015245	Enfamil® Tri-Vi-Sol® Liquid	50 mL bottle	Multivitamins

LGG® is a registered trademark of Chr. Hansen A/S.

For more information, please visit:

https://www.ramq.gouv.qc.ca/en/about-us/list-medications





## Saskatchewan Reimbursement List

The following Mead Johnson products are listed by the Saskatchewan Aids to Independent Living Program and Therapeutic Nutritional Products Program.

Mead Johnson Product Code	Product	Format	Benefit Category
1171320	Enfamil A+® EnfaCare® Powder	363 g container	Eligible Formula
1239201	Nutramigen® A+® Powder	454 g container	Eligible Formula
2015524	Nutramigen® A+® with LGG®	561 g container	Eligible Formula
2032499	Portagen® Powder	410 g container	Eligible Formula
1238161	Pregestimil® A+® Powder	454 g container	Eligible Formula
2047834	Puramino A+® Junior Powder	400 g container	Eligible Formula
1233351	Puramino A <sup>+®</sup> Powder	400 g container	Eligible Formula

LGG® is a registered trademark of Chr. Hansen A/S.

For more information, please visit:

https://www.saskatchewan.ca/residents/health/accessing-health-care-services/ health-services-for-people-with-disabilities/sail



# Healthcare Professional Resource Centre

**Your Partner in Pediatric Nutrition** 

### enfamil.ca/medical

The Mead Johnson Nutrition Healthcare Professional Resource Centre provides information, support tools, and professional education.

Date published 11/2024 – Please reference Mead Johnson Nutrition's online Product Handbook for the most up to date product details (www.enfamil.ca/medical).

